

ZR

ISSN 2095-8137 CN 53-1229/Q

Volume **38** Issue **1**
18 January **2017**

Zoological Research



CODEN: DOYADI
www.zoores.ac.cn

ZOOLOGICAL RESEARCH

Volume 38, Issue 1 18 January 2017

CONTENTS

Editorial

New Year Address of *Zoological Research* (1)

Obituary

In Memory of Academician Er-Mi Zhao (1930-2016) (2)

Research highlight

Engrafted newborn neurons could functionally integrate into the host neuronal network
..... *Zheng-Bo Wang, Dong-Dong Qin, Xin-Tian Hu* (5)

Reviews

Consequences of early adverse rearing experience (EARE) on development: insights from non-human
primate studies *Bo Zhang* (7)

Does mRNA structure contain genetic information for regulating co-translational protein folding?
..... *Jian-Rong Yang* (36)

Reports

Expression of plgR in the tracheal mucosa of SHIV/SIV-infected rhesus macaques
..... *Dong Li, Feng-Jie Wang, Lei Yu, Wen-Rong Yao, Yan-Fang Cui, Gui-Bo Yang* (44)

Nest survival rate of Reeves's pheasant (*Syrnaticus reevesii*) based on artificial nest experiments
..... *Xu Luo, Yu-Ze Zhao, Jing Ma, Jian-Qiang Li, Ji-Liang Xu* (49)

Cover image: *Macaca mulatta*. Photo by Xiao-Feng Ma

Cover design: Lin Lei

New Year Address of *Zoological Research*

The year 2016 was a wonderful and important one for *Zoological Research* (ZR). In recognition of its impressive progress and great potential to develop into a leading journal in the field, ZR is now supported by the “Project for Enhancing International Impact of China STM Journals” (PIIJ) (Class B) (2016–2018). Having wide coverage, PIJ is the most influential project in China for supporting and promoting the development of English language academic journal publications. In addition, ZR has achieved the “International Impact Academic Journal of China” title five years in a row.

The success of ZR depends on its unique niche in the field and the joint effort of its strong editorial team. Emphasizing the scope of ZR, namely, “Primates and Animal Models”, “Conservation & Utilization of Animal Resources”, and “Animal Diversity and Evolution”, two special issues were released in 2016: “*The Amphibian and Reptile Biodiversity of Qinghai-Tibetan Plateau*” and “*CRISPR/Cas9, Model Animals and Human Diseases*”. Moreover, to promote communication and collaboration among peer researchers, ZR is hosting the “2017 Frontiers in Zoology Symposium” in February 2017.

To further strengthen the editorial board and widen its scope of expertise, ZR continues to recruit local and overseas scientists to serve on its board. In 2016, ZR engaged four renowned scientists with varied expertise to join the vibrant ZR editorial team. These new board members include: Dr. Wai-Yee Chan, Chair Professor and Director, School of Biomedical Sciences, Chinese Hong Kong University, Hong Kong SAR, China; Dr. LeAnn Blomberg, Beltsville Agricultural Research Center (BARC), United States Department of Agriculture, USA; Dr. Hua-Xin (Larry) Liao, Professor, College of Life Science and Technology, Jinan University, China, and Professor, Duke University, USA; and Dr. Meng-Ji Lu, Professor, Institute of Virology, University of Duisburg-Essen, Germany. In addition, Dr. Guojie Zhang, a well-known expert on animal genomics who holds a position at the China National GeneBank, BGI-Shenzhen, and the Department of Biology, University of Copenhagen, as well as a guest professorship at the Kunming Institute of Zoology, Chinese Academy of Sciences, will also join the editorial board. With these new members, the editorial board of ZR now includes 48 distinguished scientists with expertise across multiple disciplines.

To further its scientific reach, ZR has continued to promote

and market the journal during this past year. A new and more interactive homepage was launched and, as an open access journal, full-text articles can be downloaded via the homepage, PubMed Central, and SciEngine. To expand our readership, abstracts of published articles are also available online in Chinese. Information regarding the new initiatives and measures of ZR will continue to be shared with our readers through our official WeChat account (ZoolRes). In addition, ZR has now been indexed by Scopus and other related databases, and has a Citescore value of 0.48 and ranks 786 among 1 549 journals in the subject “Medicine”.

ZR will continue to publish innovative zoological studies and advance knowledge in scientific research. With the proud tradition of attracting outstanding scholars with diverse backgrounds and expertise to write for the journal, as well as the hard work and dedication of our editors and editorial board members, ZR has become a representative international scientific journal in animal studies. With a growing number of international submissions and article citations, as well as an increasing impact factor, we believe that ZR will confidently march into its thirty-seventh year and accomplish its goal of becoming an iconic journal in animal research. ZR could not have achieved what it has without the support of you, our readers. It is a journal that belongs to all members of the animal research community, and we look forward to continuing to work with you all to nurture and carry ZR to the next stage in 2017.

Sincerely



Wai-Yee Chan, Executive Editor-in-Chief
School of Biomedical Sciences,
The Chinese University of Hong Kong, China



Yong-Gang Yao, Editor-in-Chief
Kunming Institute of Zoology,
Chinese Academy of Sciences, Kunming 650223, China



In Memory of Academician Er-Mi Zhao (1930-2016)



Professor Er-Mi Zhao, Herpetologist, 1930-2016

Academician Er-Mi Zhao, Professor of the Chengdu Institute of Biology, Chinese Academy of Sciences, and Sichuan University, passed away on 24 December, 2016, to the great loss of both Chinese and world herpetology. As one of the most internationally renowned Chinese herpetologists, Academician Zhao represented a remarkable era of Chinese herpetology and made a significant contribution to global research, both academically and spiritually. He played a significant role in our understanding of the amphibians and reptiles of Tibet, herpetofauna of the Hengduan Mountains, taxonomy of Chinese snakes, and biogeography of the East Asian islands. His distinguished studies laid a solid and substantial foundation for research by future generations of young herpetologists, and had an extensive and profound impact on the development of the field.

As a prestigious scientist, Academician Zhao won the recognition and respect of his international peers and assured

his place on the worldwide stage based on his remarkable academic achievements. "China should be so proud to have you representing her in herpetology research", Roger Conant, a senior American scientist and *Agkistrodon* expert, wrote to Academician Zhao in 1989. In 1993, *Herpetology of China* (Zhao & Adler, 1993), the first book on the herpetofauna of China, which systematically described the 661 known species at the time, was co-authored by Academician Zhao and American scholar Prof. Kraig Adler. This great work took nearly half a century of meticulous investigation and ten years of compilation to achieve. The book was highly praised by Academician Ilya Darevsky of the Russian Academy of Sciences and Academician David Wake of the American Academy of Sciences, who claimed it to be "a milestone of herpetology literature" and "an epoch-making book, [which] inaugurates a new era in exploring the fauna of amphibians and reptiles of this vast land".

As a diligent researcher, Academician Zhao devoted himself to herpetology research for 60 years. During his academic career, he published more than 140 articles, 43 books, including *Fauna Sinica* (Zhao et al., 1998), *Snakes of China* (Zhao, 2006), *Atlas of Chinese Snakes* (Hu et al., 1980), *Amphibian Zoogeographic Division of China* (Zhao, 1995), and *Herpetology of China* (Zhao & Adler, 1993) in five translations, and established four periodicals. Many of his works were compulsory study for herpetologists and students in China, America, England, and various other countries.

For herpetological classification and zoogeographic regionalization, Academician Zhao discovered a new record family of snake, *Cylindrophis ruffus* (Zhao & Adler, 1989), and 17 new record species of amphibians and reptiles in China. In addition, two new genera of amphibians, *Satobius* (*Hynobius retardatus*) (Adler & Zhao 1990) and *Liua* (*Liua shihi*) (Zhao & Hu, 1983), and 41 new species (or subspecies) were described and named by him, including *Japalura szechwanensis* (Hu & Zhao 1966), *Achalinus meiguensis* (Hu & Zhao, 1966), *Natrix optata* (Hu & Zhao, 1966), *Ranodon tsinpaensis* (Hu et al., 1966), *Megophrys nankiangensis* (Hu et al., 1966), *Hyla tsinlingensis* (Hu et al., 1966), *Rana kuangwuensis* (Hu et al., 1966), *Microhyla mixtura* (Hu et al., 1966), *Leiopisma tsinlingensis* (Hu et al., 1966), *Xenopeltis hainanensis* (Hu et al., 1975), *Dinodon rosozonatum* (Hu et al., 1975), *Achalinus hainanus* (Hu et al., 1975), *Trimeresurus medoensis* (Zhao & Jiang, 1977), *Ovophis monticola zayuensis* (Zhao & Jiang,

DOI: 10.13918/j.issn.2095-8137.2017.006

1977), *Plagiopholis unipostocularis* (Zhao et al., 1978), *Opisthotropis guangxiensis* (Zhao et al., 1978), *Agkistrodon shedaoensis* (Zhao, 1979a), *Trimeresurus xiangchengensis* (Zhao, 1979b), *Phrynocephalus vlangalii hongyuanensis* (Jiang et al., 1980), *Oligodon multizonatum* (Zhao & Jiang, 1981), *Macropisthodon rudis multiprefrontalis* (Zhao & Jiang, 1981), *Scincella huanrenensis* (Zhao & Huang, 1982), *Scutiger ruginosus* (Zhao & Jiang, 1982), *Rhabdophis nuchalis pentasupralabialis* (Jiang & Zhao, 1983), *Platymantis reticulatus* (Zhao & Li, 1984a), *Staurois liangshanensis* (Wu & Zhao, 1984), *Calotes medogensis* (Zhao & Li, 1984b), *Tenuidactylus medogensis* (Zhao & Li, 1987), *Rana tenggerensis* (Zhao et al., 1988), *Eumeces liui* (Hikida & Zhao, 1989), *Trimeresurus mangshanensis* (Zhao & Chen, 1990), *Cuora zhoui* (Zhao et al., 1990), *Amphiesma metusia* (Inger et al., 1990), *Oreolalax multipunctatus* (Wu et al., 1993), *Rana robertingeri* (Wu & Zhao, 1995), *Rhabdophis adleri* (Zhao, 1997), *Laudakia wui* (Zhao, 1998a), *Laudakia papenfussi* (Zhao, 1998b), *Rana zhengi* (Zhao, 1999), *Opisthotropis cheni* (Zhao, 1999), *Rhacophorus hainanus* (Zhao et al., 2005), and *Gloydus lijianlii* (Jiang & Zhao, 2009), as well as the world-renowned Shedao Island pit viper (*Gloydus shedaoensis*) (Zhao, 1979a), Medog green pit viper (*Viridovipera medoensis*) (Zhao & Jiang, 1977), and Mt. Mang pit viper (*Protobothrops mangshanensis*) (Zhao & Chen, 1990). He also added eight new species for Tibet and 10 newly recorded species in China, and was the first to report on the king cobra in Medog (Zhao & Li, 1983) and expand its distribution northwards (Zhao et al., 1986).

Academician Zhao brought new insight and perspective on the geographical division of amphibians and reptiles in China, the amphibian and reptile fauna in the Hengduan Mountain area, the classification of snakes distributed in China, and the zoological geography of the East Asian Islands. He zoned the southern slopes of the Himalayas as a new sub-southwest region of the Indo-Chinese sub-region in the oriental realm according to the actual distribution of species in Tibet, and first proposed a new "South of the Himalayan Sub-region" based on the distribution of reptiles (Zhao et al., 1986). In addition, Academician Zhao also proposed new distribution patterns of amphibians in extra-tropical East Asia and the formation of amphibians and reptiles in the Western Pacific island chain (Zhao, 1989).

Professor Zhao also did outstanding work in scientific practice. He developed measures for the prevention and management of venomous snakes in the grasslands in western Xinjiang, and advanced the concept of medical geography of venomous snake bites in China (Qian et al., 1991) and Yunnan snake medicine to guide the practice of snake bite control. Collaborating with colleges and universities in Hainan, Professor Zhao recruited Masters and Doctoral aquaculture students to cultivate professional personnel for python breeding. He actively cooperated with Hainan python farms by providing systematic scientific knowledge.

Academician Zhao was not only a devoted scholar, but also played active roles in helping young researchers and in disseminating scientific knowledge. He served on the Executive Committee of the International Amphibian and Reptilian Society

in 1983, and as the Chairman of Chinese Herpetologists in the IUCN (1991-2001). During his tenure as an academic advisor on the Editorial Board of *Zoological Research* from 1997 to 2014, he fulfilled his duty faithfully and diligently, and provided vital and constructive promotion of the journal. *Zoological Research* is extremely grateful for his exceptional contribution and dedication, and it is with deep regret and sorrow we note his passing. His deep passion, persistence, optimism, and strict scientific attitude in chasing the truth showed his invaluable spiritual wealth, and is an inspiration to every one of us.

Editor's note: The facts and material stated here in regards to Academician Zhao's academic achievements were provided by his students, Dr. Jia-Tang Li (Chengdu Institute of Biology, CAS), Dr. Jing Che (Kunming Institute of Zoology, CAS), Dr. Song Huang (Huangshan University, China), Peng Guo (Yibin University, China) and Dr. Yue-Zhao Wang (Chengdu Institute of Biology, CAS). This essay was compiled by *Zoological Research*.

REFERENCES

- Adler KA, Zhao EM. 1990. Studies on hynobiid salamanders, with description of a new genus. *Asiatic Herpetological Research*, **3**: 37-45.
- Hikida T, Zhao EM. 1989. *Eumeces liui*: a new species of blue-tailed skink (Lacertilia: Scincidae) from China. *Copeia*, **1989**(1): 110-114.
- Hu BQ, Huang MH, Xie ZT, Zhao EM, Jiang YM, Huang QY, Zong Y, Ma JF. 1980. *Atlas of Chinese Snakes*. Shanghai: Shanghai Scientific and Technical Publishers. (in Chinese)
- Hu SQ, Zhao EM, Liu CZ. 1966. A herpetological survey of the Tsinling and Ta-Pa Shan Region. *Acta Zoologica Sinica*, **18**(1): 57-92. (in Chinese)
- Hu SQ, Zhao EM. 1966. Three new species of reptiles from Szechwan. *Acta Zootaxonomica Sinica*, **3**(2): 158-164. (in Chinese)
- Sichuan Biological Research Institute, Beijing Institute of Zoology. 1975. Three new species of reptiles from Hainan Island, Guangdong Province. *Acta Zoologica Sinica*, **21**(4): 379-384. (in Chinese)
- Inger RF, Zhao EM, Shaffer BH, Wu GF. 1990. *Report on A Collection of Amphibians and Reptiles from Sichuan, China*. Chicago: Field Museum of Natural.
- Jiang F, Zhao EM. 2009. *Gloydus lijianlii*, a new species from the Northern Coastal Islands along Shandong Peninsula (Reptilia, Squamata, Viperidae). *Acta Zootaxonomica Sinica*, **34**(3): 642-646. (in Chinese)
- Jiang YM, Huang QY, Zhao EM. 1980. A new subspecies of *Phrynocephalus vlangalii* (Strauch) and preliminary observations on its ecology. *Acta Zoologica Sinica*, **26**(2): 178-183. (in Chinese)
- Jiang YM, Zhao EM. 1983. Studies on amphibians and reptiles of Mt. Gongga region, Sichuan, China. 3. A study of speciesgroup nuchalis, genus *Rhabdophis*. *Acta Herpetologica Sinica*, **2**(1): 59-62. (in Chinese)
- Qian Y, Zhao EM, Zhao KT. 1991. Animal Science Research: A Symposium Issued to Celebrate the 90th Birthday of the Professor Mangven Ly Chang. Beijing: Chinese Forestry Press, 208-216. (in Chinese)
- Wu GF, Zhao EM. 1984. A rare karyotype of anurans, the karyotype of *Rana phrynoides*. *Acta Herpetologica Sinica*, **3**(4): 5-10. (in Chinese)
- Wu GF, Zhao EM, Inger RF, Shaffer HB. 1993. A new frog of the genus *Oreolalax* (Pelobatidae) from Sichuan, China. *Journal of Herpetology*, **27**(4): 410-413.
- Wu GF, Zhao EM. 1995. A new ranid species of the spinosae group from

- Sichuan. In: Zhao EM. Amphibian Zoogeographic Division of China. A Symposium Issued to Celebrate the Second Asian Herpetological Meeting Held at Ashgabat, Turkmenistan 6 to 10 September 1995. Sichuan Journal of Zoology, Supplement: 52-54. Chengdu, China. (in Chinese) Herpetological Department, Sichuan Biological Research Institute. 1977. A survey of reptiles in Xizang Autonomous Region, with faunal analysis and descriptions of new forms. *Acta Zoologica Sinica*, **23**(1): 64-71. (in Chinese)
- Zhao EM, Jiang YM, Huang QY. 1978. Three new Colubrid species in China. *Materials for Herpetological Research*, **4**: 21. (in Chinese)
- Zhao EM. 1979a. A new Agkistrodon from Shedao (Snake Island), Liaoning. *Acta Herpetologica Sinica*, **1**(1): 4-6. (in Chinese)
- Zhao EM. 1979b. A new snake of the genus *Trimeresurus* from Sichuan, China. *Acta Zootaxonomica Sinica*, **4**(4): 422-424. (in Chinese)
- Zhao EM, Jiang YM. 1981. Studies on amphibians and reptiles of Mt. Gongga Shan, Sichuan, China. I. A new species and a new subspecies of snakes from Sichuan. *Acta Herpetologica Sinica*, **5**(7): 53-58. (in Chinese)
- Zhao EM, Huang KC. 1982. A survey of amphibians and reptiles in Liaoning Province. *Acta Herpetologica Sinica*, **1**(1): 1-23. (in Chinese)
- Zhao EM, Jiang YM. 1982. Studies on amphibians and reptiles of Mt. Gongga Shan, Sichuan, China. 2. A new species of genus *Scutiger* (Amphibia: Pelobatidae). *Acta Herpetologica Sinica*, **1**(1): 79-82. (in Chinese)
- Zhao EM, Hu QX. 1983. Taxonomy and evolution of Hynobiidae in western China, with description of a new genus. *Acta Herpetologica Sinica*, **2**(2): 29-35. (in Chinese)
- Zhao EM, Li SQ. 1983. *Ophiophagus hannah* (Cantor), a Record New to Xizang (Tibet) Autonomous Region, China. *Acta Herpetologica Sinica*, **2**(4): 44. (in Chinese)
- Zhao EM, Li SQ. 1984a. A new species of the genus *Platymantis* (Amphibia: Ranidae) from Xizang. *Acta Herpetologica Sinica*, **3**(3): 55-57. (in Chinese)
- Zhao EM, Li SQ. 1984b. A new species of *Calotes* (Lacertilia: Agamidae) from Xizang (Tibet). *Acta Herpetologica Sinica*, **3**(4): 77-78. (in Chinese)
- Zhao EM, Jiang YM, Li SQ. 1986. Reptilian faunal analysis and zoogeographical division of Xizang Autonomous Region. *Acta Herpetologica Sinica*, **5**(3): 199-203. (in Chinese)
- Zhao EM, Li SQ. 1987. A new lizard of *Tenuidactylus* and a new Tibetan snake record of *Amphiesma*. *Acta Herpetologica Sinica*, **6**(1): 48-51. (in Chinese)
- Zhao EM, Macey JR, Papenfuss TJ. 1988. A new species of *Rana* from Ningxia Hui Autonomous Region. *Chinese Herpetological Research*, **2**(1): 1-3.
- Zhao EM, Adler K. 1989. Asian pipe snake—A snake family new to China. *Sichuan Journal of Zoology*, **8**(2): 26. (in Chinese)
- Zhao EM. 1989. Biogeography of the Amphibians and Reptiles of the Main Islands of East Asia. London: Q. E. D. Recording Services Ltd., Tape 7.
- Zhao EM, Chen YH. 1990. Description of a new species of the genus *Trimeresurus*. *Sichuan Journal of Zoology*, **9**(1): 11-12. (in Chinese)
- Zhao EM, Zhou T, Ye P. 1990. A new Chinese box turtle (Testudinata: Emydidae)-*Cuora zhoui*. In: Zhao EM (editor). From Water onto Land. A symposium issued to commemorate the ninetieth birthday of Professor Cheng-Chao Liu, a distinguished Herpetologist of China. Beijing: Chinese Forestry Press, 213-216. (in Chinese)
- Zhao EM, Adler K. 1993. Herpetology of China. Oxford, Ohio: Society for the study of Amphibians & Reptiles.
- Zhao EM. 1995. Amphibian Zoogeographic Division of China. A Symposium Issued to Celebrate the Second Asian Herpetological Meeting Held at Ashgabat, Turkmenistan 6 to 10 September 1995. Sichuan Journal of Zoology, Supplement: 52-54. Chengdu, China. (in Chinese)
- Zhao EM. 1997. A new species of *Rhabdophis* (Serpentes: Colubridae) from Hainan Island, China. *Asiatic Herpetological Research*, **7**: 166-169.
- Zhao EM, Huang MH, Zong Yu et al. 1998. *Fauna Sinica*: Reptilia, Vol. 3, Squamata, Serpentes. Beijing: Science Press. (in Chinese)
- Zhao EM. 1998a. A new species of the genus *Laudakia* from Xizang (Tibet) Autonomous Region (Sauria: Agamidae). *Acta Zootaxonomica Sinica*, **23**(4): 440-444.
- Zhao EM. 1998b. Description of a new species of the genus *Laudakia* from Xizang (Tibet) (Sauria: Agamidae). *Zoological Research*, **19**(5): 401-404.
- Zhao EM. 1999. Diagnoses of a new frog and a new snake from China. *Sichuan Journal of Zoology*, **18**(3): cover 2. (in Chinese)
- Zhao EM, Wang LJ, Shi HT, Wu GF, Zhao H. 2005. Chinese Rhacophorid frogs and description of a new Species of *Rhacophorus*. *Sichuan Journal of Zoology*, **24**(3): 297-300. (in Chinese)
- Zhao EM. 2006. Snakes of China. Hefei: Anhui Science and Technology Publishing House. (in Chinese)

Engrafted newborn neurons could functionally integrate into the host neuronal network

Zheng-Bo Wang^{1,#}, Dong-Dong Qin^{1,#}, Xin-Tian Hu^{1,2,*}

¹ Key Laboratory of Animal Models and Human Disease Mechanisms of Chinese Academy of Sciences & Yunnan Province, Kunming Institute of Zoology, Kunming Yunnan 650223, China

² CAS Center for Excellence in Brain Science and Intelligence Technology, Chinese Academy of Sciences, Shanghai 200031, China

The limited capability to regenerate new neurons following injuries of the central neural system (CNS) still remains a major challenge for basic and clinical neuroscience. Neural stem cells (NSCs) could nearly have the potential to differentiate into all kinds of neural cells *in vitro*. Previous studies verified that exogenous transplanted NSCs are capable of differentiating into neurons and projecting onto the host neurons in the rat brain (Tabar et al., 2005; Dong JR et al., 2012), which could lead to behavioral recovery from neuronal damages such as spinal cord injury (McDonald et al., 1999), Parkinson's disease (Gonzalez et al., 2015; Kim et al., 2002; Lindvall, 2001), and stroke (Zhang et al., 2016). In January, 2013, the Food and Drug Administration (FDA) of United States has given Neuralstem clearance to commence its human spinal stem cell transplantation in chronic spinal cord injury (Orelli, 2013). In addition, this company is evaluating the safety of stem cell transplantation for the treatment of major depressive disorder (MDD), chronic traumatic encephalopathy (CTE), Alzheimer's disease, and post-traumatic stress disorder (PTSD) (Neuralstem Inc., 2013). However, the mechanism underlying this treatment is controversial. It is still unknown whether the secretion of trophic factors by the transplanted NSCs slowed or prevented the deterioration of the degenerating neurons (Breunig et al., 2011; Xiao et al., 2015; Zuo et al., 2015), or the new neurons differentiated from the transplanted NSCs substituted the injured or lost neurons by functionally integrating into the host neural circuitry (Englund et al., 2002; Weick et al., 2011). To illustrate this question and find out how likely the injured neurons were substituted with new neurons, the functional examination is preferentially to be carried out on an intact animal model (or awake animal). Generally, stem cell clusters transplanted into the brain parenchyma will migrate as single cells towards diverse brain regions. In view of this, approaches that could lock the engrafted cell mass in the specific site were required (Yang SC et al., 2011). Herein, a new technique was developed where a small 'hole' was created in the inferior colliculus (IC) of rhesus monkeys to lock the transplanted NSCs *in situ* and investigate their integration into the host auditory neural network. The results showed that a substantial portion of transplanted cells differentiated into

mature neurons and formed synaptic input/output connections with the host neurons. Under awake condition, c-fos expression in newborn neurons was found to be significantly increased after acoustic stimulation and multichannel recordings indicated IC specific tuning activities in response to auditory stimulation were also recorded in newborn neurons as the reported IC neurons (Zwiers et al., 2004). These results suggest that the transplanted cells have functionally integrated into the host neural network in the awake monkey brain and provided a strong foundation for the future stem cell treatment of the CNS injuries (Wei et al., 2016).

This is the first time evaluating the neurons differentiated from the transplanted NSCs in awake animal, which was further confirmed by Mark Hübener and his group working at the Max Planck Institute of Neurobiology (Falkner et al., 2016). They also found that the transplanted neural stem cells could differentiate into local neuronal phenotype, target projection and functionally integrate with the host neurons. Both of the findings revealed that engrafted stem cells could differentiate into mature neurons, form synaptic with the host neurons, successfully substitute the injured or lost neurons and functionally integrate with the host neuronal network, which might be necessary for the normal brain function of awake animal.

The current findings provide impetus for stem-cell therapy, but if you want to fully assess the ability of transplanted stem cells to differentiate and replace lost neurons in a damaged brain, further sophisticated experimentation should be conducted. For example, the high-throughput dual-color precision imaging (Gong et al., 2016) or transparent intact brain technique (Ke et al., 2013; Susaki et al., 2014; Yang et al., 2014)

Received: 20 December 2016; Accepted: 10 January 2017

Foundation items: This research was supported by National Program on Key Basic Research Project (973 Programs 2015CB755605) and National Natural Science Foundation of China (81471312)

[#]These authors contributed equally to this work

*Corresponding author, E-mail: xthu@mail.kiz.ac.cn

DOI:10.13918/j.issn.2095-8137.2017.005

can be used to reconstruct the engrafted neurons' projections with the host, from which we can overview the newborn neurons and their integration with the host from a systems perspective. All in all, these findings provide encouragement that stem cell replacement therapy will become a reality in the near future.

REFERENCES

- Breunig JJ, Haydar TF, Rakic P. 2011. Neural stem cells: historical perspective and future prospects. *Neuron*, **70**(4): 614-625.
- Dong JR, Guo LY, Qu JG, Qi RL, Wang WC, Xiao CJ, Wang ZB. 2012. Rhesus monkey embryonic stem cells differentiation, proliferation and allotransplantation. *Zoological Research*, **33**(1):43-48.
- Englund U, Björklund A, Wictorin K, Lindvall O, Kokaia M. 2002. Grafted neural stem cells develop into functional pyramidal neurons and integrate into host cortical circuitry. *Proceedings of the National Academy of Sciences of the United States of America*, **99**(26): 17089-17094.
- Falkner S, Grade S, Dimou L, Conzelmann KK, Bonhoeffer T, Götz M, Hübener M. 2016. Transplanted embryonic neurons integrate into adult neocortical circuits. *Nature*, **539**(7628): 248-253.
- Gong H, Xu DL, Yuan J, Li XN, Guo CD, Peng J, Li YX, Schwarz LA, Li AA, Hu BH, Xiong BY, Sun QT, Zhang YL, Liu JP, Zhong QY, Xu TH, Zeng SQ, Luo QM. 2016. High-throughput dual-colour precision imaging for brain-wide connectome with cytoarchitectonic landmarks at the cellular level. *Nature Communications*, **7**: 12142.
- Gonzalez R, Garitaonandia I, Crain A, Poustovoitov M, Abramihina T, Noskov A, Jiang C, Morey R, Laurent LC, Elsworth JD, Snyder EY, Redmond DE, Jr., Semechkin R. 2015. Proof of concept studies exploring the safety and functional activity of human parthenogenetic-derived neural stem cells for the treatment of Parkinson's disease. *Cell Transplantation*, **24**(4): 681-690.
- Ke MT, Fujimoto S, Imai T. 2013. SeeDB: a simple and morphology-preserving optical clearing agent for neuronal circuit reconstruction. *Nature Neuroscience*, **16**(8): 1154-1161.
- Kim JH, Auerbach JM, Rodríguez-Gómez JA, Velasco I, Gavin D, Lumelsky N, Lee SH, Nguyen J, Sánchez-Pernaute R, Bankiewicz K, McKay R. 2002. Dopamine neurons derived from embryonic stem cells function in an animal model of Parkinson's disease. *Nature*, **418**(6893): 50-56.
- Lindvall O. 2001. Parkinson disease. Stem cell transplantation. *Lancet*, **358** Suppl: S48.
- McDonald JW, Liu XZ, Qu Y, Liu S, Mickey SK, Turetsky D, Gottlieb DI, Choi DW. 1999. Transplanted embryonic stem cells survive, differentiate and promote recovery in injured rat spinal cord. *Nature Medicine*, **5**(12): 1410-1412.
- Neuralstem, Inc. 2013. Neuralstem Receives FDA Approval To Commence Spinal Cord Injury Trial.
- Orelli B. 2013. FDA OKs Neuralstem Trial Treating Spinal Cord Injuries With Stem Cells.
- Susaki EA, Tainaka K, Perrin D, Kishino F, Tawara T, Watanabe TM, Yokoyama C, Onoe H, Eguchi M, Yamaguchi S, Abe T, Kiyonari H, Shimizu Y, Miyawaki A, Yokota H, Ueda HR. 2014. Whole-brain imaging with single-cell resolution using chemical cocktails and computational analysis. *Cell*, **157**(3): 726-739.
- Tabar V, Panagiotakos G, Greenberg ED, Chan BK, Sadelain M, Gutin PH, Studer L. 2005. Migration and differentiation of neural precursors derived from human embryonic stem cells in the rat brain. *Nature Biotechnology*, **23**(5): 601-606.
- Wei JK, Wang WC, Zhai RW, Zhang YH, Yang SC, Rizak J, Li L, Xu LQ, Liu L, Pan MK, Hu YZ, Ghanemi A, Wu J, Yang LC, Li H, Lv LB, Li JL, Yao YG, Xu L, Feng XL, Yin Y, Qin DD, Hu XT, Wang ZB. 2016. Neurons differentiated from transplanted stem cells respond functionally to acoustic stimuli in the awake monkey brain. *Cell Reports*, **16**(4): 1016-1025.
- Weick JP, Liu Y, Zhang SC. 2011. Human embryonic stem cell-derived neurons adopt and regulate the activity of an established neural network. *Proceedings of the National Academy of Sciences of the United States of America*, **108**(50): 20189-20194.
- Xiao JJ, Yin M, Wang ZJ, Wang XP. 2015. Transplanted neural stem cells: playing a neuroprotective role by ceruloplasmin in the substantia nigra of pd model rats? *Oxidative Medicine and Cellular Longevity*, **2015**: 618631.
- Yang B, Treweek JB, Kulkarni RP, Deverman BE, Chen CK, Lubeck E, Shah S, Cai L, Gradinaru V. 2014. Single-cell phenotyping within transparent intact tissue through whole-body clearing. *Cell*, **158**(4): 945-958.
- Yang SC, Dong JR, Qu JG, Hu XT, Wang ZB. 2011. A traumatic brain injury model for distinguishing between transplanted neural cells and host cells in vivo. *Zoological Research*, **32**(4):421-427.
- Zhang RL, Zhang ZG, Chopp M. 2016. Function of neural stem cells in ischemic brain repair processes. *Journal of Cerebral Blood Flow & Metabolism*, **36**(12): 2034-2043.
- Zuo FX, Bao XJ, Sun XC, Wu J, Bai QR, Chen G, Li XY, Zhou QY, Yang YF, Shen Q, Wang RZ. 2015. Transplantation of human neural stem cells in a parkinsonian model exerts neuroprotection via regulation of the host microenvironment. *International Journal of Molecular Sciences*, **16**(11): 26473-26492.
- Zwiers MP, Versnel H, Van Opstal AJ. 2004. Involvement of monkey inferior colliculus in spatial hearing. *Journal of Neuroscience*, **24**(17): 4145-4156.

Consequences of early adverse rearing experience (EARE) on development: insights from non-human primate studies

Bo Zhang^{1,2,3,*}

¹ Yunnan Key Laboratory of Primate Biomedical Research, Kunming Yunnan 650500, China

² Institute of Primate Translational Medicine, Kunming University of Science and Technology, Kunming Yunnan 650500, China

³ National Institute of Health, Bethesda, Maryland, USA

ABSTRACT

Early rearing experiences are important in one's whole life, whereas early adverse rearing experience (EARE) is usually related to various physical and mental disorders in later life. Although there were many studies on human and animals, regarding the effect of EARE on brain development, neuroendocrine systems, as well as the consequential mental disorders and behavioral abnormalities, the underlying mechanisms remain unclear. Due to the close genetic relationship and similarity in social organizations with humans, non-human primate (NHP) studies were performed for over 60 years. Various EARE models were developed to disrupt the early normal interactions between infants and mothers or peers. Those studies provided important insights of EARE induced effects on the physiological and behavioral systems of NHPs across life span, such as social behaviors (including disturbance behavior, social deficiency, sexual behavior, etc), learning and memory ability, brain structural and functional developments (including influences on neurons and glia cells, neuroendocrine systems, e.g., hypothalamic-pituitary-adrenal (HPA) axis, etc). In this review, the effects of EARE and the underlying epigenetic mechanisms were comprehensively summarized and the possibility of rehabilitation was discussed.

Keywords: Early adverse rearing experience; Non-human primates

INTRODUCTION

One of factors affecting life-long health of humans is the stability of early childhood, especially children's relationship with their mothers. John Bowlby's attachment theory suggests that

individual's social relationship throughout life is influenced by the initial attachment with the mother (Bowlby, 1969). Attachment theory is a psychological, evolutionary and ethological theory concerning relationships among humans. Within the theory, attachment means an affectional bond or tie between an individual and an attachment figure (usually a caregiver). The core is that a child needs to build relationship with at least one primary caregiver to develop normal social and emotional behaviors. In many orphans, the lack of normal attachment to parents would cause behavioral and physical problems in childhood and possibly continuing throughout adult life (McEwen, 2003). Adults with adverse experience were more vulnerable to physical, psychosocial and mental disorders (Maughan & McCarthy, 1997; Pirkola et al., 2005). In human, early adverse rearing experience (EARE) usually refers to child abuse, which is a worldwide problem and is defined as neglect or physical, sexual or emotional mistreatment or abuse of children (Newton & Vandeven, 2009, 2010). Although human based studies revealed compelling associations between EARE and psychological outcomes, both retrospective and prospective studies showed their limits, e.g., inaccurate self-report due to biased or even false memory, failure in controlling accompanying environmental and genetic factors. Therefore, the long-term effects of EARE on subjects were usually not the direct consequences, but were inevitably intervened or masked by uncontrollable factors. However, experimental animals can be raised in laboratory environments, therefore allow researchers to carry out randomized prospective longitudinal studies, e.g., rigorously control or systematically manipulate early experiences throughout the entire period of investigation.

Received: 01 December 2016; Accepted: 30 December 2016

Foundation items: This research was supported by Hainan special fund project for science and technology (KJHZ2015-20)

*Corresponding author, E-mail: bozhangpp@foxmail.com

DOI:10.13918/j.issn.2095-8137.2017.002

Rodents are easy to manipulate genetically, and the related studies indicate EARE as a developmental risk factor with profound, long-term effects on later life (Meaney, 2001; Pryce et al., 2005b; Sánchez et al., 2001). Whereas the high similarities of NHPs with humans make it irreplaceable in investigating the effects of EARE on physiological and behavioral development, e.g., NHPs and chimpanzees in specific, share over 90% and 98.8% genomes with human beings, respectively (Lovejoy, 1981). High similarities were found in both biological (Azmitia & Gannon, 1986; Uylings & van Eden, 1991) and socio-ecological aspects, e.g., social organizations and clear dominance hierarchies (Bailey & Aunger, 1990; DeVore, 1990; Wright, 1990). The phenomenon that in NHPs, 2%-10% of infants were physically abused or neglected by their mothers in group-living conditions, allow the possible screening of natural child abuse models (Maestriperi & Carroll, 1998; Maestriperi et al., 1997). Moreover, like humans NHPs has prolonged postnatal period of maturation during which mother–infant relationship and neural system development can be influenced by environment and early life experience (Levine & Wiener, 1988; Suomi, 2005).

Harlow (Harlow & Harlow, 1965) introduced the concept of affectional systems to characterize the relationships in the social groups of primates, and five distinct affectional systems were described, including the infant-mother affectional system, the maternal affectional system, the age-mate/peer affectional system, the heterosexual affectional system and the paternal affectional system. The infant-mother and the maternal affectional systems in Harlow's affectional systems are similar to Bowlby's concept of mother-infant attachment theory in humans. In normal living group, most monkey infants virtually spend all of their initial days or weeks of life clinging with their biological mothers, ventral to ventral, during which, specific and strong attachment bonds are built. When about 2-month old, infants begin to explore the physical and social environment, spending increasing amount of time participating social interactions, especially playing with peers. From 6-month of age until puberty, playing with peers becomes the major social activity (Hinde & Spencer-Booth, 1967; Suomi, 1997, 2005). In fact, the infant and juvenile monkeys always maintain a close social relationship with their mothers, while the mother plays the role of protector especially under stressful situations, and mentor in teaching developing appropriate social behaviors. Accordingly, the studies regarding EARE usually involve disruption of the normal infant-mother relationships, by maternal deprivation of newborns, maternal separation or induced stress on older infants and juvenile monkeys. Although some epidemiological studies in humans suggest possible direct relationships between EARE and abnormal behaviors in later life, no solid evidence was raised to prove the precise impact of childhood adversities on psychiatric disorders (Benjet, 2010; Bick & Nelson, 2016; Gershon et al., 2013; Kessler et al., 1997; Kessler & Wang, 2008; Klein et al., 2013; Sheridan et al., 2010). The over 60 years NHP studies shed lights on the understanding of the influences of EARE on physiological and behavioral development, including social behaviors (e.g., disturbance behavior, social deficiency, sexual behavior, etc),

learning and memory ability, brain structural and functional development (e.g., development of neurons and glia cells, neuroendocrine dysregulation, etc). In this review, the previous findings on EARE were systematically summarized, and the underlying epigenetic mechanisms and the potential methods of rehabilitation were thoroughly discussed.

EARE MODELS IN NON-HUMAN PRIMATES

Controlled rearing conditions in standard laboratory settings are designed to simulate natural environments. The infants are reared by their mothers and live in a group consisted of other infants, juveniles and adults, allowing infants to be exposed to complex social interactions. In abnormal rearing conditions, the mother deprivation method is applied. The newborn is taken away from their mothers at birth and is reared in incubators with regular medical attention and laboratory nursery. A period of time (usually 1-month) later, when able to feed themselves, infants are moved to other rearing conditions depending on aims of research, e.g., be reared alone in social isolation condition, with nursery/peers of the same age in nursery/peer rearing condition, with a surrogate in surrogate mother/foster rearing condition, etc. They could also be separated from mothers at later time for once (temporary maternal separations) or several times (repetitive maternal separations); or even though staying with their mother all the time, but still suffer from EARE (maternal neglect).

Social isolation

Social isolation (including total and partial social isolation) is initially described in early 1960s by Harlow and his colleagues, and has been used ever since to raise monkeys in simulating social behavior deficits in humans (Table 1). In total isolation, the infant is reared in a cage alone without any auditory, visual, olfactory and tactile contact with conspecifics, including mothers, peers and other monkeys (Baysinger et al., 1972; Harlow & Harlow, 1962; Harlow et al., 1964, 1965). In partial social isolation, although infants are separately caged from their mothers, peers, and social groups, they have auditory, visual, and olfactory but not tactile contact with their conspecifics (Cross & Harlow, 1965; Mason & Sponholz, 1963; Struble & Riesen, 1978; Suomi et al., 1971). These early studies by Harlow and his colleagues, especially their extreme manipulations, including total isolation, "pit of despair" and "rape rack" devices, were controversial and were most likely forbidden to perform due to ethical issues. In 1950s, many researchers assumed that the only necessity of mother was supplying food to infants, whereas excessive intimacy between mother and infant would hinder the growth of infant, or even induce over dependence in adulthood. Harlow disagreed with the viewpoints; performed a series of isolation studies on primates to prove that to acquire necessary social skills, to obtain both physically and psychologically healthy development, infants need mothers' affection, as well as normal social interaction and emotional relationship with peers. However, their intention was to prove the essential role of mother's love to infants, in the

Table 1 Early adverse rearing experience (EARE) methods

Paradigms		Description			References
Social Isolation	Total	Infants are reared in a cage alone at birth, no any auditory, visual, olfactory and tactile contact with conspecifics is allowed			Baysinger et al., 1972; Harlow et al., 1965; Harlow & Harlow, 1962; Harlow et al., 1964
	Partial	Infants are separately caged at birth, reared with auditory, visual, and olfactory, but not tactile contact with conspecifics			Cross & Harlow, 1965; Mason & Sponholz, 1963; Struble & Riesen, 1978; Suomi et al. 1971
Maternal separation	Permanent	Peer-rearing	Continuous	Infants are reared by pairs throughout development	Chamove et al., 1973; Erwin et al., 1973; Sackett, 1967; Worlein & Sackett, 1997
			Intermittent	Peers are allowed to contact with each other for a limited period of time and then infants are housed singly during the rest of the time	Rommeck et al., 2009b
			Rotational	Infants are continuously housed with different peers	Novak & Sackett, 1997; Rommeck et al., 2009b
	Temporary	Surrogate mothers rearing (SMR)	Inanimate objects are placed into the cage as an artificial surrogate mother		Capitanio & Mason, 2000; Dettmer et al., 2008; Schneider & Suomi, 1992; Suomi, 1973
		Surrogate-peer rearing (SPR)	Combination of SMR and PR		Bastian et al., 2003; Lutz et al., 2007; Meyer et al., 1975
		One time	Repetitive	Infants are taken away from their mothers at later stages of life for a period of time, followed by mother-infant reunion	Hinde & McGinnis, 1977; Hinde et al., 1966; Kaufman & Rosenblum, 1967; Seay et al., 1962; Spencer-Booth & Hinde, 1971
Repeatedly separating infants from their natal group for relatively short periods of time, followed by repeated reunions	Clarke et al., 1998; Dettling et al., 2002a, b; Levine & Mody, 2003; Sánchez et al., 2005; Suomi et al., 1983				
Maternal neglect			Infant-mother was confronted with various foraging conditions to induce different levels of stress in the mother	Andrews & Rosenblum, 1991; Coplan et al., 1996; Rosenblum & Andrews, 1994; Rosenblum & Pauly, 1984	

form of her availability all the time, her physical touching, caring and protection, which was an obvious fact to us today without any necessity to prove.

However, although isolation models are important in highlighting the devastating consequences of maternal deprivation, the extreme manipulations could induce severe cognitive and emotional deficits, or even self-injurious behaviors, which are very difficult to remediate in primates. Therefore, less severe rearing conditions were developed afterwards at least partially due to ethical considerations.

Maternal separations

Peer-rearing (PR) (or nursery rearing, NR) (including continuous pair rearing, intermittent and rotational peer rearing) is another widely used rearing condition, in which infants were reared together with peers of the same age (Chamove et al., 1973; Erwin et al., 1973; Sackett, 1967; Worlein & Sackett, 1997) (Table 1). In continuous pair rearing condition, infants are usually reared by pairs throughout development (Chamove et al., 1973; Fekete et al., 2000; Hotchkiss & Paule, 2003; Novak & Sackett, 1997). Intermittent peer rearing allow peers to contact with each other for a limited period of time, and then infants are housed singly during the rest of the time (Rommeck et al., 2009b). Within the rotational peer rearing condition, infants are continuously peer housed with different infant partners (Novak & Sackett, 1997; Rommeck et al., 2009b).

Previous study showed that continuous rotational pairing induces a behavioral profile quite similar with that of mother rearing in socially complex environment (Rommeck et al., 2011). Compared with social isolation, PR is less severe and thus more widely used in recent NHP EARE studies. Surrogate mothers rearing (SMR) is another early rearing method, in which inanimate object is placed into the cage as an artificial surrogate mother (Capitaniao & Mason, 2000; Dettmer et al., 2008; Eastman & Mason, 1975; Harlow, 1958; Harlow & Zimmermann, 1959; Hennessy & Kaplan, 1982; Kaplan, 1974; Mason & Berkson, 1975; Roy et al., 1978; Schneider & Suomi, 1992; Suomi, 1973). Infants could quickly develop attachment with surrogate mothers, and some studies indicated that the infants usually preferred cloth surrogate mothers than wired ones (Harlow, 1958; Harlow & Zimmermann, 1959). Previous study reported that surrogate mothers could affect the behaviors of infants, and different characters of surrogate mothers such as mobility and orientation had different influences (Dettmer et al., 2008). Surrogate-peer rearing (SPR) method is a combination of SMR and PR, in which the infants are reared with inanimate surrogate mothers (SMR condition) during the initial several months of life, and then are allowed to have peer interactions for a limited period of time (PR condition) (Bastian et al., 2003; Lutz et al., 2007; Meyer et al., 1975). Comparing with permanent removal of the mother, infants are not separated from their mothers right away at born in

temporary maternal separations, but after a period of time usually several hours, days or weeks, following by mother-infant reunion (Hinde et al., 1966; Hinde & McGinnis, 1977; Kaufman & Rosenblum, 1967; Seay et al., 1962; Spencer-Booth & Hinde, 1971). Temporary maternal separation usually contains a one-time separation although different time delay could be adopted. A modified version of one-time separation is repetitive mother-infant separation, in which infants are separated from and reunited with their natal group repeatedly for relatively short periods of time (Clarke et al., 1998; Dettling et al., 2002b; Levine & Mody, 2003; Sánchez et al., 2005; Suomi et al., 1983). The impact of these procedures appeared to be further intensified if the separations were unpredictable (Levine, 2000; Sánchez et al., 2005). Unlike social isolation, maternal separation adopted relatively mild manipulations, the presence of surrogate mothers and the opportunity of direct contact with mothers and peers added social complexity to the infants' living environment, therefore could avoid severe social and emotional deficits associated with mothers' absence.

Maternal neglect

Compared with isolation and maternal separation methods described above, maternal abuse and neglect during early life are more common in humans, therefore are more widely used on NHPs to study adult mood and anxiety disorders. In NHP maternal neglect models, in order to induce stress in the mother, infant mothers are confronted with various foraging conditions, such as variable/unpredictable foraging demand (VFD), consistently low foraging demand (LFD) and consistently high (but predictable) foraging demand (HFD). Mothers in LFD condition have easy access to food while those in high foraging

demand have to work hard to get food (Andrews & Rosenblum, 1991; Coplan et al., 1996; Rosenblum & Andrews, 1994; Rosenblum & Pauly, 1984). The advantage of this model is that even though infants are still in adverse situation, the severe adverse experience of mother and peer deprivation can be avoided. In addition, other rearing strategies are applied in this model, i.e., infants were reared by a female which was not their biological mother (Maestripieri, 2005; Novak & Suomi, 1991); infants were housed with non-reproductive female adults (Champoux et al., 1989b).

EARE EFFECTS

Although partial social isolation tends to induce less severe defects than total social isolation, the expression of behavior defects is similar. Isolated monkeys reared without exposure to companions during early life, especially the first 6 months, develop a pervasive pattern of abnormalities referred to as the isolation syndrome. Mason (Mason, 1968) summarized the syndrome under four headings: (1) abnormal posturing and movements, such as rocking; (2) motivational disturbances, such as excessive fearfulness or arousal; (3) poor integration of motor patterns, such as inadequate sexual behavior; (4) deficiencies in social communication, such as failure to withdraw after being threatened by an aggressing animal. In this section, the effects of EARE on social behaviors, learning and memory ability, brain structural and functional developments, including influences on neurons and glia cells, neuroendocrine dysregulation, especially stress related HPA axis will be reviewed.

Social behavior

Effects of EARE on social behaviors are detailed in Table 2.

Table 2 Effects of EARE on social behaviors

Behavior types	Behavior descriptions
Stereotypic behaviors	Whole-body stereotypes (e.g., rocking, pacing, bouncing, swing, and back-flipping) Self-directed stereotypes (e.g., saluting, digit-sucking, self-clutching, self-clasping, eye-poking, eye-covering and hair-pulling)
Self-directed behaviors	Self-manipulation, self-scratching, self-grasping, self-rubbing Self-injurious behavior (SIB)
Aggression	Less aggression during infancy and more aggression during later life
Affiliative behavior	Tend to show more affiliative behavior during infancy but less affiliative behavior during adulthood
Social and environmental exploration	Decreased social and environmental exploration
Social dominance	Tend to show low dominance rank
Sexual behaviors	Less and abnormal sexual behaviors
Others	Polyphagia and polydipsia

Disturbance behavior

Monkeys exposed to adverse early experience tended to show more disturbance behaviors, such as stereotypic and self-directed behaviors, motivational disturbances and social deficiency. The isolated monkeys appeared to show more disturbance behaviors, including crouching, clutching, rocking, pacing, flipping, hugging, clasping, thumb-sucking (Harlow &

Harlow, 1962; Harlow & Suomi, 1971a; Mason & Sponholz, 1963; Mitchell, 1968; Suomi et al., 1971). Among these monkeys, some abnormal movements, such as rocking and self-grasping, could present very early in their lives, even at the first month (Baysinger et al., 1972). Additionally, some of these behaviors could turn into stereotypic behaviors, including repetitive movements or postures, as well as ritualized movements, and could be divided into whole-body stereotypes

(e.g., rocking, pacing, bouncing, swing, and back-flipping), self-directed stereotypes (e.g., saluting, digit-sucking, self-clutching, self-clasping, eye-poking, eye-covering and hair-pulling) and other idiosyncratic behaviors (e.g., teeth grinding, head tossing, or making noise by blowing air into the cheeks). It was reported that whole-body stereotypes were much more common than self-directed stereotypes (Lutz et al., 2003). Previous studies indicated that isolated monkeys showed more repetitive whole-body stereotypes (Mitchell, 1968), while PR monkeys showed more self-directed stereotypes (Lutz et al., 2003; Suomi et al., 1971).

EARE exposed monkeys tended to show more self-directed behaviors. Isolated monkeys showed self-manipulation, self-scratching, self-grasping, self-rubbing, and autoeroticism while in isolation (Baysinger et al., 1972), or showed remarkable increases in self-clasping soon after removal from isolation (Harlow et al., 1965; Suomi et al., 1974), or self-clutching after surrogate mother removing (Harlow & Zimmermann, 1959). PR reared infants and juvenile monkeys showed increased self-stimulation behaviors, including self-sucking, self-clinging, self-clasping and other self-directed behaviors (Champoux et al., 1991; Lutz et al., 2003; Suomi et al., 1971). Moreover, short-term stress by temporary physical restrictions could also induce significant increases in self-clasping and huddling behaviors when the infants returned to their home cages (Harlow & Suomi, 1971a). Those self-directed behaviors often turned into self-injurious behavior (SIB), with males showing a much higher level of vulnerability than females (Cross & Harlow, 1965; Gluck & Sackett, 1974; Lutz et al., 2003; Suomi et al., 1971), and PR monkeys usually much more vulnerable than MR monkeys (Rommeck et al., 2009a). Surrogate mothers appeared to provide a certain degree of contact acceptability, security and trust sufficient for isolated monkeys to suppress existing self-directed disturbance activity, and to initiate crude social interactions with other isolated monkeys (Harlow & Suomi, 1971b). However, Lutz et al. (Lutz et al., 2007) reported that SPR monkeys showed significantly more self-biting comparing to PR and MR reared animals, and it was suggested that surrogate rearing in combination with lower levels of social contact during play may be risk factors for the later development of self-biting behavior. Actually, self-directed behaviors were hypothesized to result from the redirection of normal social behaviors toward one's own body and were suggested to be symptoms of some mental diseases (Goosen, 1981; Mason & Berkson, 1975). These findings indicate that EARE exposed monkeys could be used as an ideal model of related human mental disorders from behavioral perspective.

Social deficiency

In natural environments, infants and juvenile monkeys are supposed to be more active in joining the social play with peers, but monkeys exposed to EARE show decreased social playing. Isolated monkeys showed less (Harlow et al., 1965; Mitchell, 1968), or even no contact playing at all (Harlow et al., 1965). Pair and peer reared infants (Chamove et al., 1973), VFD reared infants (Andrews & Rosenblum, 1991; Rosenblum & Pauly, 1984), repeated parental deprivation infants (Dettling et al., 2002b; Levine & Mody, 2003) all showed less social playing

compared with MR infants. Lack of sufficient social interaction led to the fact that EARE exposed monkeys could not successfully adapt to living in a large social group (Griffin & Harlow, 1966; Harlow & Harlow, 1962; Mason & Sponholz, 1963; Ruppenthal et al., 1991). Not only social interaction, studies also showed decreased environmental exploration in isolated monkeys (Griffin & Harlow, 1966; Mason & Sponholz, 1963; Mitchell, 1968), VFD and PR monkeys (Rosenblum & Pauly, 1984; Ruppenthal et al., 1991). Another major index of exploratory behavior is locomotor activity, while some NHP studies showed less locomotion in isolated adults (Harlow & Suomi, 1971a; Mason & Sponholz, 1963; Mitchell, 1968) and PR infants (Feng et al., 2011), others found no differences in PR adults (Winslow et al., 2003), or even higher activity levels in PR infants during the first month after isolation (Champoux et al., 1991). Therefore, there was no agreed tendency of EARE influence on locomotor activity in monkeys, making it an invalid measure of exploratory behavior if used alone (Wright, 1983).

Another domain of EARE induced social deficiency is social dominance. In monkey society, social dominance is a complex phenomenon mediated by different mechanisms and various factors such as kinship, age, sex, and physical factors like body weight, appearance and health (Bernstein & Cooper, 1999; Bernstein & Mason, 1963; Morgan et al., 2000; Sprague, 1998; Takahashi, 2002). Kinship seemed to be the major factor in determining dominant rank at least until puberty (Koford, 1963; Koyama, 1967), but became weaker during the development (Bernstein & Williams, 1983). Both dominance formation and maintenance among males in a living group are usually achieved by aggressive behavior such as fighting, with the stronger and more aggressive subjects winning and thus becoming dominant. However, appropriate use of aggression is critical for both acquiring and maintaining social status, as overly aggressive monkeys may risk social ostracism from their conspecifics. Moreover, aggressive behavior was not indispensable to obtain and keep dominance status and dominance sustained without aggression was more stable than that formed on the basis of aggression (Fonberg, 1988).

Monkeys exposed to EARE tended to show less aggression during infancy (Chamove et al., 1973; Harlow et al., 1965), and more aggression during later life (Chamove et al., 1973; Mitchell, 1968; Suomi et al., 1974; Winslow et al., 2003). The aggressive monkeys exposed to EARE may repeatedly attack a helpless infant or attempt to attack a dominant male, while infant-directed aggression is abnormal adult-directed aggression is both abnormal and suicidal (Chamove et al., 1973; Mitchell, 1968; Suomi et al., 1974; Winslow et al., 2003). On the other hand, studies showed EARE exposed monkeys showed heightened fear in all age stages (Champoux et al., 1991; Dettling et al., 2002b; Levine & Mody, 2003; Mitchell, 1968). It seems that EARE makes monkeys more emotional in two opposite directions, both aggression and fear. In addition to aggression, affiliative behavior, such as grooming and proximity, is also important in establishing and maintaining alliances and reinforcing the dominance hierarchy. Affiliative behavior was suggested to be more positively related to dominance rank than kinship in Japanese monkeys (Singh et al., 1992). On the

contrary to aggression, EARE exposed monkeys showed more affiliative behavior during infancy (Chamove et al., 1973; Rosenblum & Pauly, 1984; Ruppenthal et al., 1991), but less affiliative behavior during adulthood (Kraemer & McKinney, 1979; Rosenblum & Pauly, 1984; Winslow et al., 2003). With more aggressive and less affiliative behavior which both contribute to acquiring and reinforcing social dominance, EARE exposed adult monkeys are supposed to have low social dominant rank in a living group, and studies indeed indicated that both isolated and PR adult monkeys showed low social dominance (Kraemer & McKinney, 1979; Mitchell, 1968; Ruppenthal et al., 1991).

Sexual behavior

Monkeys exposed to EARE demonstrated less or abnormal sexual behaviors (Chamove et al., 1973; Harlow et al., 1966; Harlow, 1962; Harlow et al., 1965; Mitchell, 1968). Abnormal sexual behaviors (abortive mount) is defined as any improperly oriented mount, accompanied by pelvic thrusting including standing-to-head, standing-to-side and ventral lie-on (Wallen et al., 1981). Males usually were not mount properly as they engaged in varied but misplaced heterosexual efforts, while females were not maintain the sexual present (stood quadrupedally with the perineal area directed towards the recipient) or turned their bodies when mounted. Mount behavior includes no-foot-clasp mount and foot-clasp mount, which could be differentially affect by different EARE. Males with short access periods with peers (0.5 h) rarely or never foot-clasp-mounted peers, while those given 24 h access regularly foot-clasp-mounted peers (Wallen et al., 1981). Isosexually reared males showed less foot-clasp mounting and more presenting than heterosexual males, while conversely, isosexually reared females showed statistically more mounting and less presenting than heterosexual females (Goldfoot et al., 1984). Moreover, females exposed to EARE also showed abnormal maternal behaviors, in a way that those never experienced mother caring not only were unable to exhibit caring to their own offspring, but also far more likely to display inadequate, abusive or neglectful behavior toward their offspring (Bridges et al., 2008; Champoux et al., 1992; Harlow & Suomi, 1971b; Seay et al., 1964; Suomi, 1978; Suomi et al., 1974; Suomi & Ripp, 1983), consistent with human findings showing abusive behavior appeared to be transmitted across generations (Roustit et al., 2009).

Primate studies also showed other EARE induced behavioral effects besides listed above, including polyphagia and polydipsia in isolated adults (Miller et al., 1969), more vulnerable to excessive alcohol consumption (Fahlke et al., 2000; Higley et al., 1991) and elevated response to both aversive and rewarding stimuli (Nelson et al., 2009) in PR monkeys and abnormal sleep rhythmicity (Barrett et al., 2009; Boccia et al., 1989; Kaemingk & Reite, 1987; Reite et al., 1974; Reite & Short, 1978). An interesting research showed EARE significantly influenced the development of lateralisation, as PR monkeys demonstrated greater left-hand bias compared to MR reared monkeys (Bennett et al., 2008). Despite of EARE effects described above, recent research suggested that modern PR practices might not result in inevitable perturbations in

aggressive, rank-related, sexual, and emotional behavior in rhesus monkeys (Bauer & Baker, 2016).

Learning and memory

Early primate studies showed EARE exposed adults performed adequately on simple discriminations or delayed-response (Gluck et al., 1973), but showed impairments in certain complex tasks such as those requiring engaging working memory with dynamic rules or delays or response inhibition (Beauchamp & Gluck, 1988; Beauchamp et al., 1991; Gluck et al., 1973; Gluck & Sackett, 1976; Sánchez et al., 1998). These results were obtained mostly from adult monkeys separated from their mothers at birth and reared in total isolation for 9-12 months. PR reared juvenile monkeys also showed cognitive deficits, they had more difficulty acquiring the delayed non-matching to sample (DNMS) task and were also impaired in object but not spatial reversal learning (Sánchez et al., 1998). Moreover, even brief social isolation impaired performance in a multiple video-task assessment in adult rhesus monkeys (Washburn & Rumbaugh, 1991) and impaired reversal learning and behavioral inhibition in adult marmosets (Pryce et al., 2004a, b). These results were consistent with the results of human studies, which showed the post institutionalized children (Bauer et al., 2009) and childhood exposed to neglect and abuse (Majer et al., 2010) were associated with impaired learning and memory during adulthood. Although those studies revealed EARE induced impairment of learning and memory ability in a task dependent way in adult monkeys, other primate studies indicate exposure to mild early life stress improves prefrontal dependent response inhibition in primates, suggesting its beneficial effect on cognitive control (Parker et al., 2005, 2012).

Brain structure and function

The first documentation of the effects of negative early experiences on monkey brain was provided by Martin et al. (1991), which showed significant alterations in the chemo architecture of the striatum 19-24 years after social deprivation. Additionally, Siegel et al. (1993) demonstrated that early social deprivation resulted in an increase in the amount of non-phosphorylated neurofilament protein in hippocampal dentate gyrus granule cells in rhesus monkeys. Further studies showed structure and function changes in many brain regions including amygdala, hippocampus, prefrontal cortex (PFC), anterior cingulate cortex (ACC), corpus callosum and cerebellum etc, both in humans and animals exposed to EARE (Andersen, 2015; Bick & Nelson, 2016; Gilmer & McKinney, 2003; Gorman et al., 2002; Hart & Rubia, 2012; Korosi et al., 2012; McEwen, 2003; Worlein, 2014)(Table 3).

Amygdala

Amygdala is a group of almond-shaped nuclei located deep within the medial temporal lobes of the brain in complex vertebrates. It was considered as the emotion center and responsible for emotion reactions like reward, fear and anxiety (Davis, 1992; Gallagher & Chiba, 1996; Ledoux, 2003; Phelps, 2006). Rodent studies showed acceleration of amygdala development in early weaning rodents (Kikusui & Mori, 2009;

Table 3 Effects of EARE on brain structure and function

	Outcomes		Human Studies	Primate Studies
Amygdala	Children	No significant volumes changes	De Bellis et al., 2001; De Brito et al., 2013; Hanson et al., 2010; Woon & Hedges, 2008	No significant volume changes (Howell et al., 2014);
		Larger volume and elevated response	Lupien et al., 2011; Mehta et al., 2009; Tottenham et al., 2010	Decreased SERT binding potential (Ichise et al., 2006);
		Decreased volume	Edmiston et al., 2011; Hanson et al., 2015; Luby et al., 2013	Differential expression of one gene GUCY1A3 (Sabatini et al., 2007)
	Adults	No significant volume changes	Bremner et al., 1997; Cohen et al., 2006	
		Larger volume	Evans et al., 2016; Lyons-Ruth et al., 2016	
		Elevated activity	Casement et al., 2014; Javanbakht et al., 2015; Kim et al., 2013	
Hippocampus	Children	Decreased volume	Edmiston et al., 2011; Hanson et al., 2015; Luby et al., 2013	No significant volume change (Law et al., 2009a, b; Sánchez et al., 1998; Spinelli et al., 2009)
		No significant volume change	Carrion et al., 2001; De Bellis et al., 2001; De Bellis et al., 1999; De Bellis et al., 2002; Mehta et al., 2009; Tottenham et al., 2010; Woon & Hedges, 2008	
	Adults	Decreased volume	Bremner et al., 1997; Cohen et al., 2006; Stein et al., 1997; Woon & Hedges, 2008	
Prefrontal cortex (PFC)	Children	No significant volume changes	De Bellis et al., 1999	Greater enlarged medial prefrontal cortex (mPFC) size (Spinelli et al., 2009)
			De Bellis et al., 2002; Edmiston et al., 2011; Hanson et al., 2010; Morey et al., 2016; Thomaes et al., 2010	
		Decreased volume	Carrion et al., 2009; Richert et al., 2006	
		Larger volume	Tomoda et al., 2009; van Harmelen et al., 2010	
	Adults	Decreased volume	Casement et al., 2015; Kim et al., 2013; Romens et al., 2015; Schweizer et al., 2016	
		Reduced activity	Casement et al., 2014; Javanbakht et al., 2015; Jedd et al., 2015; Wang et al., 2016; White et al., 2015	
		Increased response		

Ono et al., 2008). The limited amount of primate studies found no significant amygdala volume changes (Howell et al., 2014), but functional changes including decreased SERT binding potential (Ichise et al., 2006) and differential expression of one gene GUCY1A3 (Sabatini et al., 2007) in amygdala of EARE exposed monkeys. However, human studies in maltreated children showed contrary results, with some studies found no volume changes (De Bellis et al., 2001; De Brito et al., 2013; Hanson et al., 2010; Woon & Hedges, 2008), while others revealed decreased volume (Edmiston et al., 2011; Hanson et al., 2015; Luby et al., 2013) or greater volume and elevated response (Lupien et al., 2011; Mehta et al., 2009; Tottenham et al., 2010). Furthermore, those studies found greater volume and elevated response of amygdala (Mehta et al., 2009; Tottenham et al., 2010) were performed several years after the institutionalized children adopted by high socio-economic status families. These data suggested that EARE modified amygdala changes was resistant to recovery, and it was consistent with primate research that suggested abnormal behaviors was resistant to environmental enrichment treatments (Lutz et al., 2004; Lutz & Novak, 2005; Novak et al., 1998; Rommек et al., 2009a). Similarly, in adults exposed to EARE some studies

found no significant changes of amygdala volume (Bremner et al., 1997; Cohen et al., 2006), while others found larger volume (Evans et al., 2016; Lyons-Ruth et al., 2016), interrupted regulation of negative emotion (Kim et al., 2013), increased response to potential rewards (Casement et al., 2014), elevated amygdala responses to threat but not happy faces (Javanbakht et al., 2015). In addition to amygdala structure and activity changes, its connectivity with other brain regions was also affected (Barch et al., 2016; Jedd et al., 2015). Despite those controversial results, the influence of EARE on emotion such as the elevated response to emotion stimuli both in human and primates (Casement et al., 2014; Javanbakht et al., 2015; Nelson et al., 2009) should be mainly achieved through its influence on amygdala.

Hippocampus

Hippocampus, a major component of the brains located inside the medial temporal lobe and beneath the cortical surface, is involved in episodic, declarative, contextual, and spatial learning and memory, as well as being a component in the control of autonomic and vegetative functions (Buckley, 2005; Eichenbaum, 2001; Eichenbaum et al., 1992, 1996; Manns &

Eichenbaum, 2006; Opitz, 2014; Shohamy & Turk-Browne, 2013). In human studies, EARE induced significant reduction of hippocampal volume was an consistent finding in adults (Bremner et al., 1997; Cohen et al., 2006; Hart & Rubia, 2012; McCrory et al., 2011; Stein et al., 1997; Woon & Hedges, 2008). However, children and adolescents studies showed inconsistent results, with few found decreased volume (Edmiston et al., 2011; Hanson et al., 2015; Luby et al., 2013), while most found no significant change (Carrion et al., 2001; De Bellis et al., 2001, 1999, 2002; Mehta et al., 2009; Tottenham et al., 2010; Woon & Hedges, 2008). Primate studies also found no significant hippocampal volume change in PR (Sánchez et al., 1998; Spinelli et al., 2009) and repeated mother deprived (Law et al., 2009b) juvenile monkeys, suggesting changes of hippocampus seemed to happen later in life compared to early life amygdala changes. Two possible explanations could account for the discrepancy of children and adult findings. Firstly, that might due to the fact that the hippocampus develops mainly in the first years of life, therefore less affected by exposure to adversity in childhood and adolescence (Houston et al., 2014; Lenroot & Giedd, 2006; Richards & Xie, 2015). Another possibility is that EARE might not have an immediate effect on the hippocampus but induced changes over time, and long-term effects of EARE exposure may be delayed and became manifest only in later phases of development when the vulnerable brain reaches maturation (Andersen & Teicher, 2004; Brunson et al., 2005; Gluckman & Hanson, 2004; Gluckman et al., 2007; Sapolsky et al., 1985). Moreover, human studies found interesting results concerned with influence of EARE exposure on structure and activity of hippocampus and amygdala, with decreased hippocampal volume and activity in humans exposed to adulthood stress (Bremner et al., 2007; Lupien et al., 2007; Rauch et al., 2000) and adults experiencing EARE (Bremner et al., 1997; Cohen et al., 2006; Stein et al., 1997; Woon & Hedges, 2008), while increased amygdala volume and activity in humans exposed to adulthood stress (Bremner et al., 2007; Lupien et al., 2007; Rauch et al., 2000) and adults experiencing EARE (Mehta et al., 2009; Tottenham et al., 2010). Although the biological mechanism and meaning of this phenomenon remains unclear, that might contribute to or even be the direct reason for the impaired learning and memory ability (decreased hippocampal volume and activity related) and elevated response to emotional stimuli (increased amygdala volume and activity related) described above.

Prefrontal cortex

The prefrontal cortex (PFC) is the anterior part of the frontal lobes of the brain and implicated in planning complex cognitive behaviors, personality expression, decision making and moderating correct social behavior. Children and adolescents studies showed inconsistent results of EARE induced PFC structural changes, with findings of either no significant differences (De Bellis et al., 1999), or significantly smaller volume (De Bellis et al., 2002; Edmiston et al., 2011; Hanson et al., 2010; Morey et al., 2016; Thomaes et al., 2010) or significantly larger volume (Carrion et al., 2009; Richert et al., 2006). In contrast, decreased PFC volume in adults exposed to

childhood maltreatment was a consistent finding (Tomoda et al., 2009; van Harmelen et al., 2010). That might due to the fact that PFC continues to develop during adolescence (Houston et al., 2014; Lenroot & Giedd, 2006; Richards & Xie, 2015), therefore might be particularly vulnerable to the effects of stress during adolescence. In addition to the structural changes, EARE could also induce PFC functional changes, with some human adults exposed to EARE showing reduced prefrontal cortex activity during monetary reward anticipation and emotion regulation (Casement et al., 2015; Kim et al., 2013; Romens et al., 2015; Schweizer et al., 2016), while others showing increased response to potential rewards and threatening faces and in passive viewing conditions (Casement et al., 2014; Javanbakht et al., 2015; Jedd et al., 2015; Wang et al., 2016; White et al., 2015). One primate report indicated PR juvenile monkeys showed greater enlarged medial prefrontal cortex (mPFC) size (Spinelli et al., 2009). Moreover, both rodent and primate studies revealed the direct underlying epigenetic mechanisms of EARE on PFC through influencing differential gene expression, histone acetylation and DNA methylation (Blaze et al., 2015a; Provençal et al., 2012; Wall et al., 2012). Studies regarding EARE effects on PFC in primates are rare, and further investigations are necessary.

Other brain regions

The anterior cingulate cortex (ACC) is the frontal part of the cingulate cortex, and appears to play a role in a wide variety of rational cognitive functions, such as reward anticipation, decision-making, empathy and emotion (Devinsky et al., 1995; Drevets et al., 2008). It can be divided anatomically into dorsal and ventral components, with dorsal part connected with PFC making its involvement in cognition possible, and the ventral part connected with amygdala making its involvement in emotion possible (Bush et al., 2000; Morecraft et al., 2007). Human studies showed reduced volume of adult ACC in people with mood disorders (Botteron et al., 2002; Drevets et al., 1997; Yamasue et al., 2003), adults exposed to early life stress (ELS) (Cohen et al., 2006) and abuse-related Posttraumatic stress disorder (PTSD) (Kitayama et al., 2006; Thomaes et al., 2010) and major depressive disorder (Treadway et al., 2009). On the contrary, a primate study found enlarged ACC in PR juvenile monkeys (Spinelli et al., 2009). Moreover, an epigenetic study showed parental separations in infant marmoset affected expression of genes in the ACC of adolescent monkeys (Law et al., 2009a). Additionally, both human and primate studies revealed EARE affected cerebellum, with human studies showing decreased cerebellum (Bauer et al., 2009; Edmiston et al., 2011), while a primate study revealing larger cerebellar vermis area in PR juvenile monkeys (Spinelli et al., 2009). EARE effect on primate cerebellum might due to the fact that macaque cerebellum has high density of glucocorticoid receptors (GRs) (Sánchez et al., 2000), which put it particularly vulnerable to stress hormones related over stimulation. Striatum was another brain region affected by EARE, with increased response to potential rewards (Casement et al., 2014) and elevated dopamine responses to amphetamine (Oswald et al.,

2014), and a potential neurobiological mechanism linking early-life adversity and altered ventral striatal development was indicated (Goff & Tottenham, 2015). In addition to those specific regional changes, PR chimpanzees showed less global white-to-grey matter volume and cortical folding (Bogart et al., 2014). Structural connectivity between different brain regions was also affected by EARE, as studies showed affected corpus callosum, a wide and flat bundle of axons beneath the cortex connecting left and right cerebral hemispheres and facilitating inter-hemispheric communication, in a inconsistent way that most human studies showing EARE reduced corpus callosum (De Bellis et al., 1999; Rinne-Albers et al., 2016; Teicher et al., 2004, 1997), while few showing no significant changes (Mehta et al., 2009). Primate studies also found either decreased corpus callosum size (Sánchez et al., 1998) or no significant changes (Spinelli et al., 2009).

Neurons and glia cells

Neurons are the basic unit of brain. Neuronal network is responsible for the daily cognitive and emotional behaviors. Glia cell is a group of non-neuronal cells that support and protect the neurons in the brain. Rodent studies showed that maternal separation could induce morphological alteration of the apical dendrites of CA3 pyramidal neurons (Kwak et al., 2008); could increase corticotropin releasing factor (CRF)-containing neurons in amygdala (Becker et al., 2007); and could decrease *in vivo* firing activity of amygdala neurons (Adams & Rosenkranz, 2016) and sex related neurogenesis (Oomen et al., 2009). Chronic stress could induce atrophy of dendrites in hippocampus of rats (Brunson et al., 2005; McEwen, 1999) and tree shrews (Magariños et al., 1996), and could induce hippocampal neuroplasticity changes (Fenoglio et al., 2006). Bartesaghi and colleagues used guinea-pig as animal model to investigate the effects of early isolation on neurons, and they found that early isolation could induce morphologic changes of neurons in entorhinal cortex and hippocampus (Bartesaghi et al., 2003a, b; Bartesaghi & Serrai, 2001, 2004). Although primate studies found neuronal morphological changes in EARE exposed monkeys (Bryan & Riesen, 1989; Floeter & Greenough, 1979; Stell & Riesen, 1987; Struble & Riesen, 1978), these early findings were limited to cerebellum, somatosensory and motor cortex, with limited information on other important brain regions, such as hippocampus, amygdala and PFC. Recent studies showed that different environments could induce neuron plasticity changes in the key brain regions involved in learning and memory. Complex environment could enhance complexity of the dendritic tree and density of dendritic spine in hippocampus and PFC in monkeys (Kozorovitskiy et al., 2005). Early parental deprivation in the marmoset monkey could produce long-term changes in hippocampal expression of genes involved in synaptic plasticity and implicated in mood disorder (Law et al., 2009b). So these neuron morphological and plasticity changes might explain and account for how EARE take effects on cell level, and then further more leading to behavioral changes.

As EARE effects on glia cells, rodent studies revealed that EARE could induce long-term changes of astrocyte density and

numbers in many brain regions, including PFC, mPFC, hippocampus, cingulate cortex and amygdala (Leventopoulos et al., 2007), and could alter behavioral, autonomic and endocrine responses to environmental challenge (Musholt et al., 2009; Rüedi-Bettschen et al., 2006). Although there was no direct evidence pointing out that glia cell changes were responsible for those altered responses in rats, those studies at least suggested the involvement of glia cell in EARE induced effects. Moreover, human studies showed that glial cell depletion in many brain regions was related to mood disorders, as the number of glia cell was reduced in PFC of both major depressive disorder (MDD) and bipolar disorder (BD) patients (Öngür et al., 1998), in the amygdala of major depressive disorder patients (Bowley et al., 2002) and in anterior cingulate cortex of major depressive disorder and schizophrenia patients (Cotter et al., 2001). Considering the important trophic influence of glia on neurons, glia cell deficits induced by EARE could possibly be responsible for EARE effects on neurons and furthermore to abnormal behavioral function. If that is true, how does it happen? Rodent Studies showed that stress related hormone glucocorticoid receptors (GRs) were also expressed in glia cells (Bohn et al., 1991; Jung-Testas & Baulieu, 1998; Vielkind et al., 1990). Glucocorticoid is the product of the HPA axis, so EARE might take effects through its influence on stress related hormones, like glucocorticoid, and then exert influence on glia cells leading to various effects (Jauregui-Huerta et al., 2010). Indeed, *in vitro* and *in vivo* studies showed that glucocorticoids could influence gene expression in glia cells (Bohn et al., 1994; Kumar et al., 1985) and could regulate the concentration of glial fibrillary acidic proteins (O'Callaghan et al., 1989). By playing central roles in learning and memory, hippocampal astrocyte number was dose-dependently increased by corticosterone treatment (Bridges et al., 2008), and glial responses in hippocampus was also regulated by glucocorticoid through influencing gene expression (Nichols et al., 2005). However, few studies were performed to investigate this issue and EARE affected glia cell changes were link directly to behavioral outcomes without solid evidence. In primate studies, there are lack of evidence to support that EARE affects glia cell structural and functional changes, and furthermore, induces behavioral outcomes.

Lateralisation

Some studies suggested that the influence of EARE on different brain hemisphere might be different, and different type of EARE might take effects differentially on the same brain structure. A human study found that the institutionalized children had greater right amygdala volume, while the left amygdala volume was smaller in the children experienced longer periods of deprivation (Mehta et al., 2009). Another human study showed that patients with child abuse-related complex PTSD showed reduced gray matter concentration in right hippocampus and right dorsal ACC, but not in the left areas (Thomaes et al., 2010). In primate studies, maternal separation was associated with activation in the right dorsolateral PFC and decreased activity in the left dorsolateral PFC of juvenile rhesus monkeys (Rilling et al., 2001). Not only brain structure and function

showed lateralisation affection by early experiences, behavioral research also found lateralisation in primates, as an interesting research showed that PR monkeys demonstrated greater left-hand bias compared to MR reared monkeys (Bennett et al., 2008). The number of lateralisation related studies is limited and the underlying mechanism remains unknown, which certainly adds complexity to the understanding of the influence EARE on brain structure and functional changes and the related abnormal behavioral outcomes.

Other EARE effects

Young animals are particularly vulnerable to EARE effects

Adverse experience has its influence over all life stages, including early, middle and later life, in which infants are especially vulnerable to EARE and the consequences could be persistent into later life. That might be due to the fact that the most sensitive period of the whole life is the early stage, during which the body is undergoing profound physiological development, such as HPA axis, and brain is also undergoing profound neural development, such as neurogenesis. The amygdala develops rapidly during the early postnatal period in animals, e.g., in rats, cats and primates (Kikusui & Mori, 2009; Lupien et al., 2009; Payne et al., 2010; Wakefield & Levine, 1985). Stress related hormones and receptors were also maximally expressed in the brain early in development (Avishai-Eliner et al., 1996; Baram & Hataalski, 1998; Meaney & Szyf, 2005; Pryce et al., 2005a; Vazquez et al., 2006). These early physiological development heighten the vulnerability of the brain to environmental exposures. On the other hand, the proper development needs proper environmental stimuli, and the natural and best stimuli during early life is the attachment between caregivers, especially mothers, and infants, as mothers could supply tactile contact, physical warmth, nourishment, and psychological comforts. As stated in attachment theory and affectional system, infants need to develop a stable relationship with the mother for social and emotional development to occur normally, while various EARE intervene the forming of the bonds of this relation, therefore both short-term and long-term devastating influence are inevitable.

Sexual differences in EARE effects

Human studies showed that affectability of various mental disorders were sex-related during development, with boys showing higher tendencies to develop aggression and novelty seeking behaviors (Farrington & Loeber, 2000) while girls more susceptible to anxiety and depression (Kessler, 2003). Additionally, EARE influence might also be sex related, e.g., corpus callosum volume reduction was only found in EARE exposed males (De Bellis et al., 1999). Similarly, animal studies also revealed the vulnerability of males to EARE in rodents (Galea et al., 1997; Kikusui & Mori, 2009) and primates (Clarke, 1993; Cross & Harlow, 1965; Mitchell, 1968; Rommeck et al., 2009a; Suomi et al., 1971). On the contrary, other studies showed preference of EARE on females in humans (Heim & Nemeroff, 2001; Klimes-Dougan et al., 2001), rodents (Hoyer et al., 2013; Ziabreva et al., 2003b) and primates (Sánchez et al., 2005). Previous studies showed that stress could induce

decreasing in number and length of apical dendritic branch of medial prefrontal cortex in male rats, whereas increasing in apical dendritic length in female rats (Garrett & Wellman, 2009). Isolated males showed less dendritic branches, shorter dendritic length and smaller dendritic spine density than control males, while isolated females had more dendritic branches than control females in guinea pig (Bartasaghi et al., 2003a). Neurogenesis was significantly increased in male but decreased in female offspring after maternal deprivation in rats (Oomen et al., 2009). The mechanism of those sexual differences remains unclear, but one possible explanation is the gender related physiological differences, such as neuroendocrine system and brain structure and function, which may induce different behavioral and physiological responses in male and female subjects.

Time effects of EARE

Early life is a time of heightened susceptibility to EARE and expression of adverse experiences induced effects would be different across life time, therefore the time of administration of adverse experiences and subjects age of measurement might partially explain the discrepant findings across studies (Tottenham & Sheridan, 2009).

The time of adverse experiences administration is important, as different brain regions might have unique windows of vulnerability to stress, e.g., human studies indicate that the time window of hippocampus, corpus callosum and frontal cortex is at ages of 3-5, 9-10 and 14-16 years, respectively (Andersen et al., 2008). Rodent studies revealed the critical importance of specific time windows early in life for the outcome of maternal separation (Bock et al., 2005; Gos et al., 2008; Pryce et al., 2005b). Early primate studies by Harlow et al. showed the importance of administration time of adverse experiences (isolation), in a way that isolation beginning at birth generated most severe effects and persisting abnormalities (Harlow et al., 1965; Mitchell, 1968), while the isolation starting until later in life would produce less severe effects and persistent abnormalities (Harlow et al., 1965; Mitchell, 1968). Moreover, different lasting period of EARE also produced different effects even was all initiated at birth, i.e., 3 months isolation only induced reversible debilitating behavioral deficits, while at least six months isolation generated most severe effects and persisting abnormalities; 3 months isolation induced least, 6 months isolation induced moderate and 12 months isolation induced most severe defects (Griffin & Harlow, 1966; Harlow et al., 1965; Mitchell, 1968). These studies suggested that both the time point of administration of EARE and the lasting period have different influences on behavioral and biological outcomes.

Human studies showed different, or even contrary effects of EARE in children and adults, suggesting EARE might induce differential outcomes across lifespan. For example, childhood abuse induced significant reduction of hippocampal volume in adults (Bremner et al., 1997; Cohen et al., 2006; Stein et al., 1997; Woon & Hedges, 2008) but not in children (Carion et al., 2001; De Bellis et al., 2001, 1999; Woon & Hedges, 2008); EARE induced hypercortisolism in children (Essex et al., 2002;

Fernald & Gunnar, 2009; Flinn & England, 1997; Kaufman et al., 1997) but hypocortisolism in adults (Carpenter et al., 2009; Elzinga et al., 2008); adults with abuse related PTSD showed ACC volume reductions (Kitayama et al., 2006; Thomaes et al., 2010), whereas pediatric PTSD showed increased ACC (Richert et al., 2006). Primate studies also found similar results, e.g., monkeys exposed to EARE showed less aggression during infancy (Chamove et al., 1973; Harlow et al., 1965) but more aggression during latter life (Chamove et al., 1973; Mitchell, 1968), whereas showed more affiliative behavior during infancy (Chamove et al., 1973; Rosenblum & Pauly, 1984; Ruppenthal et al., 1991) but less during adulthood (Kraemer & McKinney, 1979; Rosenblum & Pauly, 1984; Winslow et al., 2003). Monkeys exposed to EARE showed more activity during infants (Champoux et al., 1991) but less activity during adulthood (Harlow & Suomi, 1971a; Mason & Sponholz, 1963; Mitchell, 1968); The number and style of stereotypies exhibited in monkeys also varied by age, e.g., the number of whole-body stereotypies were negatively correlated with age, whereas self-directed stereotypies were positively correlated; moreover younger monkeys exhibited

more pacing, body-flipping, and swinging, while older ones exhibited more hair-pulling and saluting (Lutz et al., 2003). These studies showed different, or even opposite effects of EARE on behavioral and biological outcomes between infants and adults, indicating EARE induce different outcomes across lifespan.

MECHANISMS UNDERLYING EARE INDUCED EFFECTS

Neuroendocrinological mechanisms

Some recent study linked behavioral outcomes with EARE affected neuroendocrine systems, and suggested that EARE might modulate subsequent social behaviors through regulating both the production and body's sensitivity to neurotransmitters and hormones (Cushing & Kramer, 2005). Moreover, studies indicate that the involved neurotransmitters and hormones were mainly monoamine neurotransmitter serotonergic systems, including serotonergic system and catecholamine system (both noradrenergic system and dopaminergic system), and glucocorticoid hormones (cortisol in non-human primates and humans), oxytocin and growth hormone (GH)(Table 4).

Table 4 EARE induced effects on neuroendocrine systems

	Outcome		Primate studies	Human studies
Serotonin system		PR and SPR monkeys showed decreased CSF levels of 5-HIAA	Fahlke et al., 2000; Higley et al., 1996a; Maestripieri et al., 2006; Shannon et al., 2005	
		PR monkeys showed decreased SERT binding potential	Ichise et al., 2006	
		VFD monkeys were hyporesponsive to the serotonergic probe mCPP	Rosenblum et al., 1994	
Catecholamine system		PR monkeys showed lower CSF concentrations of HVA and attenuated NE secretion	Clarke et al., 1999; Clarke et al., 1996	
		VFD monkeys were hyper responsive to the noreadrenergic probe yohimbine	Rosenblum et al., 1994	
		PR monkeys showed significantly lower DOPAC concentrations	Clarke et al., 1999; Clarke et al., 1996	
HPA axis dysregulation	Hypocortisolism	Hormone level	Barrett et al., 2009; Coplan et al., 2005; Essex et al., 2002; Flinn Coplan et al., 1996; Coplan et al., 2001; & England, 1997; Gunnar et al., 2001; Kaufman et al., 1997	
		Increased HPA response	Fahlke et al., 2000; Higley et al., 1992; Heim et al., 2000b; Kikusui Kraemer et al., 1983, 1984; Sánchez et al., 2005; Suomi, 1991	
	Hypocortisolism	Hormone level	Clarke et al., 1998; Capitanio et al., 2005; Shannon et al., 1998	
		Decreased HPA response	Barr et al., 2004; Capitanio et al., 2005; Carpenter et al., 2009; Clarke, 1993; Dettling et al., 1998; Dettling Elzinga et al., 2008; Hart et al., 2002a, b; Lyons et al., 2000; Parker et al., 1995 et al., 2004	

Monoamine and hormone systems

The serotonergic system was shown to moderate the effects of EARE on the risk of depression in humans (Eley et al., 2004; Kaufman et al., 2004), and primate studies also indicate the role

of serotonin system in regulating the effects of EARE. Maternal rejected, PR and SPR reared infant monkeys exhibited lower CSF 5-HIAA concentrations (Fahlke et al., 2000; Higley et al., 1996a; Maestripieri et al., 2006; Shannon et al., 2005); PR monkeys showed decreased SERT binding potential across a

range of brain areas (Ichise et al., 2006); VFD reared monkeys were hyporesponsive to the serotonergic probe meta-Chlorophenylpiperazine (mCPP) (Rosenblum et al., 1994). Moreover, epigenetic studies also indicate the role of serotonin system plays in EARE induced HPA axis dysfunction (Barr et al., 2004; Rosenblum et al., 1994; Shannon et al., 2005; Spinelli et al., 2007) and subsequent abnormal behavioral outcomes (Barr et al., 2003, 2004; Law et al., 2009b; Maestriperi et al., 2006; Vicentic et al., 2006; Ziabreva et al., 2003a). Many studies showed catecholamine system is another candidate through which EARE takes its effect. PR monkeys showed attenuated Norepinephrine (NE) secretion (Clarke et al., 1999, 1996) and reduced CSF concentrations of catecholamine metabolite (Clarke et al., 1999, 1996), while VFD reared monkeys were hyper responsive to the noreadrenergic probe yohimbine (Rosenblum et al., 1994). It was further suggested that EARE might influence the differentiation of noradrenergic neurons and thus alter HPA responses stress during adulthood (Liu et al., 2000). Dopamine system (another catecholamine system) might also be involved in EARE effects, as significantly lower concentrations of dopamine metabolite were revealed in PR infant monkeys (Clarke et al., 1999, 1996) and history of childhood adversity was positively associated with striatal dopamine responses to amphetamine (Oswald et al., 2014).

Additionally, primate studies indicate potential hormonal pathways through which EARE takes effects, including oxytocin, growth hormone (GH), and most importantly, cortisol. Monkeys exposed to EARE showed abnormal aggressive and affiliative behaviors, and oxytocin was suggested to be a neuropeptide for affiliation and involved in the regulation of social bonding behaviors (Insel, 1992; Lim & Young, 2006). Therefore, oxytocin is a possible pathway for EARE to take effects, which indeed was probed by Winslow et al. (Winslow et al., 2003), showing that the decrease in affiliative behavior in PR rhesus monkeys was significantly and positively correlated with cerebrospinal oxytocin. Another hormone, GH, was also related to early adversity, as PR and social separation experiences in infant monkeys showed abnormal GH levels (Champoux et al., 1989a; Laudenslager et al., 1995). Most importantly, the main target of EARE under investigation is HPA axis. While some studies showed blunt HPA response, and thus decreased cortisol and ACTH levels (Barr et al., 2004; Capitanio et al., 2005; Clarke, 1993; Dettling et al., 1998, 2002a, b; Lyons et al., 2000; Parker et al., 2004), others showed the opposite (Barrett et al., 2009; Coplan et al., 2005, 1996, 2001; Suomi, 1991). Although consistent results were not achieved, the importance of HPA dysregulation in EARE induced effects was suggested.

Hypothalamic-pituitary-adrenal (HPA) axis dysregulation

EARE is associated with elevated levels of stress and fear. The adverse impact of stress on brain development was suggested to be largely through hypothalamic-pituitary-adrenal (HPA) axis both in humans (Loman & Gunnar, 2010) and primates (Sanchez, 2006). The effects of EARE on HPA circadian rhythmicity and the function of HPA axis were reviewed in this section.

Circadian rhythmicity

The HPA axis is a complex set of direct influences and feedback interactions among three endocrine glands, i.e., hypothalamus, pituitary gland, and adrenal glands. Under basal conditions, HPA axis exhibits a circadian rhythmicity with a peak around the time of waking and a trough during the quiescent time of the activity cycle (Dickmeis et al., 2013; Leliavski et al., 2015; Tsang et al., 2016, 2014). So cortisol levels typically follow the circadian rhythm with levels highest occurring about 20 minutes after awakening in the morning (cortisol awakening response, CAR) and declining throughout the day. Alterations in the normal pattern of HPA rhythmicity, including CAR response and diurnal decrease of cortisol, were found in human studies. Most studies found higher morning cortisol level than controls in maltreated children (Cicchetti & Rogosch, 2001; Cutuli et al., 2010) and EARE exposed adults (Gonzalez et al., 2009; Gustafsson et al., 2010), while some found lower morning cortisol level (Carlson & Earls, 1997). Moreover, different kind of EARE might have differential influence on morning cortisol values, as studies indicate more emotionally and sexually abused children showed higher morning cortisol values, whereas more severe physically neglected and abused children showed lower levels (Bruce et al., 2009; Cicchetti & Rogosch, 2001). Additionally, EARE exposed children also showed higher incidences of atypical diurnal rhythmicity patterns, such as a peaking in the afternoon or evening (Cicchetti et al., 2010; Dozier et al., 2006). Similarly, abnormal HPA circadian rhythmicity were also found in limited amount of primate studies on rhesus monkeys, with morning peak occurring late in PR infants (Thomas et al., 1995) and flattened diurnal rhythm in repetitive maternal separation exposed infants (Sánchez et al., 2005). However, a recent study found no shift in diurnal patterns of cortisol in PR reared juvenile rhesus monkeys (Barrett et al., 2009). Although how EARE induces those abnormal HPA axis circadian rhythmicity, and its different or even contrary effects remains unknown, these HPA axis circadian rhythmicity abnormalities certainly contribute to various abnormal behavioral outcomes.

HPA axis dysregulation

In humans, the HPA axis develops over the initial several years of life and is highly sensitive to EARE (De Weerth et al., 2003; Watamura et al., 2004). The key elements of the HPA axis are as following: the hypothalamus synthesizes and secretes corticotropin-releasing hormone (CRH); CRH stimulates the secretion of adrenocorticotrophic hormone (ACTH) in pituitary gland; ACTH acts on the adrenal cortices, which then produces glucocorticoid hormones (mainly cortisol in NHPs and humans); glucocorticoids in turn act back on the hypothalamus and pituitary to suppress CRH and ACTH production in a negative feedback cycle. When activated in response to a stressor, the HPA axis participates in a cascade of neuroendocrine responses, and a typical HPA stress response involves a period of increased glucocorticoids in circulation induced by stimulation of elevated levels of CRH and ACTH, followed by a return to baseline levels induced by negative feedback of glucocorticoids

(Herman & Cullinan, 1997). Thus CRH, ACTH and glucocorticoids levels could indicate the reactivity levels of HPA axis, with CRH an important neurotransmitter in HPA axis to initiate the autonomic and behavioral changes in response to stress (Heinrichs et al., 1995; Krohg et al., 2008; Ohmura & Yoshioka, 2009; Smagin & Dunn, 2000). Studies showed EARE induced elevated cerebrospinal fluid (CSF) concentrations of CRH levels in mother deprived rats (Ladd et al., 1996) and VFD reared infant monkeys (Coplan et al., 1996). Not only the CRH levels was increased, a study showed that EARE increased the density of CRH binding sites in many brain regions, including PFC cortex, amygdala and hippocampus (Anisman et al., 1998). As analysis of CRH requires sampling of CSF, it was hard to perform the experiment on healthy humans. On the other hand, analysis of ACTH and glucocorticoids (cortisol) only requires blood or urine sampling, so they are more widely investigated in humans.

Glucocorticoids was revealed to be released from the adrenal cortex during neuroendocrine responses to stress (Herman et al., 2003, 1996), and then regulate HPA axis via negative feedback by binding to two types of receptors, mineralocorticoid receptors (MRs) with high affinity (important in proactive maintenance of HPA basal activity), and GRs with low affinity (primarily responsible for negative feedback). Glucocorticoids could pass through the blood-brain barrier to influence brain function (Zarrow et al., 1970), and MRs' expression was significantly greater in monkey infants than other ages (Pryce et al., 2005a). Therefore, HPA axis was one of the major pathways through which EARE induces stress and shapes brain development, particular in infants. Glucocorticoids could facilitate HPA axis activation by occupying its receptors in amygdala, leading CRH increase within amygdala (Kolber et al., 2008), whereas it could also suppress HPA axis by occupying its hippocampal receptors (van Haast et al., 1997). Amygdala and hippocampus are important brain regions for socio-emotional functioning and learning and memory throughout development, and they have a high density of receptors for unbound glucocorticoids, therefore are major targets of EARE affected HPA axis (Johnson et al., 2005; Sánchez et al., 2000). Additionally, EARE could affect HPA axis function bidirectionally, with some studies showing attenuated basal and challenge induced levels of cortisol (hypocortisolism), while others showing elevated levels in both conditions (hypercortisolism).

Hypercortisolism

Human studies showed that EARE could induce hypercortisolism of basal cortisol level in children, reflected by elevated levels of cortisol (Essex et al., 2002; Flinn & England, 1997; Gunnar et al., 2001) and ACTH (Kaufman et al., 1997) in EARE exposed children. Primate studies also showed EARE induced hypercortisolism, reflected by increased plasma cortisol and ACTH in PR infants and juvenile monkeys (Barrett et al., 2009; Suomi, 1991). The elevated cortisol levels in hairs of PR infants indicate the long time accumulation of EARE outcomes (Dettmer et al., 2012). Increased basal cortisol levels were found to be induced by prenatal stress (Pryce et al., 2011).

Persistently elevated CSF concentrations of CRF in both infants and mothers under VFD conditions were reported (Coplan et al., 2005, 1996). Not only EARE could affect infants and children, it was suggested that childhood abuse was associated with a persistent sensitization of the HPA axis to stress in human adults (Elzinga et al., 2008), e.g., adults exposed to EARE had higher HPA reactivity during the Trier Social Stress Test (TSST) (Heim et al., 2000b; Pesonen et al., 2010). Animal studies also showed hyper-response of HPA axis activity when facing stress in both infants and adults exposed to EARE. Rodent studies revealed higher HPA response to novelty stress in early-weaned mice (Kikusui & Mori, 2009). Primate studies showed hyper-responsiveness in EARE reared monkeys, reflected by increased cortisol response to stress in monkeys exposed to PR rearing (Fahlke et al., 2000; Suomi, 1991), VFD rearing (Coplan et al., 2001), repetitive maternal separation (Sánchez et al., 2005) and parental deprivation (Higley et al., 1992; Sánchez et al., 2005). Amphetamine challenge test also revealed neurochemical and behavioral hyper-responsiveness in isolated monkeys (Kraemer et al., 1983, 1984). All those studies suggested EARE induced hypercortisolism, reflected by elevated basal and stress or challenge facing levels of cortisol, ACTH or CRF.

Hypocortisolism

EARE induced hypocortisolism was also a common finding (Gunnar & Vazquez, 2001), e.g., maltreated children showed decreased basal levels of cortisol (Brand et al., 2010; Heim et al., 2000a) and ACTH (De Bellis et al., 1994). Primate studies revealed attenuated basal levels of cortisol and ACTH in PR monkeys (Capitanio et al., 2005; Clarke et al., 1998; Shannon et al., 1998). When facing stress or challenge, EARE exposed children and adults both showed blunt cortisol response and thus reduced cortisol level (Carpenter et al., 2009; Elzinga et al., 2008; Hart et al., 1995). Similarly, primate studies showed blunt HPA responses during stress and thus decreased cortisol and ACTH levels in PR reared infants (Barr et al., 2004; Capitanio et al., 2005; Clarke, 1993), in young adults exposed to maternal deprivation and intermittent separation (Capitanio et al., 2005; Lyons et al., 2000) and in the hairs of PR infants (Feng et al., 2011) of rhesus monkeys. EARE induced hypocortisolism was also found in other monkey species, including maternal neglect exposed juvenile Goeldi's monkeys (Dettling et al., 1998), intermittent stress exposed squirrel monkeys (Parker et al., 2004) and parental deprivation exposed marmosets (Dettling et al., 2002a, b). All those studies indicate hypocortisolism reflected by decreased basal and stress or challenge facing levels of cortisol or ACTH.

As described above, it is controversial as to the effects of EARE on HPA axis, with some studies showing hypercortisolism while other showing hypocortisolism. There are several possible reasons. Firstly, different types of EARE vary between different research, and even for a same type of EARE, the procedures, manipulations, tests and measuring indexes could be different in different experiments. Secondly, different genotype among human races or animal species could contribute to the divergence as well, in a way that individuals with certain

genotype may be more sensitive to a particular type of EARE than others. In addition, subjects' personality or temperaments could also partially contribute to the divergence, e.g., children with inhibited temperaments tended to have higher cortisol levels than extroverted children (Gunnar et al., 1995; Kagan et al., 1988), indicating that long-term consequences of EARE may not uniform across subject populations..

A sample of EARE induced neurotransmitter and hormonal changes related behavioral outcomes - social status of primates

Studies suggest that EARE could induce abnormal changes of neurotransmitters and hormones and then influence social status of primates. Serotonergic system was the most widely studied neurotransmitter involved. Primate studies showed that different levels of CSF serotonin (5-HT) or its main metabolite 5-Hydroxyindoleacetic acid (5HIAA) were related to different social status, with higher levels related to more dominant status (Higley et al., 1996b; Raleigh et al., 1983). Additionally, serotonergic drugs were found to be able to influence dominance status, in a way that serotonergic enhancing drugs increase social dominance while serotonergic reducing drugs decrease dominance (Raleigh et al., 1991). 5-HT seems to be positively related to social dominance status, and studies suggested that might due to its influence on affiliative and aggressive behaviors which are important factors in dominance formation and maintenance. Primate studies showed positive correlation between CSF 5-HIAA and affiliative approaching and grooming behavior (Mehlman et al., 1995; Raleigh et al., 1985) and negative correlation between CSF 5-HIAA and aggressive behavior (Higley et al., 1996a), which were consistent with the previous presumption that affiliative behavior was much more effective in acquiring and reinforcing social dominance than aggressive behaviors. Supporting evidence also came from another genetic primate study that suggested certain serotonin transporter (5-HTT/SERT) diplotypes might modulate acquisition of dominance status (Miller-Butterworth et al., 2007). Beside serotonergic system, dopaminergic systems might also affect social dominance status, as dopamine transporter (DAT) gene variants were suggested to be associated with social rank in cynomolgus monkeys (Miller-Butterworth et al., 2008). As to hormones, although a study showed that cortisol concentration was significantly higher in dominant monkeys (Czoty et al., 2009), most studies failed to find the relationship between cortisol level and social rank (Czoty et al., 2009; Goo & Sassenrath, 1980; Morgan et al., 2000; Stavisky et al., 2001). Those studies suggested that EARE could influence social status of primates through its influence on neurotransmitters and hormones. The effects of EARE should not be limited on social status but also might on some other abnormal behaviors.

Genetic and epigenetic influences of EARE effects

Developing is a dynamic process involving constant and reciprocal interactions between organisms and the environments. Emerging evidence suggests that epigenetic modifications may serve as a critical mechanism through which experiences occurring during the lifespan can have sustained

effects in developmental outcomes (Daskalakis et al., 2013). Epigenetics refers to the study of inherited changes in phenotype (appearance) or gene expression caused by mechanisms other than changes in the underlying DNA sequence, such as modifications of transcription of the genome by chemical markers regulation, and variation in gene expression rather than gene sequence is the key concept. Moreover, epigenetics is used to describe the dynamic interactions between genome and the environment (Jablonka & Lamb, 2002). Research suggested that environmental events can modify the epigenetic status of the genome by activating intracellular pathways to regulate interaction between transcription factors and their DNA binding sites, leading to changes in gene expression and eventually different levels of proteins (Bagot & Meaney, 2010; Zhang & Meaney, 2010). This is the biological basis for the interplay between environmental factors and the genome in the regulation of individual differences in behaviors and cognition. Both animal and human studies suggest that EARE can lead to lasting changes in neurotransmitter systems and brain function, and then induce cognitive and behavioral changes. However, there was remarked inter-individual variations in responses to adversity (Collishaw et al., 2007; Rutter, 2007), and these variations might be due to different genotype, different living environment and interaction between the genome and environment.

Genetic influences

Different genotype could induce different behavioral outcomes. Allelic variation of the monoamine oxidase A (MAOA) gene was implicated in aggressive behaviors (Volavka et al., 2004). Both human (Caspi et al., 2002; Craig, 2005; Kim-Cohen et al., 2006) and primate (Karere et al., 2009) studies showed that genotype conferring low MAOA activity was related to mental health problems. These findings may partially explain the variability in developmental outcomes associated with maltreatment, e.g., why not all victims of maltreatment grow up to with abnormal behaviors like antisocial problems, and they provide epidemiological evidence that genotypes can moderate children's sensitivity to environmental insults. Similar results was revealed in 5-HTT genotype, as short promoter region of the serotonin transporter (5-HTTLPR) allele was related to increased anxious behavior in primates (McCormack et al., 2009) and highest emotional problem scores in human (Kumsta et al., 2010). Those evidences suggest the importance of genotype in behavioral outcomes.

Epigenetic influences of EARE on gene expression

Environmental and life experience could exert influences on gene expression and time course analysis indicate that maternally induced epigenetics might emerge during the postnatal period and could sustain into adulthood (Weaver et al., 2004). Epigenetic regulation of gene expression is particularly important during the early stages of development, and it is one of the main mechanisms mediating the long-term effects of maternal care on development (Champagne, 2008; Champagne & Curley, 2009; Diorio & Meaney, 2007; Meaney, 2001; Zhang et al., 2006). For example, rodent studies showed that postnatal

maternal licking/grooming (LG) behavior could induce increased hippocampal GR expression (Caldji et al., 1998; Francis et al., 1999; Liu et al., 1997; Weaver, 2007), while low levels of LG during neonates led to reduced expression of estrogen receptor in hypothalamus and reduced response to estrogen (Champagne et al., 2001, 2006, 2003). As to the effects of EARE on gene expression, isolation attenuated social interaction induced gene expression in rodents (Ahern et al., 2016; Lukkes et al., 2012, 2013; Shishkina et al., 2015; Wall et al., 2012). A human study also showed EARE related down regulation of genes containing GR response elements (Miller et al., 2009). In primate studies, early maternal separation could lead to gene expression changes in many brain regions, including differential expression of gene GUCY1A3 in amygdala, decreases in hippocampal growth associated protein 43 (GAP-43) mRNA and 5-HT receptor mRNA (Law et al., 2009b) and a selective long-term effect on expression of genes in ACC (Law et al., 2009a). Moreover, epigenetics is not a binary response across the whole brain. Different genes in different brain regions can be affected in different ways, e.g., early maternal deprivation could either induce reduction of gene expression (Liu et al., 1997; Rocerri et al., 2002) or up-regulation of gene expressions (Plotsky et al., 2005; Ziabreva et al., 2000).

Epigenetic influences of EARE on neurobiological and behavioral outcomes

Epigenetic influences on gene expression may lead to different expression patterns of proteins, thus different levels of hormones and neurotransmitters, ultimately lead to different behavioral outcomes. Studies suggested that EARE might exert its effects on behavioral outcomes independent of genotype. Primate studies revealed the importance of environment and life experience independent of genotype. MR monkeys showed significantly up-regulated level of 5-HTT during maternal separation, while NR monkeys did not (Kinnally et al., 2008). With the same low-activity Monoamine oxidase A (MAOA) genotype, MR reared monkeys were more aggressive than the PR monkeys (Newman et al., 2005). Higher 5-HTT cytosine-phosphate-guanosine (CpG) methylation, but not rh5-HTTLPR genotype, exacerbated the effects of early life stress on behavioral stress reactivity in infant monkeys (Kinnally et al., 2010). Additionally, from behavioral perspective alone, infant monkeys exposed to mother abuse showed the same tendency to their offspring, regardless of whether they were reared by their biological mothers or by foster mothers (Maestripieri, 2005). Rodent studies also supported the important role of environment and life experience on behavioral outcomes. Rodent studies suggested that maternal care behaviors especially postnatal maternal LG could be transmitted from the mother to her female offspring, so female offspring who received low levels of LG also provided low levels of this form of maternal care to their own offspring (Fleming et al., 2002). Cross-fostering studies in rodents indicate that this intergenerational transmission of behaviors was the result of early experience rather than genetic inheritance (Champagne & Meaney, 2001). For example, the biological offspring of low-LG mothers reared by high-LG dams resembled the normal

offspring of high-LG mothers (Francis et al., 1999). All those studies indicate the importance of environment and life experience, especially maternal interactions on the subsequent expression of behaviors, rather than the genetic contributions. Moreover, primate studies showed the importance of the interaction between genetic factors and environmental experience on neurobiological outcomes, as CSF 5-HIAA concentrations were significantly influenced by genotype in the PR but not MR reared monkeys (Bennett et al., 2002), and 5-HTT gene variation affected HPA axis activity in response to stress in a way that cortisol levels increased during separation in MR but decreased in PR monkeys (Barr et al., 2004).

Transgenerational epigenetic programming

Among the epigenetic mechanisms of EARE, such as DNA methylation, histone modifications, and micro-RNA expression, DNA methylation was the most intensively studied epigenetic phenomenon (Babenko et al., 2015; Blaze et al., 2015b; Jawahar et al., 2015; Provençal et al., 2015; Vaiserman, 2015a, b; Zheng et al., 2014). Actually, not only EARE exposure of the offspring themselves could lead to long time biological and behavioral outcomes, EARE exposure of parents could also influence their offspring, which was defined as transgenerational epigenetic programming phenomenon and has drawn much attention recently. Related studies suggested the underlying epigenetic mechanisms of maternal transgenerational influence to be DNA methylation, histone modifications, and micro-RNA expression (Babenko et al., 2015; Bale, 2014; Blaze & Roth, 2015; Gröger et al., 2016; Miska & Ferguson-Smith, 2016; Nagy & Turecki, 2015).

In addition to maternal influences, fathers can exert influences on offspring development either through direct care in living social environment, or indirectly through interacting with maternal influences. Some recent studies showed the important influence of paternal early experiences on infant development through non-social mechanisms, even in the absence of direct contact with offspring. This emerging field focuses on how environmental influences can epigenetically alter paternal sperm DNA methylation, histone modification and micro-RNA expression, and ultimately change the phenotype and behavior of offspring (Braun & Champagne, 2014; Curley et al., 2011; Day et al., 2016; Kinnally & Capitanio, 2015; Rodgers et al., 2013; Yuan et al., 2016).

DISCUSSION

Rehabilitation

Human studies indicate the possibility of rehabilitation of EARE induced deficits, e.g., Fisher et al. (2000, 2006, 2007) suggested that the improvements of caring following EARE had the potential to prevent or reverse EARE induced HPA axis dysfunction, such as normalizing perturbed diurnal cortisol patterns and reducing basal salivary cortisol level (Fernald & Gunnar, 2009). In primate studies, total social isolation was once considered to induce permanent defect, which was described as learning deficit in some studies, because isolated monkeys were lack of physical interactions and had no

opportunity for social learning with conspecifics, or to gradually develop sophisticated social behaviors (Mitchell, 1968; Sackett, 1969; Suomi et al., 1974). These abnormalities might be rehabilitated by socializing the isolate monkeys with conspecifics. Some studies reported that after the isolated monkeys was paired with “therapist” monkeys, less self-directed disturbance activities, or stereotypic behaviors, but more social contact and exploratory behaviors were observed (Harlow & Suomi, 1971b; Suomi, 1973). Less severe self-injurious behaviors were found when isolated monkeys were reared with surrogates (Brunelli et al., 2014; Harlow & Suomi, 1971b) or social housing (Lutz & Novak, 2005). Additionally, environmental enrichment treatments were used to eliminate abnormal behaviors and to normalize the behavioral repertoire of EARE exposed monkeys (Lutz & Novak, 2005; Novak et al., 1998; Rommeck et al., 2009a). It was suggested that environmental enrichment devices could only ameliorate less severe forms of abnormal behavior but not more severe forms of self-injurious or non-injurious self-abuse behaviors (Rommeck et al., 2009a). Those studies indicate rehabilitation is possible, at least partially, but it requires combination of multiple rehabilitation methods, such as socializing, environmental enrichment, and considerable time and effort.

Establishing NHP mental disorder models with EARE methods

EARE exposed NHPs showed signs of various mental disorders, including anxiety, autism and depression etc., making it a potential animal model to study human mental disorders, among which depression model was the mostly investigated one (Gilmer & McKinney, 2003; Pryce et al., 2005b; Worlein, 2014). The influence of EARE on NHPs is through daily life and accumulates over a period of time, making it a more natural model of mental disorders than those induced by drugs or invasive surgeries. The diagnosis of mental disorders in humans usually depends on questionnaire investigation and verbal communication between patients and doctors, which are not doable in NHPs. Therefore NHPs studies usually include daily group living or single subject observation and behavioral analysis, biochemical index analysis (e.g., hormones), brain structural and functional changes analysis by using modern imaging methods. However, due to the complexity of mental disorders, it is very difficult to diagnose mental disorders in NHPs. Different disorders might show very similar behavioral symptoms and biochemical abnormalities, therefore additional indexes are necessary in diagnosing. For example, in NHPs, anxiety and depression share symptoms of stereotypic behaviors and elevated cortisol level in response to stressor, so, more depression specific symptoms, such as lacking of responses to stimulus are needed for diagnosing. It is difficult to differentiate if a monkey was depressed or autistic as in both cases, preference of staying away from social activity and being alone in a corner would be shown. Certain mental disorders could be divided into many subtypes, e.g., depression includes unipolar, bipolar and atypical depression, etc, which certainly adds more complexity in establishing NHP animal models. These might explain the fact that despite being an ideal and

irreplaceable animal model, studies on EARE induced NHP mental disorder models are limited.

CONCLUDING REMARKS

In summary, as an irreplaceable animal model, NHP EARE experiments were performed for over 60 years and revealed important insights into understanding the effects of EARE on development and underlying mechanisms of related physiological and psychological diseases. Although much has been learned to date, there is much more to understand about EARE impact on developmental trajectory. Now, with the help of emerging cutting edge technologies, such as new brain imaging method, gene modification, optogenetics, etc, future EARE studies will further clarify these issues and help to cure the diseases.

ACKNOWLEDGEMENTS

The author wishes to thank Prof. Yuan-Ye Ma and the anonymous reviewers and editors of Zoological Research for their valuable opinions and suggestions.

REFERENCES

- Adams T, Rosenkranz JA. 2016. Social isolation during postweaning development causes hypoactivity of neurons in the medial nucleus of the male rat amygdala. *Neuropsychopharmacology*, **41**(7): 1929-1940.
- Ahern M, Goodell DJ, Adams J, Bland ST. 2016. Brain regional differences in social encounter-induced Fos expression in male and female rats after post-weaning social isolation. *Brain Research*, **1630**: 120-133.
- Andersen SL, Teicher MH. 2004. Delayed effects of early stress on hippocampal development. *Neuropsychopharmacology*, **29**(11): 1988-1993.
- Andersen SL, Tomada A, Vincow ES, Valente E, Polcari A, Teicher MH. 2008. Preliminary evidence for sensitive periods in the effect of childhood sexual abuse on regional brain development. *The Journal of Neuropsychiatry and Clinical Neurosciences*, **20**(3): 292-301.
- Andersen SL. 2015. Exposure to early adversity: points of cross-species translation that can lead to improved understanding of depression. *Development and Psychopathology*, **27**(2): 477-491.
- Andrews MW, Rosenblum LA. 1991. Attachment in monkey infants raised in variable- and low-demand environments. *Child Development*, **62**(4): 686-693.
- Anisman H, Zaharia MD, Meaney MJ, Merali Z. 1998. Do early-life events permanently alter behavioral and hormonal responses to stressors? *International Journal of Developmental Neuroscience*, **16**(3-4): 149-164.
- Avishai-Eliner S, Yi SJ, Baram TZ. 1996. Developmental profile of messenger RNA for the corticotropin-releasing hormone receptor in the rat limbic system. *Developmental Brain Research*, **91**(2): 159-163.
- Azmitia EC, Gannon PJ. 1986. The primate serotonergic system: a review of human and animal studies and a report on Macaca fascicularis. *Advances in Neurology*, **43**: 407-468.
- Babenko O, Kovalchuk I, Metz GAS. 2015. Stress-induced perinatal and transgenerational epigenetic programming of brain development and mental health. *Neuroscience & Biobehavioral Reviews*, **48**: 70-91.
- Bagot RC, Meaney MJ. 2010. Epigenetics and the biological basis of gene × environment interactions. *Journal of the American Academy of Child & Adolescent Psychiatry*, **49**(8): 752-771.

- Bailey RC, Aunger R. 1990. Humans as primates: the social relationships of Efe pygmy men in comparative perspective. *International Journal of Primatology*, **11**(2): 127-146.
- Bale TL. 2014. Lifetime stress experience: transgenerational epigenetics and germ cell programming. *Dialogues in Clinical Neuroscience*, **16**(3): 297-305.
- Baram TZ, Hatalski CG. 1998. Neuropeptide-mediated excitability: a key triggering mechanism for seizure generation in the developing brain. *Trends in Neurosciences*, **21**(11): 471-476.
- Barch D, Pagliaccio D, Belden A, Harms MP, Gaffrey M, Sylvester CM, Tillman R, Luby J. 2016. Effect of hippocampal and amygdala connectivity on the relationship between preschool poverty and school-age depression. *American Journal of Psychiatry*, **173**(6): 625-634.
- Barr CS, Newman TK, Becker ML, Parker CC, Champoux M, Lesch KP, Goldman D, Suomi SJ, Higley JD. 2003. The utility of the non-human primate model for studying gene by environment interactions in behavioral research. *Genes, Brain and Behavior*, **2**(6): 336-340.
- Barr CS, Newman TK, Shannon C, Parker C, Dvoskin RL, Becker ML, Schwandt M, Champoux M, Lesch KP, Goldman D, Suomi SJ, Higley JD. 2004. Rearing condition and rh5-HTTLPR interact to influence limbic-hypothalamic-pituitary-adrenal axis response to stress in infant macaques. *Biological Psychiatry*, **55**(7): 733-738.
- Barrett CE, Noble P, Hanson E, Pine DS, Winslow JT, Nelson EE. 2009. Early adverse rearing experiences alter sleep-wake patterns and plasma cortisol levels in juvenile rhesus monkeys. *Psychoneuroendocrinology*, **34**(7): 1029-1040.
- Bartesaghi R, Serrai A. 2001. Effects of early environment on granule cell morphology in the dentate gyrus of the guinea-pig. *Neuroscience*, **102**(1): 87-100.
- Bartesaghi R, Raffi M, Severi S. 2003a. Effects of early isolation on layer ii neurons in the entorhinal cortex of the guinea pig. *Neuroscience*, **120**(3): 721-732.
- Bartesaghi R, Severi S, Guidi S. 2003b. > effects of early environment on pyramidal neuron morphology in field CA1 of the guinea-pig. *Neuroscience*, **116**(3): 715-732.
- Bartesaghi R, Severi S. 2004. effects of early environment on field CA2 pyramidal neurons in the guinea-pig. *Neuroscience*, **123**(3): 703-714.
- Bastian ML, Sponberg AC, Sponberg AC, Suomi SJ, Higley JD. 2003. Long-term effects of infant rearing condition on the acquisition of dominance rank in juvenile and adult rhesus macaques (*Macaca mulatta*). *Developmental Psychobiology*, **42**(1): 44-51.
- Bauer PM, Hanson JL, Pierson RK, Davidson RJ, Pollak SD. 2009. Cerebellar volume and cognitive functioning in children who experienced early deprivation. *Biological Psychiatry*, **66**(12): 1100-1106.
- Bauer SA, Baker KC. 2016. Persistent effects of peer rearing on abnormal and species-appropriate activities but not social behavior in group-housed rhesus macaques (*Macaca mulatta*). *Comparative Medicine*, **66**(2): 129-136.
- Baysinger CM, Brandt EM, Mitchell G. 1972. Development of infant social isolate monkeys (*Macaca mulatta*) in their isolation environments. *Primates*, **13**(3): 257-270.
- Beauchamp AJ, Gluck JP. 1988. Associative processes in differentially reared monkeys (*Macaca mulatta*): sensory preconditioning. *Developmental Psychobiology*, **21**(4): 355-364.
- Beauchamp AJ, Gluck JP, Fouty HE, Lewis MH. 1991. Associative processes in differentially reared rhesus monkeys (*Macaca mulatta*): blocking. *Developmental Psychobiology*, **24**(3): 175-189.
- Becker K, Abraham A, Kindler J, Helmeke C, Braun K. 2007. Exposure to neonatal separation stress alters exploratory behavior and corticotropin releasing factor expression in neurons in the amygdala and hippocampus. *Developmental Neurobiology*, **67**(5): 617-629.
- Benjet C. 2010. Childhood adversities of populations living in low-income countries: prevalence, characteristics, and mental health consequences. *Current Opinion in Psychiatry*, **23**(4): 356-362.
- Bennett AJ, Lesch KP, Heils A, Long JC, Lorenz JG, Shoaf SE, Champoux M, Suomi SJ, Linnoila MV, Higley JD. 2002. Early experience and serotonin transporter gene variation interact to influence primate CNS function. *Molecular Psychiatry*, **7**(1): 118-122.
- Bennett AJ, Suomi SJ, Hopkins WD. 2008. Effects of early adverse experiences on behavioural lateralisation in rhesus monkeys (*Macaca mulatta*). *Laterality: Asymmetries of Body, Brain and Cognition*, **13**(3): 282-292.
- Bernstein IS, Mason WA. 1963. Group formation by rhesus monkeys. *Animal Behaviour*, **11**(1): 28-31.
- Bernstein IS, Williams LE. 1983. Ontogenetic changes and the stability of rhesus monkey dominance relationships. *Behavioural Processes*, **8**(4): 379-392.
- Bernstein IS, Cooper MA. 1999. Dominance in assamese macaques (*Macaca assamensis*). *American Journal of Primatology*, **48**(4): 283-289.
- Bick J, Nelson CA. 2016. Early adverse experiences and the developing brain. *Neuropsychopharmacology*, **41**(1): 177-196.
- Blaze J, Roth TL. 2015. Evidence from clinical and animal model studies of the long-term and transgenerational impact of stress on DNA methylation. *Seminars in Cell & Developmental Biology*, **43**: 76-84.
- Blaze J, Asok A, Roth TL. 2015a. Long-term effects of early-life caregiving experiences on brain-derived neurotrophic factor histone acetylation in the adult rat mPFC. *Stress*, **18**(6): 607-615.
- Blaze J, Asok A, Roth TL. 2015b. The long-term impact of adverse caregiving environments on epigenetic modifications and telomeres. *Frontiers in Behavioral Neuroscience*, **9**: 79.
- Boccia ML, Reite M, Kaemingk K, Held P, Laudenslager M. 1989. Behavioral and autonomic responses to peer separation in pigtail macaque monkey infants. *Developmental Psychobiology*, **22**(5): 447-461.
- Bock J, Gruss M, Becker S, Braun K. 2005. Experience-induced changes of dendritic spine densities in the prefrontal and sensory cortex: correlation with developmental time windows. *Cerebral Cortex*, **15**(6): 802-808.
- Bogart SL, Bennett AJ, Schapiro SJ, Reamer LA, Hopkins WD. 2014. Different early rearing experiences have long-term effects on cortical organization in captive chimpanzees (*Pan troglodytes*). *Developmental Science*, **17**(2): 161-174.
- Bohn MC, Howard E, Vielkind U, Krozowski Z. 1991. Glial cells express both mineralocorticoid and glucocorticoid receptors. *The Journal of Steroid Biochemistry and Molecular Biology*, **40**(1-3): 105-111.
- Bohn MC, O'banion MK, Young DA, Giuliano R, Hussain S, Dean DO, Cunningham LA. 1994. *In vitro* studies of glucocorticoid effects on neurons and astrocytes. *Annals of the New York Academy of Sciences*, **746**: 243-258.
- Botteron KN, Raichle ME, Drevets WC, Heath AC, Todd RD. 2002.

- Volumetric reduction in left subgenual prefrontal cortex in early onset depression. *Biological Psychiatry*, **51**(4): 342-344.
- Bowlby J. 1969. Attachment and Loss: Volume I: Attachment. London: The Hogarth Press and the Institute of Psycho-Analysis.(请核对出版信息)
- Bowley MP, Drevets WC, Öngür D, Price JL. 2002. Low glial numbers in the amygdala in major depressive disorder. *Biological Psychiatry*, **52**(5): 404-412.
- Brand SR, Brennan PA, Newport DJ, Smith AK, Weiss T, Stowe ZN. 2010. The impact of maternal childhood abuse on maternal and infant HPA axis function in the postpartum period. *Psychoneuroendocrinology*, **35**(5): 686-693.
- Braun K, Champagne FA. 2014. Paternal influences on offspring development: behavioural and epigenetic pathways. *Journal of Neuroendocrinology*, **26**(10): 697-706.
- Bremner JD, Randall P, Vermetten E, Staib L, Bronen RA, Mazure C, Capelli S, McCarthy G, Innis RB, Charney DS. 1997. Magnetic resonance imaging-based measurement of hippocampal volume in posttraumatic stress disorder related to childhood physical and sexual abuse—a preliminary report. *Biological Psychiatry*, **41**(1): 23-32.
- Bremner JD, Elzinga B, Schmahl C, Vermetten E. 2007. Structural and functional plasticity of the human brain in posttraumatic stress disorder. *Progress in Brain Research*, **167**: 171-186.
- Bridges N, Slais K, Syková E. 2008. The effects of chronic corticosterone on hippocampal astrocyte numbers: a comparison of male and female Wistar rats. *Acta Neurobiologiae Experimentalis*, **68**(2): 131-138.
- Bruce J, Fisher PA, Pears KC, Levine S. 2009. Morning cortisol levels in preschool-aged foster children: differential effects of maltreatment type. *Developmental Psychobiology*, **51**(1): 14-23.
- Brunelli RL, Blake J, Willits N, Rommeck I, Mccowan B. 2014. Effects of a mechanical response-contingent surrogate on the development of behaviors in nursery-reared rhesus macaques (*Macaca mulatta*). *Journal of the American Association for Laboratory Animal Science*, **53**(5): 464-471.
- Brunson KL, Kramár E, Lin B, Chen YC, Colgin LL, Yanagihara TK, Lynch G, Baram TZ. 2005. Mechanisms of late-onset cognitive decline after early-life stress. *Journal of Neuroscience*, **25**(41): 9328-9338.
- Bryan GK, Riesen AH. 1989. Deprived somatosensory-motor experience in stump-tailed monkey neocortex: dendritic spine density and dendritic branching of layer IIIB pyramidal cells. *The Journal of Comparative Neurology*, **286**(2): 208-217.
- Buckley MJ. 2005. The role of the perirhinal cortex and hippocampus in learning, memory, and perception. *The Quarterly Journal of Experimental Psychology Section B: Comparative and Physiological Psychology*, **58**(3-4): 246-268.
- Bush G, Luu P, Posner MI. 2000. Cognitive and emotional influences in anterior cingulate cortex. *Trends in Cognitive Sciences*, **4**(6): 215-222.
- Caldji C, Tannenbaum B, Sharma S, Francis D, Plotsky PM, Meaney MJ. 1998. Maternal care during infancy regulates the development of neural systems mediating the expression of fearfulness in the rat. *Proceedings of the National Academy of Sciences of the United States of America*, **95**(9): 5335-5340.
- Capitano JP, Mason WA. 2000. Cognitive style: problem solving by rhesus macaques (*Macaca mulatta*) reared with living or inanimate substitute mothers. *Journal of Comparative Psychology*, **114**(2): 115-125.
- Capitano JP, Mendoza SP, Mason WA, Maninger N. 2005. Rearing environment and hypothalamic-pituitary-adrenal regulation in young rhesus monkeys (*Macaca mulatta*). *Developmental Psychobiology*, **46**(4): 318-330.
- Carlson M, Earls F. 1997. Psychological and neuroendocrinological sequelae of early social deprivation in institutionalized children in Romania. *Annals of the New York Academy of Sciences*, **807**: 419-428.
- Carpenter LL, Tyrka AR, Ross NS, Khoury L, Anderson GM, Price LH. 2009. Effect of childhood emotional abuse and age on cortisol responsivity in adulthood. *Biological Psychiatry*, **66**(1): 69-75.
- Carrion VG, Weems CF, Eliez S, Patwardhan A, Brown W, Ray RD, Reiss AL. 2001. Attenuation of frontal asymmetry in pediatric posttraumatic stress disorder. *Biological Psychiatry*, **50**(12): 943-951.
- Carrion VG, Weems CF, Watson C, Eliez S, Menon V, Reiss AL. 2009. Converging evidence for abnormalities of the prefrontal cortex and evaluation of midsagittal structures in pediatric posttraumatic stress disorder: an MRI study. *Psychiatry Research: Neuroimaging*, **172**(3): 226-234.
- Casement MD, Guyer AE, Hipwell AE, Mcaloon RL, Hoffmann AM, Keenan KE, Forbes EE. 2014. Girls' challenging social experiences in early adolescence predict neural response to rewards and depressive symptoms. *Developmental Cognitive Neuroscience*, **8**: 18-27.
- Casement MD, Shaw DS, Sitnick SL, Musselman SC, Forbes EE. 2015. Life stress in adolescence predicts early adult reward-related brain function and alcohol dependence. *Social Cognitive and Affective Neuroscience*, **10**(3): 416-423.
- Caspi A, McClay J, Moffitt TE, Mill J, Martin J, Craig IW, Taylor A, Poulton R. 2002. Role of genotype in the cycle of violence in maltreated children. *Science*, **297**(5582): 851-854.
- Chamove AS, Rosenblum LA, Harlow HF. 1973. Monkeys (*Macaca mulatta*) raised only with peers. A pilot study. *Animal Behaviour*, **21**(2): 316-325.
- Champagne F, Diorio J, Sharma S, Meaney MJ. 2001. Naturally occurring variations in maternal behavior in the rat are associated with differences in estrogen-inducible central oxytocin receptors. *Proceedings of the National Academy of Sciences of the United States of America*, **98**(22): 12736-12741.
- Champagne F, Meaney MJ. 2001. Like mother, like daughter: evidence for non-genomic transmission of parental behavior and stress responsivity. *Progress in Brain Research*, **133**: 287-302.
- Champagne FA, Weaver ICG, Diorio J, Sharma S, Meaney MJ. 2003. Natural variations in maternal care are associated with estrogen receptor α expression and estrogen sensitivity in the medial preoptic area. *Endocrinology*, **144**(11): 4720-4724.
- Champagne FA, Weaver ICG, Diorio J, Dymov S, Szyf M, Meaney MJ. 2006. Maternal care associated with methylation of the estrogen receptor- α 1b promoter and estrogen receptor- α expression in the medial preoptic area of female offspring. *Endocrinology*, **147**(6): 2909-2915.
- Champagne FA. 2008. Epigenetic mechanisms and the transgenerational effects of maternal care. *Frontiers in Neuroendocrinology*, **29**(3): 386-397.
- Champagne FA, Curley JP. 2009. Epigenetic mechanisms mediating the long-term effects of maternal care on development. *Neuroscience & Biobehavioral Reviews*, **33**(4): 593-600.
- Champoux M, Coe CL, Schanberg SM, Kuhn CM, Suomi SJ. 1989a. Hormonal effects of early rearing conditions in the infant rhesus monkey. *American Journal of Primatology*, **19**(2): 111-117.
- Champoux M, Metz B, Suomi SJ. 1989b. Rehousing Nonreproductive

- Rhesus Macaques with Weanlings: I. Behavior of Adults toward Weanlings. *Laboratory Primate Newsletter*, **28**(4): 1-4.
- Champoux M, Metz B, Suomi SJ. 1991. Behavior of nursery/peer-reared and mother-reared rhesus monkeys from birth through 2 years of age. *Primates*, **32**(4): 509-514.
- Champoux M, Byrne E, Delizio R, Suomi SJ. 1992. Motherless mothers revisited: rhesus maternal behavior and rearing history. *Primates*, **33**(2): 251-255.
- Cicchetti D, Rogosch FA. 2001. Diverse patterns of neuroendocrine activity in maltreated children. *Development and Psychopathology*, **13**(3): 677-693.
- Cicchetti D, Rogosch FA, Gunnar MR, Toth SL. 2010. The differential impacts of early physical and sexual abuse and internalizing problems on daytime cortisol rhythm in school-aged children. *Child Development*, **81**(1): 252-269.
- Clarke AS. 1993. Social rearing effects on HPA axis activity over early development and in response to stress in rhesus monkeys. *Developmental Psychobiology*, **26**(8): 433-446.
- Clarke AS, Hedeker DR, Ebert MH, Schmidt DE, McKinney WT, Kraemer GW. 1996. Rearing experience and biogenic amine activity in infant rhesus monkeys. *Biological Psychiatry*, **40**(5): 338-352.
- Clarke AS, Kraemer GW, Kupfer DJ. 1998. Effects of rearing condition on HPA axis response to fluoxetine and desipramine treatment over repeated social separations in young rhesus monkeys. *Psychiatry Research*, **79**(2): 91-104.
- Clarke AS, Ebert MH, Schmidt DE, McKinney WT, Kraemer GW. 1999. Biogenic amine activity in response to fluoxetine and desipramine in differentially reared rhesus monkeys. *Biological Psychiatry*, **46**(2): 221-228.
- Cohen RA, Grieve S, Hoth KF, Paul RH, Sweet L, Tate D, Gunstad J, Stroud L, McCaffery J, Hitsman B, Niaura R, Clark CR, MacFarlane A, Bryant R, Gordon E, Williams LM. 2006. Early life stress and morphometry of the adult anterior cingulate cortex and caudate nuclei. *Biological Psychiatry*, **59**(10): 975-982.
- Collishaw S, Pickles A, Messer J, Rutter M, Shearer C, Maughan B. 2007. Resilience to adult psychopathology following childhood maltreatment: evidence from a community sample. *Child Abuse & Neglect*, **31**(3): 211-229.
- Coplan JD, Andrews MW, Rosenblum LA, Owens MJ, Friedman S, Gorman JM, Nemeroff CB. 1996. Persistent elevations of cerebrospinal fluid concentrations of corticotropin-releasing factor in adult nonhuman primates exposed to early-life stressors: implications for the pathophysiology of mood and anxiety disorders. *Proceedings of the National Academy of Sciences of the United States of America*, **93**(4): 1619-1623.
- Coplan JD, Smith ELP, Altemus M, Scharf BA, Owens MJ, Nemeroff CB, Gorman JM, Rosenblum LA. 2001. Variable foraging demand rearing: sustained elevations in cisternal cerebrospinal fluid corticotropin-releasing factor concentrations in adult primates. *Biological Psychiatry*, **50**(3): 200-204.
- Coplan JD, Altemus M, Mathew SJ, Smith ELP, Sharf B, Coplan PM, Kral JG, Gorman JM, Owens MJ, Nemeroff CB, Rosenblum LA. 2005. Synchronized maternal-infant elevations of primate CSF CRF concentrations in response to variable foraging demand. *CNS Spectrums*, **10**(7): 530-536.
- Cotter D, Mackay D, Landau S, Kerwin R, Everall I. 2001. Reduced glial cell density and neuronal size in the anterior cingulate cortex in major depressive disorder. *Archives of General Psychiatry*, **58**(6): 545-553.
- Craig IW. 2005. The role of monoamine oxidase A, MAOA, in the aetiology of antisocial behaviour: the importance of gene-environment interactions. *Novartis Foundation Symposium*, **268**: 227-237; discussion 237-241, 242-253. (请核对页码)
- Cross HA, Harlow HF. 1965. Prolonged and progressive effects of partial isolation on the behavior of macaque monkeys. *Journal of Experimental Research in Personality*, **1**(1): 39-49.
- Curley JP, Mashoodh R, Champagne FA. 2011. Epigenetics and the origins of paternal effects. *Hormones and Behavior*, **59**(3): 306-314.
- Cushing BS, Kramer KM. 2005. Mechanisms underlying epigenetic effects of early social experience: the role of neuropeptides and steroids. *Neuroscience & Biobehavioral Reviews*, **29**(7): 1089-1105.
- Cutuli JJ, Wiik KL, Herbers JE, Gunnar MR, Masten AS. 2010. Cortisol function among early school-aged homeless children. *Psychoneuroendocrinology*, **35**(6): 833-845.
- Czoty PW, Gould RW, Nader MA. 2009. Relationship between social rank and cortisol and testosterone concentrations in male cynomolgus monkeys (*Macaca fascicularis*). *Journal of Neuroendocrinology*, **21**(1): 68-76.
- Daskalakis NP, Bagot RC, Parker KJ, Vinkers CH, de Kloet ER. 2013. The three-hit concept of vulnerability and resilience: toward understanding adaptation to early-life adversity outcome. *Psychoneuroendocrinology*, **38**(9): 1858-1873.
- Davis M. 1992. The role of the amygdala in fear and anxiety. *Annual Review of Neuroscience*, **15**: 353-375.
- Day J, Savani S, Krempel BD, Nguyen M, Kitlinska JB. 2016. Influence of paternal preconception exposures on their offspring: through epigenetics to phenotype. *American Journal of Stem Cells*, **5**(1): 11-18.
- De Bellis MD, Chrousos GP, Dorn LD, Burke L, Halmers K, Kling MA, Trickett PK, Putnam FW. 1994. Hypothalamic-pituitary-adrenal axis dysregulation in sexually abused girls. *The Journal of Clinical Endocrinology and Metabolism*, **78**(2): 249-255.
- De Bellis MD, Keshavan MS, Clark DB, Casey BJ, Giedd JN, Boring AM, Frustaci K, Ryan ND. 1999. Developmental traumatology part II: brain development. *Biological Psychiatry*, **45**(10): 1271-1284.
- De Bellis MD, Hall J, Boring AM, Frustaci K, Moritz G. 2001. A pilot longitudinal study of hippocampal volumes in pediatric maltreatment-related posttraumatic stress disorder. *Biological Psychiatry*, **50**(4): 305-309.
- De Bellis MD, Keshavan MS, Shifflett H, Iyengar S, Beers SR, Hall J, Moritz G. 2002. Brain structures in pediatric maltreatment-related posttraumatic stress disorder: a sociodemographically matched study. *Biological Psychiatry*, **52**(11): 1066-1078.
- De Brito SA, Viding E, Sebastian CL, Kelly PA, Mechelli A, Maris H, McCrory EJ. 2013. Reduced orbitofrontal and temporal grey matter in a community sample of maltreated children. *Journal of Child Psychology and Psychiatry*, **54**(1): 105-112.
- De Weerth C, Ziji RH, Buitelaar JK. 2003. Development of cortisol circadian rhythm in infancy. *Early Human Development*, **73**(1-2): 39-52.
- Detting A, Pryce CR, Martin RD, Döbeli M. 1998. Physiological Responses to parental separation and a strange situation are related to parental care received in juvenile Goeldi's monkeys (*Callimico goeldii*). *Developmental Psychobiology*, **33**(1): 21-31.
- Detting AC, Feldon J, Pryce CR. 2002a. Early deprivation and behavioral and physiological responses to social separation/novelty in the marmoset. *Pharmacology Biochemistry and Behavior*, **73**(1): 259-269.
- Detting AC, Feldon J, Pryce CR. 2002b. Repeated parental deprivation in

- the infant common marmoset (*Callithrix jacchus*, primates) and analysis of its effects on early development. *Biological Psychiatry*, **52**(11): 1037-1046.
- Dettmer AM, Ruggiero AM, Novak MA, Meyer JS, Suomi SJ. 2008. Surrogate mobility and orientation affect the early neurobehavioral development of infant rhesus macaques (*Macaca mulatta*). *Developmental Psychobiology*, **50**(4): 418-422.
- Dettmer AM, Novak MA, Suomi SJ, Meyer JS. 2012. Physiological and behavioral adaptation to relocation stress in differentially reared rhesus monkeys: hair cortisol as a biomarker for anxiety-related responses. *Psychoneuroendocrinology*, **37**(2): 191-199.
- Devinsky O, Morrell MJ, Vogt BA. 1995. Contributions of anterior cingulate cortex to behaviour. *Brain*, **118**(1): 279-306.
- DeVore I. 1990. Introduction: current studies on primate socioecology and evolution. *International Journal of Primatology*, **11**(1): 1-5.
- Dickmeis T, Weger BD, Weger M. 2013. The circadian clock and glucocorticoids – interactions across many time scales. *Molecular and Cellular Endocrinology*, **380**(1-2): 2-15.
- Diorio J, Meaney MJ. 2007. Maternal programming of defensive responses through sustained effects on gene expression. *Journal of Psychiatry and Neuroscience*, **32**(4): 275-284.
- Dozier M, Manni M, Gordon MK, Peloso E, Gunnar MR, Stovall-McClough KC, Eldreth D, Levine S. 2006. Foster children's diurnal production of cortisol: an exploratory study. *Child Maltreatment*, **11**(2): 189-197.
- Drevets WC, Price JL, Simpson JR Jr, Todd RD, Reich T, Vannier M, Raichle ME. 1997. Subgenual prefrontal cortex abnormalities in mood disorders. *Nature*, **386**(6627): 824-827.
- Drevets WC, Savitz J, Trimble M. 2008. The subgenual anterior cingulate cortex in mood disorders. *CNS Spectrums*, **13**(8): 663-681.
- Eastman RF, Mason WA. 1975. Looking behavior in monkeys raised with mobile and stationary artificial mothers. *Developmental Psychobiology*, **8**(3): 213-221.
- Edmiston EE, Wang F, Mazure CM, Guiney J, Sinha R, Mayes LC, Blumberg HP. 2011. Corticostriatal-limbic gray matter morphology in adolescents with self-reported exposure to childhood maltreatment. *Archives of Pediatrics & Adolescent Medicine*, **165**(12): 1069-1077.
- Eichenbaum H, Otto T, Cohen NJ. 1992. The hippocampus—what does it do? *Behavioral and Neural Biology*, **57**(1): 2-36.
- Eichenbaum H, Schoenbaum G, Young B, Bunsey M. 1996. Functional organization of the hippocampal memory system. *Proceedings of the National Academy of Sciences of the United States of America*, **93**(24): 13500-13507.
- Eichenbaum H. 2001. The hippocampus and declarative memory: cognitive mechanisms and neural codes. *Behavioural Brain Research*, **127**(1-2): 199-207.
- Eley TC, Sugden K, Corsico A, Gregory AM, Sham P, McGuffin P, Plomin R, Craig IW. 2004. Gene-environment interaction analysis of serotonin system markers with adolescent depression. *Molecular Psychiatry*, **9**(10): 908-915.
- Elzinga BM, Roelofs K, Tollenaar MS, Bakvis P, van Pelt J, Spinhoven P. 2008. Diminished cortisol responses to psychosocial stress associated with lifetime adverse events: a study among healthy young subjects. *Psychoneuroendocrinology*, **33**(2): 227-237.
- Erwin J, Mitchell G, Maple T. 1973. Abnormal behavior in non-isolate-reared rhesus monkeys. *Psychological Reports*, **33**(2): 515-523.
- Essex MJ, Klein MH, Cho E, Kalin NH. 2002. Maternal stress beginning in infancy may sensitize children to later stress exposure: effects on cortisol and behavior. *Biological Psychiatry*, **52**(8): 776-784.
- Evans GW, Swain JE, King AP, Wang X, Javanbakht A, Ho SS, Angstadt M, Phan KL, Xie H, Liberzon I. 2016. Childhood cumulative risk exposure and adult amygdala volume and function. *Journal of Neuroscience Research*, **94**(6): 535-543.
- Fahlke C, Lorenz JG, Long J, Champoux M, Suomi SJ, Higley JD. 2000. Rearing experiences and stress-induced plasma cortisol as early risk factors for excessive alcohol consumption in nonhuman primates. *Alcoholism: Clinical and Experimental Research*, **24**(5): 644-650.
- Farrington DP, Loeber R. 2000. Epidemiology of juvenile violence. *Child and Adolescent Psychiatric Clinics of North America*, **9**(4): 733-748.
- Fekete JM, Norcross JL, Newman JD. 2000. Artificial turf foraging boards as environmental enrichment for pair-housed female squirrel monkeys. *Journal of the American Association for Laboratory Animal Science*, **39**(2): 22-26.
- Feng XL, Wang LN, Yang SC, Qin DD, Wang JH, Li CL, Lv LB, Ma YY, Hu XT. 2011. Maternal separation produces lasting changes in cortisol and behavior in rhesus monkeys. *Proceedings of the National Academy of Sciences of the United States of America*, **108**(34): 14312-14317.
- Fenoglio KA, Brunson KL, Baram TZ. 2006. Hippocampal neuroplasticity induced by early-life stress: functional and molecular aspects. *Frontiers in Neuroendocrinology*, **27**(2): 180-192.
- Fernald LCH, Gunnar MR. 2009. Poverty-alleviation program participation and salivary cortisol in very low-income children. *Social Science & Medicine*, **68**(12): 2180-2189.
- Fisher PA, Gunnar MR, Chamberlain P, Reid JB. 2000. Preventive intervention for maltreated preschool children: impact on children's behavior, neuroendocrine activity, and foster parent functioning. *Journal of the American Academy of Child & Adolescent Psychiatry*, **39**(11): 1356-1364.
- Fisher PA, Gunnar MR, Dozier M, Bruce J, Pears KC. 2006. Effects of therapeutic interventions for foster children on behavioral problems, caregiver attachment, and stress regulatory neural systems. *Annals of the New York Academy of Sciences*, **1094**: 215-225.
- Fisher PA, Stoolmiller M, Gunnar MR, Burraston BO. 2007. Effects of a therapeutic intervention for foster preschoolers on diurnal cortisol activity. *Psychoneuroendocrinology*, **32**(8-10): 892-905.
- Fleming AS, Kraemer GW, Gonzalez A, Lovic V, Rees S, Melo A. 2002. Mothering begets mothering: the transmission of behavior and its neurobiology across generations. *Pharmacology Biochemistry and Behavior*, **73**(1): 61-75.
- Flinn MV, England BG. 1997. Social economics of childhood glucocorticoid stress response and health. *American Journal of Physical Anthropology*, **102**(1): 33-53.
- Floeter MK, Greenough WT. 1979. Cerebellar plasticity: modification of Purkinje cell structure by differential rearing in monkeys. *Science*, **206**(4415): 227-229.
- Fonberg E. 1988. Dominance and aggression. *International Journal of Neuroscience*, **41**(3-4): 201-213.
- Francis D, Diorio J, Liu D, Meaney MJ. 1999. Nongenomic transmission across generations of maternal behavior and stress responses in the rat. *Science*, **286**(5442): 1155-1158.
- Galea LAM, McEwen BS, Tanapat P, Deak T, Spencer RL, Dhabhar FS. 1997. Sex differences in dendritic atrophy of CA3 pyramidal neurons in

- response to chronic restraint stress. *Neuroscience*, **81**(3): 689-697.
- Gallagher M, Chiba AA. 1996. The amygdala and emotion. *Current Opinion in Neurobiology*, **6**(2): 221-227.
- Garrett JE, Wellman CL. 2009. Chronic stress effects on dendritic morphology in medial prefrontal cortex: sex differences and estrogen dependence. *Neuroscience*, **162**(1): 195-207.
- Gershon A, Sudheimer K, Tirouvanziam R, Williams LM, O'hara R. 2013. The long-term impact of early adversity on late-life psychiatric disorders. *Current Psychiatry Reports*, **15**: 352.
- Gilmer WS, McKinney WT. 2003. Early experience and depressive disorders: human and non-human primate studies. *Journal of Affective Disorders*, **75**(2): 97-113.
- Gluck JP, Harlow HF, Schiltz KA. 1973. Differential effect of early enrichment and deprivation on learning in the rhesus monkey (*Macaca mulatta*). *Journal of Comparative and Physiological Psychology*, **84**(3): 598-604.
- Gluck JP, Sackett GP. 1974. Frustration and self-aggression in social isolate rhesus monkeys. *Journal of Abnormal Psychology*, **83**(3): 331-334.
- Gluck JP, Sackett GP. 1976. Extinction deficits in socially isolated rhesus monkeys (*Macaca mulatta*). *Developmental Psychology*, **12**(2): 173-174.
- Gluckman PD, Hanson MA. 2004. Living with the past: evolution, development, and patterns of disease. *Science*, **305**(5691): 1733-1736.
- Gluckman PD, Hanson MA, Beedle AS. 2007. Early life events and their consequences for later disease: a life history and evolutionary perspective. *American Journal of Human Biology*, **19**(1): 1-19.
- Goff B, Tottenham N. 2015. Early-life adversity and adolescent depression: mechanisms involving the ventral striatum. *CNS Spectrums*, **20**(4): 337-345.
- Goldfoot DA, Wallen K, Neff DA, McBair MC, Goy RW. 1984. Social influences on the display of sexually dimorphic behavior in rhesus monkeys: isosexual rearing. *Archives of Sexual Behavior*, **13**(5): 395-412.
- Gonzalez A, Jenkins JM, Steiner M, Fleming AS. 2009. The relation between early life adversity, cortisol awakening response and diurnal salivary cortisol levels in postpartum women. *Psychoneuroendocrinology*, **34**(1): 76-86.
- Goo GP, Sassenrath EN. 1980. Persistent adrenocortical activation in female rhesus monkeys after new breeding groups formation. *Journal of Medical Primatology*, **9**(6): 325-334.
- Goosen C. 1981. Abnormal behavior patterns in rhesus monkeys: symptoms of mental disease. *Biological Psychiatry*, **16**(8): 697-716.
- Gorman JM, Mathew S, Coplan J. 2002. Neurobiology of early life stress: nonhuman primate models. *Seminars in Clinical Neuropsychiatry*, **7**(2): 96-103.
- Gos T, Bock J, Poeggel G, Braun K. 2008. Stress-induced synaptic changes in the rat anterior cingulate cortex are dependent on endocrine developmental time windows. *Synapse*, **62**(3): 229-232.
- Griffin GA, Harlow HF. 1966. Effects of three months of total social deprivation on social adjustment and learning in the rhesus monkey. *Child Development*, **37**(3): 533-547. (与下条参考文献重复, 请核对)
- Griffin GA, Harlow HF. 1966. Effects of three months of total social deprivation on social adjustment and learning in the rhesus monkey. *Child Development*, **37**(3): 533-547.
- Gröger N, Matas E, Gos T, Lesse A, Poeggel G, Braun K, Bock J. 2016. The transgenerational transmission of childhood adversity: behavioral, cellular, and epigenetic correlates. *Journal of Neural Transmission*, **123**(9): 1037-1052.
- Gunnar MR, Porter FL, Wolf CM, Rigatuso J, Larson MC. 1995. Neonatal stress reactivity: predictions to later emotional temperament. *Child Development*, **66**(1): 1-13.
- Gunnar MR, Morison SJ, Chisholm K, Schuder M. 2001. Salivary cortisol levels in children adopted from romanian orphanages. *Development and Psychopathology*, **13**(3): 611-628.
- Gunnar MR, Vazquez DM. 2001. Low cortisol and a flattening of expected daytime rhythm: potential indices of risk in human development. *Development and Psychopathology*, **13**(3): 515-538.
- Gustafsson PE, Janlert U, Theorell T, Hammarström A. 2010. Life-course socioeconomic trajectories and diurnal cortisol regulation in adulthood. *Psychoneuroendocrinology*, **35**(4): 613-623.
- Hanson JL, Chung MK, Avants BB, Shirtcliff EA, Gee JC, Davidson RJ, Pollak SD. 2010. Early stress is associated with alterations in the orbitofrontal cortex: a tensor-based morphometry investigation of brain structure and behavioral risk. *Journal of Neuroscience*, **30**(22): 7466-7472.
- Hanson JL, Nacewicz BM, Sutterer MJ, Cayo AA, Schaefer SM, Rudolph KD, Shirtcliff EA, Pollak SD, Davidson RJ. 2015. Behavioral problems after early life stress: contributions of the hippocampus and amygdala. *Biological Psychiatry*, **77**(4): 314-323.
- Harlow HF. 1958. The nature of love. *American Psychologist*, **13**(12): 673-685.
- Harlow HF, Zimmermann RR. 1959. Affectional response in the infant monkey: orphaned baby monkeys develop a strong and persistent attachment to inanimate surrogate mothers. *Science*, **130**(3373): 421-432.
- Harlow HF. 1962. The heterosexual affectional system in monkeys. *American Psychologist*, **17**(1): 1-9.
- Harlow HF, Harlow M. 1962. Social deprivation in monkeys. *Scientific American*, **207**(5): 136-147.
- Harlow HF, Rowland GL, Griffin GA. 1964. The effect of total social deprivation on the development of monkey behavior. *Psychiatric Research Reports*, **19**: 116-135.
- Harlow HF, Dodsworth RO, Harlow MK. 1965. Total social isolation in monkeys. *Proceedings of the National Academy of Sciences of the United States of America*, **54**(1): 90-97.
- Harlow HF, Harlow MK. 1965. The affectional systems. In: Schrier AJ, Harlow HF, Stollnitz F. Behavior of Nonhuman Primates. New York: Academic Press, 287-334.
- Harlow HF, Joslyn WD, Senko MG, Dopp A. 1966. Behavioral aspects of reproduction in primates. *Journal of Animal Science*, **25**(S): 49-65.
- Harlow HF, Suomi SJ. 1971a. Production of depressive behaviors in young monkeys. *Journal of Autism and Childhood Schizophrenia*, **1**(3): 246-255.
- Harlow HF, Suomi SJ. 1971b. Social recovery by isolation-reared monkeys. *Proceedings of the National Academy of Sciences of the United States of America*, **68**(7): 1534-1538.
- Hart H, Rubia K. 2012. Neuroimaging of child abuse: a critical review. *Frontiers in Human Neuroscience*, **6**: 52.
- Hart J G, Gunnar M, Cicchetti D. 1995. Salivary cortisol in maltreated children: evidence of relations between neuroendocrine activity and social competence. *Development and Psychopathology*, **7**(1): 11-26.
- Heim C, Ehler U, Hellhammer DH. 2000a. The potential role of hypocortisolism in the pathophysiology of stress-related bodily disorders. *Psychoneuroendocrinology*, **25**(1): 1-35.

- Heim C, Newport DJ, Heit S, Graham YP, Wilcox M, Bonsall R, Miller AH, Nemeroff CB. 2000b. Pituitary-adrenal and autonomic responses to stress in women after sexual and physical abuse in childhood. *JAMA*, **284**(5): 592-597.
- Heim C, Nemeroff CB. 2001. The role of childhood trauma in the neurobiology of mood and anxiety disorders: preclinical and clinical studies. *Biological Psychiatry*, **49**(12): 1023-1039.
- Heinrichs SC, Menzaghi FE, Merlo Pich E, Britton KT, Koob GF. 1995. The role of CRF in behavioral aspects of stress. *Annals of the New York Academy of Sciences*, **771**: 92-104.
- Hennessy MB, Kaplan JN. 1982. Influence of the maternal surrogate on pituitary-adrenal activity and behavior of infant squirrel monkeys. *Developmental Psychobiology*, **15**(5): 423-431.
- Herman JP, Prewitt CMF, Cullinan WE. 1996. Neuronal circuit regulation of the hypothalamo-pituitary-adrenocortical stress axis. *Critical Reviews™ in Neurobiology*, **10**(3-4): 371-394.
- Herman JP, Cullinan WE. 1997. Neurocircuitry of stress: central control of the hypothalamo-pituitary-adrenocortical axis. *Trends in Neurosciences*, **20**(2): 78-84.
- Herman JP, Figueiredo H, Mueller NK, Ulrich-Lai Y, Ostrander MM, Choi DC, Cullinan WE. 2003. Central mechanisms of stress integration: hierarchical circuitry controlling hypothalamo-pituitary-adrenocortical responsiveness. *Frontiers in Neuroendocrinology*, **24**(3): 151-180.
- Higley JD, Hasert MF, Suomi SJ, Linnoila M. 1991. Nonhuman primate model of alcohol abuse: effects of early experience, personality, and stress on alcohol consumption. *Proceedings of the National Academy of Sciences of the United States of America*, **88**(16): 7261-7265.
- Higley JD, Suomi SJ, Linnoila M. 1992. A longitudinal assessment of CSF monoamine metabolite and plasma cortisol concentrations in young rhesus monkeys. *Biological Psychiatry*, **32**(2): 127-145.
- Higley JD, Mehlman PT, Poland RE, Taub DM, Vickers J, Suomi SJ, Linnoila M. 1996a. CSF testosterone and 5-HIAA correlate with different types of aggressive behaviors. *Biological Psychiatry*, **40**(11): 1067-1082.
- Higley JD, Suomi SJ, Linnoila M. 1996b. A nonhuman primate model of type II alcoholism? Part 2. Diminished social competence and excessive aggression correlates with low cerebrospinal fluid 5-hydroxyindoleacetic acid concentrations. *Alcoholism: Clinical & Experimental Research*, **20**(4): 643-650.
- Hinde RA, Spencer-Booth Y, Bruce M. 1966. Effects of 6-day maternal deprivation on rhesus monkey infants. *Nature*, **210**(5040): 1021-1023.
- Hinde RA, Spencer-Booth Y. 1967. The behaviour of socially living rhesus monkeys in their first two and a half years. *Animal Behaviour*, **15**(1): 169-196.
- Hinde RA, McGinnis L. 1977. Some factors influencing the effects of temporary mother-infant separation: some experiments with rhesus monkeys. *Psychological Medicine*, **7**(2): 197-212.
- Hotchkiss CE, Paule MG. 2003. Effect of pair-housing on operant behavior task performance by rhesus monkeys. *Journal of the American Association for Laboratory Animal Science*, **42**(4): 38-41.
- Houston SM, Herting MM, Sowell ER. 2014. The neurobiology of childhood structural brain development: conception through adulthood. In: Andersen SL, Pine DS. *The Neurobiology of Childhood: Current Topics in Behavioral Neurosciences*. Berlin Heidelberg: Springer, **16**: 3-17.
- Howell BR, Grand AP, McCormack KM, Shi YD, Laprarie JL, Maestripieri D, Styner MA, Sanchez MM. 2014. Early adverse experience increases emotional reactivity in juvenile rhesus macaques: relation to amygdala volume. *Developmental Psychobiology*, **56**(8): 1735-1746.
- Hoyer C, Richter H, Brandwein C, Riva MA, Gass P. 2013. Preconceptional paternal exposure to a single traumatic event affects postnatal growth of female but not male offspring. *Neuroreport*, **24**(15): 856-860.
- Ichise M, Vines DC, Gura T, Anderson GM, Suomi SJ, Higley JD, Innis RB. 2006. Effects of early life stress on [¹¹C]DASB positron emission tomography imaging of serotonin transporters in adolescent peer- and mother-reared rhesus monkeys. *Journal of Neuroscience*, **26**(17): 4638-4643.
- Insel TR. 1992. Oxytocin—a neuropeptide for affiliation: evidence from behavioral, receptor autoradiographic, and comparative studies. *Psychoneuroendocrinology*, **17**(1): 3-35.
- Jablonka E, Lamb MJ. 2002. The changing concept of epigenetics. *Annals of the New York Academy of Sciences*, **981**: 82-96.
- Jauregui-Huerta F, Ruvalcaba-Delgadillo Y, Gonzalez-Perez O, Gonzalez-Castaneda R, Garcia-Estrada J, Luquin S. 2010. Responses of glial cells to stress and glucocorticoids. *Current Immunology Reviews*, **6**(3): 195-204.
- Javanbakht A, King AP, Evans GW, Swain JE, Angstadt M, Phan KL, Liberzon I. 2015. Childhood Poverty Predicts Adult Amygdala and Frontal Activity and Connectivity in Response to Emotional Faces. *Frontiers in Behavioral Neuroscience*, **9**: 154.
- Jawahar MC, Murgatroyd C, Harrison EL, Baune BT. 2015. Epigenetic alterations following early postnatal stress: a review on novel aetiological mechanisms of common psychiatric disorders. *Clinical Epigenetics*, **7**: 122.
- Jedd K, Hunt RH, Cicchetti D, Hunt E, Cowell RA, Rogosch FA, Toth SL, Thomas KM. 2015. Long-term consequences of childhood maltreatment: altered amygdala functional connectivity. *Development and Psychopathology*, **27**(4Pt2): 1577-1589.
- Johnson LR, Farb C, Morrison JH, McEwen BS, Ledoux JE. 2005. Localization of glucocorticoid receptors at postsynaptic membranes in the lateral amygdala. *Neuroscience*, **136**(1): 289-299.
- Jung-Testas I, Baulieu EE. 1998. Steroid hormone receptors and steroid action in rat glial cells of the central and peripheral nervous system. *The Journal of Steroid Biochemistry and Molecular Biology*, **65**(1-6): 243-251.
- Kaemingk K, Reite M. 1987. Social environment and nocturnal sleep: studies in peer-reared monkeys. *Sleep*, **10**(6): 542-550.
- Kagan J, Reznick JS, Snidman N. 1988. Biological bases of childhood shyness. *Science*, **240**(4849): 167-171.
- Kaplan J. 1974. Growth and behavior of surrogate-reared squirrel monkeys. *Developmental Psychobiology*, **7**(1): 7-13.
- Karere GM, Kinnally EL, Sanchez JN, Famula TR, Lyons LA, Capitanio JP. 2009. What is an "adverse" environment? Interactions of rearing experiences and MAOA genotype in rhesus monkeys. *Biological Psychiatry*, **65**(9): 770-777.
- Kaufman IC, Rosenblum LA. 1967. The reaction to separation in infant monkeys: anaclitic depression and conservation-withdrawal. *Psychosomatic Medicine*, **29**(6): 648-675.
- Kaufman J, Birmaher B, Perel J, Dahl RE, Moreci P, Nelson B, Wells W, Ryan ND. 1997. The corticotropin-releasing hormone challenge in depressed abused, depressed nonabused, and normal control children. *Biological Psychiatry*, **42**(8): 669-679.

- Kaufman J, Yang BZ, Douglas-Palumberi H, Houshyar S, Lipschitz D, Krystal JH, Gelernter J. 2004. Social supports and serotonin transporter gene moderate depression in maltreated children. *Proceedings of the National Academy of Sciences of the United States of America*, **101**(49): 17316-17321.
- Kessler RC, Davis CG, Kendler KS. 1997. Childhood adversity and adult psychiatric disorder in the US National Comorbidity Survey. *Psychological Medicine*, **27**(5): 1101-1119.
- Kessler RC. 2003. Epidemiology of women and depression. *Journal of Affective Disorders*, **74**(1): 5-13.
- Kessler RC, Wang PS. 2008. The descriptive epidemiology of commonly occurring mental disorders in the United States. *Annual Review of Public Health*, **29**: 115-129.
- Kikusui T, Mori Y. 2009. Behavioural and neurochemical consequences of early weaning in rodents. *Journal of Neuroendocrinology*, **21**(4): 427-431.
- Kim P, Evans GW, Angstadt M, Ho SS, Sripada CS, Swain JE, Liberzon I, Phan KL. 2013. Effects of childhood poverty and chronic stress on emotion regulatory brain function in adulthood. *Proceedings of the National Academy of Sciences of the United States of America*, **110**(46): 18442-18447.
- Kim-Cohen J, Caspi A, Taylor A, Williams B, Newcombe R, Craig IW, Moffitt TE. 2006. MAOA, maltreatment, and gene-environment interaction predicting children's mental health: new evidence and a meta-analysis. *Molecular Psychiatry*, **11**(10): 903-913.
- Kinnally EL, Lyons LA, Abel K, Mendoza S, Capitanio JP. 2008. Effects of early experience and genotype on serotonin transporter regulation in infant rhesus macaques. *Genes, Brain and Behavior*, **7**(4): 481-486.
- Kinnally EL, Capitanio JP, Leibel R, Deng L, LeDuc C, Haghighi F, Mann JJ. 2010. Epigenetic regulation of serotonin transporter expression and behavior in infant rhesus macaques. *Genes, Brain and Behavior*, **9**(6): 575-582.
- Kinnally EL, Capitanio JP. 2015. Paternal early experiences influence infant development through non-social mechanisms in Rhesus Macaques. *Frontiers in Zoology*, **12**(S1): S14.
- Kitayama N, Quinn S, Bremner JD. 2006. Smaller volume of anterior cingulate cortex in abuse-related posttraumatic stress disorder. *Journal of Affective Disorders*, **90**(2-3): 171-174.
- Klein B, Gorter JW, Rosenbaum P. 2013. Diagnostic shortfalls in early childhood chronic stress: a review of the issues. *Child: Care, Health and Development*, **39**(6): 765-771.
- Klimes-Dougan B, Hastings PD, Granger DA, Usher BA, Zahn-Waxler C. 2001. Adrenocortical activity in at-risk and normally developing adolescents: individual differences in salivary cortisol basal levels, diurnal variation, and responses to social challenges. *Development and Psychopathology*, **13**(3): 695-719.
- Koford CB. 1963. Rank of mothers and sons in bands of rhesus monkeys. *Science*, **141**(3578): 356-357.
- Kolber BJ, Roberts MS, Howell MP, Wozniak DF, Sands MS, Muglia LJ. 2008. Central amygdala glucocorticoid receptor action promotes fear-associated CRH activation and conditioning. *Proceedings of the National Academy of Sciences of the United States of America*, **105**(33): 12004-12009.
- Korosi A, Naninck EFG, Oomen CA, Schouten M, Krugers H, Fitzsimons C, Lucassen PJ. 2012. Early-life stress mediated modulation of adult neurogenesis and behavior. *Behavioural Brain Research*, **227**(2): 400-409.
- Koyama N. 1967. On dominance rank and kinship of a wild Japanese monkey troop in Arashiyama. *Primates*, **8**(3): 189-216.
- Kozorovitskiy Y, Gross CG, Kopil C, Battaglia L, Mcbreen M, Stranahan AM, Gould E. 2005. Experience induces structural and biochemical changes in the adult primate brain. *Proceedings of the National Academy of Sciences of the United States of America*, **102**(48): 17478-17482.
- Kraemer GW, McKinney WT. 1979. Interactions of pharmacological agents which alter biogenic amine metabolism and depression: an analysis of contributing factors within a primate model of depression. *Journal of Affective Disorders*, **1**(1): 33-54.
- Kraemer GW, Ebert MH, Lake CR, McKinney WT. 1983. Amphetamine challenge: effects in previously isolated rhesus monkeys and implications for animal models of schizophrenia. *Progress in Clinical and Biological Research*, **131**: 199-218.
- Kraemer GW, Ebert MH, Lake CR, McKinney WT. 1984. Hypersensitivity to d-amphetamine several years after early social deprivation in rhesus monkeys. *Psychopharmacology*, **82**(3): 266-271.
- Kroh K, Hageman I, Jørgensen MB. 2008. Corticotropin-releasing factor (CRF) in stress and disease: a review of literature and treatment perspectives with special emphasis on psychiatric disorders. *Nordic Journal of Psychiatry*, **62**(1): 8-16.
- Kumar S, Sachar K, Huber J, Weingarten DP, de Vellis J. 1985. Glucocorticoids regulate the transcription of glycerol phosphate dehydrogenase in cultured glial cells. *The Journal of Biological Chemistry*, **260**(27): 14743-14747.
- Kumsta R, Stevens S, Brookes K, Schlotz W, Castle J, Beckett C, Kreppner J, Rutter M, Sonuga-Barke E. 2010. 5HTT genotype moderates the influence of early institutional deprivation on emotional problems in adolescence: evidence from the English and Romanian Adoptee (ERA) study. *Journal of Child Psychology and Psychiatry*, **51**(7): 755-762.
- Kwak HR, Lee JW, Kwon KJ, Park JI, Chun W, Kim SS, Lee HJ. 2008. Neuronal architecture of the hippocampus and the cerebral cortex in rats experiencing maternal social separation. *Clinical Psychopharmacology and Neuroscience*, **6**(2): 65-70.
- Ladd CO, Owens MJ, Nemeroff CB. 1996. Persistent changes in corticotropin-releasing factor neuronal systems induced by maternal deprivation. *Endocrinology*, **137**(4): 1212-1218.
- Laudenslager ML, Boccia ML, Berger CL, Gennaro-Ruggles MM, McFerran B, Reite ML. 1995. Total cortisol, free cortisol, and growth hormone associated with brief social separation experiences in young macaques. *Developmental Psychobiology*, **28**(4): 199-211.
- Law AJ, Pei Q, Feldon J, Pryce CR, Harrison PJ. 2009a. Gene expression in the anterior cingulate cortex and amygdala of adolescent marmoset monkeys following parental separations in infancy. *The International Journal of Neuropsychopharmacology*, **12**(6): 761-772.
- Law AJ, Pei Q, Walker M, Gordon-Andrews H, Weickert CS, Feldon J, Pryce CR, Harrison PJ. 2009b. Early parental deprivation in the marmoset monkey produces long-term changes in hippocampal expression of genes involved in synaptic plasticity and implicated in mood disorder. *Neuropsychopharmacology*, **34**(6): 1381-1394.
- LeDoux J. 2003. The emotional brain, fear, and the amygdala. *Cellular and Molecular Neurobiology*, **23**(4-5): 727-738.
- Leliavski A, Dumbell R, Ott V, Oster H. 2015. Adrenal clocks and the role of

- adrenal hormones in the regulation of circadian physiology. *Journal of Biological Rhythms*, **30**(1): 20-34.
- Lenroot RK, Giedd JN. 2006. Brain development in children and adolescents: insights from anatomical magnetic resonance imaging. *Neuroscience & Biobehavioral Reviews*, **30**(6): 718-729.
- Leventopoulos M, Rüedi-Bettschen D, Knuesel I, Feldon J, Pryce CR, Opacka-Juffry J. 2007. Long-term effects of early life deprivation on brain glia in Fischer rats. *Brain Research*, **1142**: 119-126.
- Levine S, Wiener SG. 1988. Psychoendocrine aspects of mother-infant relationships in nonhuman primates. *Psychoneuroendocrinology*, **13**(1-2): 143-154.
- Levine S. 2000. Influence of psychological variables on the activity of the hypothalamic-pituitary-adrenal axis. *European Journal of Pharmacology*, **405**(1-3): 149-160.
- Levine S, Mody T. 2003. The long-term psychobiological consequences of intermittent postnatal separation in the squirrel monkey. *Neuroscience & Biobehavioral Reviews*, **27**(1-2): 83-89.
- Lim MM, Young LJ. 2006. Neuropeptidergic regulation of affiliative behavior and social bonding in animals. *Hormones and Behavior*, **50**(4): 506-517.
- Liu D, Diorio J, Tannenbaum B, Caldji C, Francis D, Freedman A, Sharma S, Pearson D, Plotsky PM, Meaney MJ. 1997. Maternal care, hippocampal glucocorticoid receptors, and hypothalamic-pituitary-adrenal responses to stress. *Science*, **277**(5332): 1659-1662.
- Liu D, Caldji C, Sharma S, Plotsky PM, Meaney MJ. 2000. Influence of neonatal rearing conditions on stress-induced adrenocorticotropin responses and norepinephrine release in the hypothalamic paraventricular nucleus. *Journal of Neuroendocrinology*, **12**(1): 5-12.
- Loman MM, Gunnar MR. 2010. Early experience and the development of stress reactivity and regulation in children. *Neuroscience & Biobehavioral Reviews*, **34**(6): 867-876.
- Lovejoy CO. 1981. The origin of man. *Science*, **211**(4480): 341-350.
- Luby J, Belden A, Botteron K, Marrus N, Harms MP, Babb C, Nishino T, Barch D. 2013. The effects of poverty on childhood brain development: the mediating effect of caregiving and stressful life events. *JAMA Pediatrics*, **167**(12): 1135-1142.
- Lukkes JL, Burke AR, Zelin NS, Hale MW, Lowry CA. 2012. Post-weaning social isolation attenuates c-Fos expression in GABAergic interneurons in the basolateral amygdala of adult female rats. *Physiology & Behavior*, **107**(5): 719-725.
- Lukkes JL, Kopelman JM, Donner NC, Hale MW, Lowry CA. 2013. Development × environment interactions control *tph2* mRNA expression. *Neuroscience*, **237**: 139-150.
- Lupien SJ, Maheu F, Tu M, Fiocco A, Schramek TE. 2007. The effects of stress and stress hormones on human cognition: implications for the field of brain and cognition. *Brain and Cognition*, **65**(3): 209-237.
- Lupien SJ, McEwen BS, Gunnar MR, Heim C. 2009. Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nature Reviews Neuroscience*, **10**(6): 434-445.
- Lupien SJ, Parent S, Evans AC, Tremblay RE, Zelazo PD, Corbo V, Pruessner JC, Séguin JR. 2011. Larger amygdala but no change in hippocampal volume in 10-year-old children exposed to maternal depressive symptomatology since birth. *Proceedings of the National Academy of Sciences of the United States of America*, **108**(34): 14324-14329.
- Lutz C, Well A, Novak M. 2003. Stereotypic and self-injurious behavior in rhesus macaques: a survey and retrospective analysis of environment and early experience. *American Journal of Primatology*, **60**(1): 1-15.
- Lutz C, Tiefenbacher S, Meyer J, Novak M. 2004. Extinction deficits in male rhesus macaques with a history of self-injurious behavior. *American Journal of Primatology*, **63**(2): 41-48.
- Lutz CK, Novak MA. 2005. Environmental enrichment for nonhuman primates: theory and application. *ILAR Journal*, **46**(2): 178-191.
- Lutz CK, Davis EB, Ruggiero AM, Suomi SJ. 2007. Early predictors of self-biting in socially-housed rhesus macaques (*Macaca mulatta*). *American Journal of Primatology*, **69**(5): 584-590.
- Lyons DM, Yang C, Mobley BW, Nickerson JT, Schatzberg AF. 2000. Early environmental regulation of glucocorticoid feedback sensitivity in young adult monkeys. *Journal of Neuroendocrinology*, **12**(8): 723-728.
- Lyons-Ruth K, Pechtel P, Yoon SA, Anderson CM, Teicher MH. 2016. Disorganized attachment in infancy predicts greater amygdala volume in adulthood. *Behavioural Brain Research*, **308**: 83-93.
- Maestripietri D, Wallen K, Carroll KA. 1997. Infant abuse runs in families of group-living pigtail macaques. *Child Abuse & Neglect*, **21**(5): 465-471.
- Maestripietri D, Carroll KA. 1998. Risk factors for infant abuse and neglect in group-living rhesus monkeys. *Psychological Science*, **9**(2): 143-145.
- Maestripietri D. 2005. Early experience affects the intergenerational transmission of infant abuse in rhesus monkeys. *Proceedings of the National Academy of Sciences of the United States of America*, **102**(27): 9726-9729.
- Maestripietri D, McCormack K, Lindell SG, Higley JD, Sanchez MM. 2006. Influence of parenting style on the offspring's behaviour and CSF monoamine metabolite levels in crossfostered and noncrossfostered female rhesus macaques. *Behavioural Brain Research*, **175**(1): 90-95.
- Magariños AM, McEwen BS, Flügge G, Fuchs E. 1996. Chronic psychosocial stress causes apical dendritic atrophy of hippocampal CA3 pyramidal neurons in subordinate tree shrews. *Journal of Neuroscience*, **16**(10): 3534-3540.
- Majer M, Nater UM, Lin JMS, Capuron L, Reeves WC. 2010. Association of childhood trauma with cognitive function in healthy adults: a pilot study. *BMC Neurology*, **10**: 61.
- Manns JR, Eichenbaum H. 2006. Evolution of declarative memory. *Hippocampus*, **16**(9): 795-808.
- Martin LJ, Spicer DM, Lewis MH, Gluck JP, Cork LC. 1991. Social deprivation of infant rhesus monkeys alters the chemoarchitecture of the brain: I. Subcortical regions. *Journal of Neuroscience*, **11**(11): 3344-3358.
- Mason WA, Sponholz RR. 1963. Behavior of Rhesus Monkeys Raised in Isolation. *Journal of Psychiatric Research*, **1**(4): 299-306.
- Mason WA. 1968. Early social deprivation in the nonhuman primates: implications for human behavior. In: Glass DS. Environmental Influences. New York: Rockefeller University Press, 70-100.
- Mason WA, Berkson G. 1975. Effects of maternal mobility on the development of rocking and other behaviors in rhesus monkeys: a study with artificial mothers. *Developmental Psychobiology*, **8**(3): 197-211.
- Maughan B, McCarthy G. 1997. Childhood adversities and psychosocial disorders. *British Medical Bulletin*, **53**(1): 156-169.
- McCormack K, Newman TK, Higley JD, Maestripietri D, Sanchez MM. 2009. Serotonin transporter gene variation, infant abuse, and responsiveness to stress in rhesus macaque mothers and infants. *Hormones and Behavior*,

55(4): 538-547.

McCrorry E, De Brito SA, Viding E. 2011. The impact of childhood maltreatment: a review of neurobiological and genetic factors. *Frontiers in Psychiatry*, **2**: 48.

McEwen BS. 1999. Stress and hippocampal plasticity. *Annual Review of Neuroscience*, **22**: 105-122.

McEwen BS. 2003. Early life influences on life-long patterns of behavior and health. *Mental Retardation and Developmental Disabilities Research Reviews*, **9**(3): 149-154.

Meaney MJ. 2001. Maternal care, gene expression, and the transmission of individual differences in stress reactivity across generations. *Annual Review of Neuroscience*, **24**: 1161-1192.

Meaney MJ, Szyf M. 2005. Maternal care as a model for experience-dependent chromatin plasticity? *Trends in Neurosciences*, **28**(9): 456-463.

Mehlman PT, Higley JD, Faucher I, Lilly AA, Taub DM, Vickers J, Suomi SJ, Linnoila M. 1995. Correlation of CSF 5-HIAA concentration with sociality and the timing of emigration in free-ranging primates. *The American Journal of Psychiatry*, **152**(6): 907-913.

Mehta MA, Golembi NI, Nosarti C, Colvert E, Mota A, Williams SCR, Rutter M, Sonuga-Barke EJS. 2009. Amygdala, hippocampal and corpus callosum size following severe early institutional deprivation: the English and Romanian Adoptees study pilot. *Journal of Child Psychology and Psychiatry*, **50**(8): 943-951.

Meyer JS, Novak MA, Bowman RE, Harlow HF. 1975. Behavioral and hormonal effects of attachment object separation in surrogate-peer-reared and mother-reared infant rhesus monkeys. *Developmental Psychobiology*, **8**(5): 425-435.

Miller GE, Chen E, Fok AK, Walker H, Lim A, Nicholls EF, Cole S, Kobor MS. 2009. Low early-life social class leaves a biological residue manifested by decreased glucocorticoid and increased proinflammatory signaling. *Proceedings of the National Academy of Sciences of the United States of America*, **106**(34): 14716-14721.

Miller RE, Mirsky IA, Caul WF, Sakata T. 1969. Hyperphagia and polydipsia in socially isolated rhesus monkeys. *Science*, **165**(3897): 1027-1028.

Miller-Butterworth CM, Kaplan JR, Barnada MM, Manuck SB, Ferrell RE. 2007. The serotonin transporter: sequence variation in *Macaca fascicularis* and its relationship to dominance. *Behavior Genetics*, **37**(5): 678-696.

Miller-Butterworth CM, Kaplan JR, Shaffer J, Devlin B, Manuck SB, Ferrell RE. 2008. Sequence variation in the primate dopamine transporter gene and its relationship to social dominance. *Molecular Biology and Evolution*, **25**(1): 18-28.

Miska EA, Ferguson-Smith AC. 2016. Transgenerational inheritance: models and mechanisms of non-DNA sequence-based inheritance. *Science*, **354**(6308): 59-63.

Mitchell GD. 1968. Persistent behavior pathology in rhesus monkeys following early social isolation. *Folia Primatologica*, **8**(2): 132-147.

Morecraft RJ, McNeal DW, Stilwell-Morecraft KS, Gedney M, Ge JZ, Schroeder CM, van Hoesen GW. 2007. Amygdala interconnections with the cingulate motor cortex in the rhesus monkey. *The Journal of Comparative Neurology*, **500**(1): 134-165.

Morey RA, Haswell CC, Hooper SR, De Bellis MD. 2016. Amygdala, hippocampus, and ventral medial prefrontal cortex volumes differ in maltreated youth with and without chronic posttraumatic stress disorder. *Neuropsychopharmacology*, **41**(3): 791-801.

Morgan D, Grant KA, Pringle OA, Nader SH, Kaplan JR, Nader MA. 2000. Predictors of social status in cynomolgus monkeys (*Macaca fascicularis*) after group formation. *American Journal of Primatology*, **52**(3): 115-131.

Musholt K, Cirillo G, Cavaliere C, Rosaria Bianco M, Bock J, Helmeke C, Braun K, Papa M. 2009. Neonatal separation stress reduces glial fibrillary acidic protein- and S100 β -immunoreactive astrocytes in the rat medial precentral cortex. *Developmental Neurobiology*, **69**(4): 203-211.

Nagy C, Turecki G. 2015. Transgenerational epigenetic inheritance: an open discussion. *Epigenomics*, **7**(5): 781-790.

Nelson EE, Herman KN, Barrett CE, Noble PL, Wojteczko K, Chisholm K, Delaney D, Ernst M, Fox NA, Suomi SJ, Winslow JT, Pine DS. 2009. Adverse rearing experiences enhance responding to both aversive and rewarding stimuli in juvenile rhesus monkeys. *Biological Psychiatry*, **66**(7): 702-704.

Newman TK, Syagailo YV, Barr CS, Wendland JR, Champoux M, Graessle M, Suomi SJ, Higley JD, Lesch KP. 2005. Monoamine oxidase A gene promoter variation and rearing experience influences aggressive behavior in rhesus monkeys. *Biological Psychiatry*, **57**(2): 167-172.

Newton AW, Vandeven AM. 2009. Update on child maltreatment. *Current Opinion in Pediatrics*, **21**(2): 252-261.

Newton AW, Vandeven AM. 2010. Child abuse and neglect: a worldwide concern. *Current Opinion in Pediatrics*, **22**(2): 226-233.

Nichols NR, Agolley D, Zieba M, Bye N. 2005. Glucocorticoid regulation of glial responses during hippocampal neurodegeneration and regeneration. *Brain Research Reviews*, **48**(2): 287-301.

Novak MA, Suomi SJ. 1991. Social interaction in nonhuman primates: an underlying theme for primate research. *Laboratory Animal Science*, **41**(4): 308-314.

Novak MA, Kinsey JH, Jorgensen MJ, Hazen TJ. 1998. Effects of puzzle feeders on pathological behavior in individually housed rhesus monkeys. *American Journal of Primatology*, **46**(3): 213-227.

Novak MFSX, Sackett GP. 1997. Pair-rearing infant monkeys (*Macaca nemestrina*) using a "rotating-peer" strategy. *American Journal of Primatology*, **41**(2): 141-149.

O'Callaghan JP, Brinton RE, McEwen BS. 1989. Glucocorticoids regulate the concentration of glial fibrillary acidic protein throughout the brain. *Brain Research*, **494**(1): 159-161.

Ohmura Y, Yoshioka M. 2009. The roles of corticotropin releasing factor (CRF) in responses to emotional stress: is CRF release a cause or result of fear/anxiety? *CNS & Neurological Disorders - Drug Targets*, **8**(6): 459-469.

Öngür D, Drevets WC, Price JL. 1998. Glial reduction in the subgenual prefrontal cortex in mood disorders. *Proceedings of the National Academy of Sciences of the United States of America*, **95**(22): 13290-13295.

Ono M, Kikusui T, Sasaki N, Ichikawa M, Mori Y, Murakami-Murofushi K. 2008. Early weaning induces anxiety and precocious myelination in the anterior part of the basolateral amygdala of male Balb/c mice. *Neuroscience*, **156**(4): 1103-1110.

Oomen CA, Girardi CEN, Cahyadi R, Verbeek EC, Krugers H, Joëls M, Lucassen PJ. 2009. Opposite effects of early maternal deprivation on neurogenesis in male versus female rats. *PLoS One*, **4**(1): e3675.

Opitz B. 2014. Memory function and the hippocampus. *Frontiers of Neurology and Neuroscience*, **34**: 51-59.

Oswald LM, Wand GS, Kuwabara H, Wong DF, Zhu SJ, Brasic JR. 2014. History of childhood adversity is positively associated with ventral striatal

- dopamine responses to amphetamine. *Psychopharmacology*, **231**(12): 2417-2433.
- Parker KJ, Buckmaster CL, Schatzberg AF, Lyons DM. 2004. Prospective investigation of stress inoculation in young monkeys. *Archives of General Psychiatry*, **61**(9): 933-941.
- Parker KJ, Buckmaster CL, Justus KR, Schatzberg AF, Lyons DM. 2005. Mild early life stress enhances prefrontal-dependent response inhibition in monkeys. *Biological Psychiatry*, **57**(8): 848-855.
- Parker KJ, Buckmaster CL, Lindley SE, Schatzberg AF, Lyons DM. 2012. Hypothalamic-pituitary-adrenal axis physiology and cognitive control of behavior in stress inoculated monkeys. *International Journal of Behavioral Development*, **36**(1), doi: 10.1177/0165025411406864.
- Payne C, Machado CJ, Bliwise NG, Bachevalier J. 2010. Maturation of the hippocampal formation and amygdala in *Macaca mulatta*: a volumetric magnetic resonance imaging study. *Hippocampus*, **20**(8): 922-935.
- Pesonen AK, Räikkönen K, Feldt K, Heinonen K, Osmond C, Phillips DIW, Barker DJP, Eriksson JG, Kajantie E. 2010. Childhood separation experience predicts HPA axis hormonal responses in late adulthood: a natural experiment of World War II. *Psychoneuroendocrinology*, **35**(5): 758-767.
- Phelps EA. 2006. Emotion and cognition: insights from studies of the human amygdala. *Annual Review of Psychology*, **57**: 27-53.
- Pirkola S, Isometsä E, Aro H, Kestilä L, Hämäläinen J, Veijola J, Kiviruusu O, Lönnqvist J. 2005. Childhood adversities as risk factors for adult mental disorders: results from the Health 2000 study. *Social Psychiatry and Psychiatric Epidemiology*, **40**(10): 769-777.
- Plotsky PM, Thiruvikraman KV, Nemeroff CB, Caldji C, Sharma S, Meaney MJ. 2005. Long-term consequences of neonatal rearing on central corticotropin-releasing factor systems in adult male rat offspring. *Neuropsychopharmacology*, **30**(12): 2192-2204.
- Provençal N, Suderman MJ, Guillemin C, Massart R, Ruggiero A, Wang D, Bennett AJ, Pierre PJ, Friedman DP, Côté SM, Hallett M, Tremblay RE, Suomi SJ, Szyf M. 2012. The signature of maternal rearing in the methylome in rhesus macaque prefrontal cortex and T cells. *Journal of Neuroscience*, **32**(44): 15626-15642.
- Provençal N, Booij L, Tremblay RE. 2015. The developmental origins of chronic physical aggression: biological pathways triggered by early life adversity. *The Journal of Experimental Biology*, **218**(Pt 1): 123-133.
- Pryce CR, Dettling A, Spengler M, Spaete C, Feldon J. 2004a. Evidence for altered monoamine activity and emotional and cognitive disturbance in marmoset monkeys exposed to early life stress. *Annals of the New York Academy of Sciences*, **1032**: 245-249.
- Pryce CR, Dettling AC, Spengler M, Schnell CR, Feldon J. 2004b. Deprivation of parenting disrupts development of homeostatic and reward systems in marmoset monkey offspring. *Biological Psychiatry*, **56**(2): 72-79.
- Pryce CR, Feldon J, Fuchs E, Knuesel I, Oertle T, Sengstag C, Spengler M, Weber E, Weston A, Jongen-Rélo A. 2005a. Postnatal ontogeny of hippocampal expression of the mineralocorticoid and glucocorticoid receptors in the common marmoset monkey. *European Journal of Neuroscience*, **21**(6): 1521-1535.
- Pryce CR, Rüedi-Bettschen D, Dettling AC, Weston A, Russig H, Ferger B, Feldon J. 2005b. Long-term effects of early-life environmental manipulations in rodents and primates: potential animal models in depression research. *Neuroscience & Biobehavioral Reviews*, **29**(4-5): 649-674.
- Pryce CR, Aubert Y, Maier C, Pearce PC, Fuchs E. 2011. The developmental impact of prenatal stress, prenatal dexamethasone and postnatal social stress on physiology, behaviour and neuroanatomy of primate offspring: studies in rhesus macaque and common marmoset. *Psychopharmacology*, **214**(1): 33-53.
- Raleigh MJ, Brammer GL, McGuire MT. 1983. Male dominance, serotonergic systems, and the behavioral and physiological effects of drugs in vervet monkeys (*Cercopithecus aethiops sabaeus*). *Progress in Clinical and Biological Research*, **131**: 185-197.
- Raleigh MJ, Brammer GL, McGuire MT, Yuwiler A. 1985. Dominant social status facilitates the behavioral effects of serotonergic agonists. *Brain Research*, **348**(2): 274-282.
- Raleigh MJ, McGuire MT, Brammer GL, Pollack DB, Yuwiler A. 1991. Serotonergic mechanisms promote dominance acquisition in adult male vervet monkeys. *Brain Research*, **559**(2): 181-190.
- Rauch SL, Whalen PJ, Shin LM, McInerney SC, Macklin ML, Lasko NB, Orr SP, Pitman RK. 2000. Exaggerated amygdala response to masked facial stimuli in posttraumatic stress disorder: a functional MRI study. *Biological Psychiatry*, **47**(9): 769-776.
- Reite M, Short RA. 1978. Nocturnal sleep in separated monkey infants. *Archives of General Psychiatry*, **35**(10): 1247-1253.
- Reite M, Kaufman CI, Pauley DJ, Styne AJ. 1974. Depression in infant monkeys: physiological correlates. *Psychosomatic Medicine*, **36**(4): 363-367.
- Richards JE, Xie WZ. 2015. Brains for all the ages: structural neurodevelopment in infants and children from a life-span perspective. *Advances in Child Development and Behavior*, **48**: 1-52.
- Richert KA, Carrion VG, Karchemskiy A, Reiss AL. 2006. Regional differences of the prefrontal cortex in pediatric PTSD: an MRI study. *Depression and Anxiety*, **23**(1): 17-25.
- Rilling JK, Winslow JT, O'Brien D, Gutman DA, Hoffman JM, Kilts CD. 2001. Neural correlates of maternal separation in rhesus monkeys. *Biological Psychiatry*, **49**(2): 146-157.
- Rinne-Albers MAW, van der Werff SJA, van Hoof MJ, van Lang ND, Lamers-Winkelmann F, Rombouts SA, Vermeiren RRJM, van der Wee NJA. 2016. Abnormalities of white matter integrity in the corpus callosum of adolescents with PTSD after childhood sexual abuse: a DTI study. *European Child & Adolescent Psychiatry*, **25**(8): 869-878.
- Roceri M, Hendriks W, Racagni G, Ellenbroek BA, Riva MA. 2002. Early maternal deprivation reduces the expression of BDNF and NMDA receptor subunits in rat hippocampus. *Molecular Psychiatry*, **7**(6): 609-616.
- Rodgers AB, Morgan CP, Bronson SL, Revello S, Bale TL. 2013. Paternal stress exposure alters sperm microRNA content and reprograms offspring HPA stress axis regulation. *Journal of Neuroscience*, **33**(21): 9003-9012.
- Romans SE, Casement MD, McAloon R, Keenan K, Hipwell AE, Guyer AE, Forbes EE. 2015. Adolescent girls' neural response to reward mediates the relation between childhood financial disadvantage and depression. *Journal of Child Psychology and Psychiatry*, **56**(11): 1177-1184.
- Rommeck I, Anderson K, Heagerty A, Cameron A, McCowan B. 2009a. Risk factors and remediation of self-injurious and self-abuse behavior in rhesus macaques. *Journal of Applied Animal Welfare Science*, **12**(1): 61-72.
- Rommeck I, Gottlieb DH, Strand SC, McCowan B. 2009b. The effects of four nursery rearing strategies on infant behavioral development in rhesus

- macaques (*Macaca mulatta*). *Journal of the American Association for Laboratory Animal Science*, **48**(4): 395-401.
- Rommeck I, Capitanio JP, Strand SC, McCowan B. 2011. Early social experience affects behavioral and physiological responsiveness to stressful conditions in infant rhesus macaques (*Macaca mulatta*). *American Journal of Primatology*, **73**(7): 692-701.
- Rosenblum LA, Pausly GS. 1984. The effects of varying environmental demands on maternal and infant behavior. *Child Development*, **55**(1): 305-314.
- Rosenblum LA, Andrews MW. 1994. Influences of environmental demand on maternal behavior and infant development. *Acta Paediatrica*, **83**(S397): 57-63.
- Rosenblum LA, Coplan JD, Friedman S, Bassoff T, Gorman JM, Andrews MW. 1994. Adverse early experiences affect noradrenergic and serotonergic functioning in adult primates. *Biological Psychiatry*, **35**(4): 221-227.
- Roustit C, Renahy E, Guernec G, Lesieur S, Parizot I, Chauvin P. 2009. Exposure to interparental violence and psychosocial maladjustment in the adult life course: advocacy for early prevention. *Journal of Epidemiology & Community Health*, **63**(7): 563-568.
- Roy MA, Wolf RH, Martin LN, Rangan SR, Allen WP. 1978. Social and reproductive behaviors in surrogate-reared squirrel monkeys (*Saimiri sciureus*). *Laboratory Animal Science*, **28**(4): 417-421.
- Rüedi-Bettschen D, Zhang WN, Russig H, Ferger B, Weston A, Pedersen EM, Feldon J, Pryce CR. 2006. Early deprivation leads to altered behavioural, autonomic and endocrine responses to environmental challenge in adult Fischer rats. *European Journal of Neuroscience*, **24**(10): 2879-2893.
- Ruppenthal GC, Walker CG, Sackett GP. 1991. Rearing infant monkeys (*Macaca nemestrina*) in pairs produces deficient social development compared with rearing in single cages. *American Journal of Primatology*, **25**(2): 103-113.
- Rutter M. 2007. Resilience, competence, and coping. *Child Abuse & Neglect*, **31**(3): 205-209.
- Sabatini MJ, Ebert P, Lewis DA, Levitt P, Cameron JL, Mirnics K. 2007. Amygdala gene expression correlates of social behavior in monkeys experiencing maternal separation. *Journal of Neuroscience*, **27**(12): 3295-3304.
- Sackett GP. 1967. Some persistent effects of different rearing conditions on preadult social behavior of monkeys. *Journal of Comparative and Physiological Psychology*, **64**(2): 363-365.
- Sackett GP. 1969. The persistence of abnormal behaviour in monkeys following isolation rearing. *International Psychiatry Clinics*, **6**(1): 3-37.
- Sánchez MM, Hearn EF, Do D, Rilling JK, Herndon JG. 1998. Differential rearing affects corpus callosum size and cognitive function of rhesus monkeys. *Brain Research*, **812**(1-2): 38-49.
- Sánchez MM, Young LJ, Plotsky PM, Insel TR. 2000. Distribution of corticosteroid receptors in the rhesus brain: relative absence of glucocorticoid receptors in the hippocampal formation. *Journal of Neuroscience*, **20**(12): 4657-4668.
- Sánchez MM, Ladd CO, Plotsky PM. 2001. Early adverse experience as a developmental risk factor for later psychopathology: evidence from rodent and primate models. *Development and Psychopathology*, **13**(3): 419-449.
- Sánchez MM, Noble PM, Lyon CK, Plotsky PM, Davis M, Nemeroff CB, Winslow JT. 2005. Alterations in diurnal cortisol rhythm and acoustic startle response in nonhuman primates with adverse rearing. *Biological Psychiatry*, **57**(4): 373-381.
- Sanchez MM. 2006. The impact of early adverse care on HPA axis development: nonhuman primate models. *Hormones and Behavior*, **50**(4): 623-631.
- Sapolsky RM, Krey LC, McEwen BS. 1985. Prolonged glucocorticoid exposure reduces hippocampal neuron number: implications for aging. *Journal of Neuroscience*, **5**(5): 1222-1227.
- Schneider ML, Suomi SJ. 1992. Neurobehavioral assessment in rhesus monkey neonates (*Macaca mulatta*): developmental changes, behavioral stability, and early experience. *Infant Behavior and Development*, **15**(2): 155-177.
- Schweizer S, Walsh ND, Stretton J, Dunn VJ, Goodyer IM, Dalgleish T. 2016. Enhanced emotion regulation capacity and its neural substrates in those exposed to moderate childhood adversity. *Social Cognitive and Affective Neuroscience*, **11**(2): 272-281.
- Seay B, Hansen E, Harlow HF. 1962. Mother-infant separation in monkeys. *Journal of Child Psychology and Psychiatry*, **3**(3-4): 123-132.
- Seay B, Alexander BK, Harlow HF. 1964. Maternal behavior of socially deprived Rhesus monkeys. *The Journal of Abnormal and Social Psychology*, **69**(4): 345-354.
- Shannon C, Champoux M, Suomi SJ. 1998. Rearing condition and plasma cortisol in rhesus monkey infants. *American Journal of Primatology*, **46**(4): 311-321.
- Shannon C, Schwandt ML, Champoux M, Shoaf SE, Suomi SJ, Linnoila M, Higley JD. 2005. Maternal absence and stability of individual differences in CSF 5-HIAA concentrations in rhesus monkey infants. *American Journal of Psychiatry*, **162**(9): 1658-1664.
- Sheridan M, Drury S, McLaughlin K, Almas A. 2010. Early institutionalization: neurobiological consequences and genetic modifiers. *Neuropsychology Review*, **20**(4): 414-429.
- Shishkina GT, Kalinina TS, Bulygina VV, Lanshakov DA, Babluk EV, Dygalo NN. 2015. Anti-apoptotic protein Bcl-xL expression in the midbrain raphe region is sensitive to stress and glucocorticoids. *PLoS One*, **10**(12): e0143978.
- Shohamy D, Turk-Browne NB. 2013. Mechanisms for widespread hippocampal involvement in cognition. *Journal of Experimental Psychology. General*, **142**(4): 1159-1170.
- Siegel SJ, Ginsberg SD, Hof PR, Foote SL, Young WG, Kraemer GW, McKinney WT, Morrison JH. 1993. Effects of social deprivation in prepubescent rhesus monkeys: immunohistochemical analysis of the neurofilament protein triplet in the hippocampal formation. *Brain Research*, **619**(1-2): 299-305.
- Singh M, D'Souza L, Singh M. 1992. Hierarchy, kinship and social interaction among Japanese monkeys (*Macaca fuscata*). *Journal of Biosciences*, **17**(1): 15-27.
- Smagin GN, Dunn AJ. 2000. The role of CRF receptor subtypes in stress-induced behavioural responses. *European Journal of Pharmacology*, **405**(1-3): 199-206.
- Spencer-Booth Y, Hinde RA. 1971. The effects of 13 days maternal separation on infant rhesus monkeys compared with those of shorter and repeated separations. *Animal Behaviour*, **19**(3): 595-605.
- Spinelli S, Schwandt ML, Lindell SG, Newman TK, Heilig M, Suomi SJ,

- Higley JD, Goldman D, Barr CS. 2007. Association between the recombinant human serotonin transporter linked promoter region polymorphism and behavior in rhesus macaques during a separation paradigm. *Development and Psychopathology*, **19**(4): 977-987.
- Spinelli S, Chefer S, Suomi SJ, Higley JD, Barr CS, Stein E. 2009. Early-life stress induces long-term morphologic changes in primate brain. *Archives of General Psychiatry*, **66**(6): 658-665.
- Sprague DS. 1998. Age, dominance rank, natal status, and tenure among male macaques. *American Journal of Physical Anthropology*, **105**(4): 511-521.
- Stavisky RC, Adams MR, Watson SL, Kaplan JR. 2001. Dominance, cortisol, and behavior in small groups of female cynomolgus monkeys (*Macaca fascicularis*). *Hormones and Behavior*, **39**(3): 232-238.
- Stein MB, Koverola C, Hanna C, Torchia MG, McClarty B. 1997. Hippocampal volume in women victimized by childhood sexual abuse. *Psychological Medicine*, **27**(4): 951-959.
- Stell M, Riesen A. 1987. Effects of early environments on monkey cortex neuroanatomical changes following somatomotor experience: effects on layer III pyramidal cells in monkey cortex. *Behavioral Neuroscience*, **101**(3): 341-346.
- Struble RG, Riesen AH. 1978. Changes in cortical dendritic branching subsequent to partial social isolation in stump-tailed monkeys. *Developmental Psychobiology*, **11**(5): 479-486.
- Suomi SJ, Harlow HF, Kimball SD. 1971. Behavioral effects of prolonged partial social isolation in the rhesus monkey. *Psychological Reports*, **29**(3S): 1171-1177.
- Suomi SJ. 1973. Surrogate rehabilitation of monkeys reared in total social isolation. *Journal of Child Psychology and Psychiatry*, **14**(1): 71-77.
- Suomi SJ. 1978. Maternal behavior by socially incompetent monkeys: neglect and abuse of offspring. *Journal of Pediatric Psychology*, **3**(1): 28-34.
- Suomi SJ, Harlow HF, Novak MA. 1974. Reversal of social deficits produced by isolation rearing in monkeys. *Journal of Human Evolution*, **3**(6): 527-534.
- Suomi SJ, Mineka S, DeLizio RD. 1983. Short- and long-term effects of repetitive mother-infant separations on social development in rhesus monkeys. *Developmental Psychology*, **19**(5): 770-786.
- Suomi SJ, Ripp C. 1983. A history of mother-less mother monkey mothering at the University of Wisconsin Primate Laboratory. In: Reite M, Caine N. Child Abuse: The Nonhuman Primate Data. New York: Alan R Liss, 50-78.
- Suomi SJ. 1991. Early stress and adult emotional reactivity in rhesus monkeys. *Ciba Foundation Symposium*, **156**: 171-183; discussion 183-188. (请核对页码)
- Suomi SJ. 1997. Early determinants of behaviour: evidence from primate studies. *British Medical Bulletin*, **53**(1): 170-184.
- Suomi SJ. 2005. Mother-infant attachment, peer relationships, and the development of social networks in rhesus monkeys. *Human Development*, **48**(1-2): 67-79.
- Takahashi H. 2002. Changes of dominance rank, age, and tenure of wild Japanese macaque males in the kinkazan a troop during seven years. *Primates*, **43**(2): 133-138.
- Teicher MH, Ito Y, Glod CA, Andersen SL, Dumont N, Ackerman E. 1997. Preliminary evidence for abnormal cortical development in physically and sexually abused children using EEG coherence and MRI. *Annals of the New York Academy of Sciences*, **821**: 160-175.
- Teicher MH, Dumont NL, Ito Y, Vaituzis C, Giedd JN, Andersen SL. 2004. Childhood neglect is associated with reduced corpus callosum area. *Biological Psychiatry*, **56**(2): 80-85.
- Thomae K, Dorrepaal E, Draijer N, De Ruiter MB, Van Balkom AJ, Smit JH, Veltman DJ. 2010. Reduced anterior cingulate and orbitofrontal volumes in child abuse-related complex PTSD. *The Journal of Clinical Psychiatry*, **71**(12): 1636-1644.
- Thomas BW, Champoux M, Suomi SJ, Gunnar MR. 1995. Salivary cortisol in nursery-reared rhesus monkeys: reactivity to peer interactions and altered circadian activity. *Developmental Psychobiology*, **28**(5): 257-267.
- Tomoda A, Suzuki H, Rabi K, Sheu YS, Polcari A, Teicher MH. 2009. Reduced prefrontal cortical gray matter volume in young adults exposed to harsh corporal punishment. *NeuroImage*, **47**(S2): T66-T71.
- Tottenham N, Sheridan MA. 2009. A review of adversity, the amygdala and the hippocampus: a consideration of developmental timing. *Frontiers in Human Neuroscience*, **3**: 68.
- Tottenham N, Hare TA, Quinn BT, McCarry TW, Nurse M, Gilhooly T, Millner A, Galvan A, Davidson MC, Eigsti IM, Thomas KM, Freed PJ, Booma ES, Gunnar MR, Altemus M, Aronson J, Casey BJ. 2010. Prolonged institutional rearing is associated with atypically large amygdala volume and difficulties in emotion regulation. *Developmental Science*, **13**(1): 46-61.
- Treadway MT, Grant MM, Ding ZH, Hollon SD, Gore JC, Shelton RC. 2009. Early adverse events, HPA activity and rostral anterior cingulate volume in MDD. *PLoS One*, **4**(3): e4887.
- Tsang AH, Barclay JL, Oster H. 2014. Interactions between endocrine and circadian systems. *Journal of Molecular Endocrinology*, **52**(1): R1-16.
- Tsang AH, Astiz M, Friedrichs M, Oster H. 2016. Endocrine regulation of circadian physiology. *Journal of Endocrinology*, **230**(1): R1-R11.
- Uylings HBM, van Eden CG. 1991. Qualitative and quantitative comparison of the prefrontal cortex in rat and in *Primates*, including humans. *Progress in Brain Research*, **85**: 31-62.
- Vaiserman A. 2015a. Epidemiologic evidence for association between adverse environmental exposures in early life and epigenetic variation: a potential link to disease susceptibility? *Clinical Epigenetics*, **7**: 96.
- Vaiserman AM. 2015b. Epigenetic programming by early-life stress: evidence from human populations. *Developmental Dynamics*, **244**(3): 254-265.
- van Haarst AD, Oitzl MS, de Kloet ER. 1997. Facilitation of feedback inhibition through blockade of glucocorticoid receptors in the hippocampus. *Neurochemical Research*, **22**(11): 1323-1328.
- van Harmelen AL, van Tol MJ, van der Wee NJA, Veltman DJ, Aleman A, Spinhoven P, van Buchem MA, Zitman FG, Penninx BWJH, Elzinga BM. 2010. Reduced medial prefrontal cortex volume in adults reporting childhood emotional maltreatment. *Biological Psychiatry*, **68**(9): 832-838.
- Vazquez DM, Bailey C, Dent GW, Okimoto DK, Steffek A, López JF, Levine S. 2006. Brain corticotropin-releasing hormone (CRH) circuits in the developing rat: effect of maternal deprivation. *Brain Research*, **1121**(1): 83-94.
- Vicentic A, Francis D, Moffett M, Lakatos A, Rogge G, Hubert GW, Harley J, Kuhar MJ. 2006. Maternal separation alters serotonergic transporter densities and serotonergic 1A receptors in rat brain. *Neuroscience*, **140**(1): 355-365.
- Vielkind U, Walencewicz A, Levine JM, Bohn MC. 1990. Type II glucocorticoid receptors are expressed in oligodendrocytes and astrocytes.

Journal of Neuroscience Research, **27**(3): 360-373.

Volavka J, Bilder R, Nolan K. 2004. Catecholamines and aggression: the role of COMT and MAO polymorphisms. *Annals of the New York Academy of Sciences*, **1036**: 393-398.

Wakefield CL, Levine MS. 1985. Early postnatal development of basolateral amygdala in kitten: a Golgi morphometric analysis. *Brain Research Bulletin*, **14**(2): 159-167.

Wall VL, Fischer EK, Bland ST. 2012. Isolation rearing attenuates social interaction-induced expression of immediate early gene protein products in the medial prefrontal cortex of male and female rats. *Physiology & Behavior*, **107**(3): 440-450.

Wallen K, Goldfoot DA, Goy RW. 1981. Peer and maternal influences on the expression of foot-clasp mounting by juvenile male rhesus monkeys. *Developmental Psychobiology*, **14**(4): 299-309.

Wang X, Xie H, Cotton AS, Duval ER, Tamburrino MB, Brickman KR, Elhai JD, Ho SS, McLean SA, Ferguson EJ, Liberzon I. 2016. Preliminary Study of Acute Changes in Emotion Processing in Trauma Survivors with PTSD Symptoms. *PLoS One*, **11**(7): e0159065.

Washburn DA, Rumbaugh DM. 1991. Impaired performance from brief social isolation of rhesus monkeys (*Macaca mulatta*): a multiple video-task assessment. *Journal of Comparative Psychology*, **105**(2): 145-151.

Watanabe SE, Donzella B, Kertes DA, Gunnar MR. 2004. Developmental changes in baseline cortisol activity in early childhood: relations with napping and effortful control. *Developmental Psychobiology*, **45**(3): 125-133.

Weaver ICG, Cervoni N, Champagne FA, D'Alessio AC, Sharma S, Seckl JR, Dymov S, Szyf M, Meaney MJ. 2004. Epigenetic programming by maternal behavior. *Nature Neuroscience*, **7**(8): 847-854.

Weaver ICG. 2007. Epigenetic programming by maternal behavior and pharmacological intervention *Nature versus nurture: let's call the whole thing off*. *Epigenetics*, **2**(1): 22-28.

White SF, Costanzo ME, Blair JR, Roy MJ. 2015. PTSD symptom severity is associated with increased recruitment of top-down attentional control in a trauma-exposed sample. *Neuroimage: Clinical*, **7**: 19-27.

Winslow JT, Noble PL, Lyons CK, Sterk SM, Insel TR. 2003. Rearing effects on cerebrospinal fluid oxytocin concentration and social buffering in rhesus monkeys. *Neuropsychopharmacology*, **28**(5): 910-918.

Woon FL, Hedges DW. 2008. Hippocampal and amygdala volumes in children and adults with childhood maltreatment-related posttraumatic stress disorder: a meta-analysis. *Hippocampus*, **18**(8): 729-736.

Worlein JM, Sackett GP. 1997. Social development in nursery-reared

pigtailed macaques (*Macaca nemestrina*). *American Journal of Primatology*, **41**(1): 23-35.

Worlein JM. 2014. Nonhuman primate models of depression: effects of early experience and stress. *ILAR Journal*, **55**(2): 259-273.

Wright JC. 1983. The effects of differential rearing on exploratory behavior in puppies. *Applied Animal Ethology*, **10**(1-2): 27-34.

Wright PC. 1990. Patterns of paternal care in primates. *International Journal of Primatology*, **11**(2): 89-102.

Yamasue H, Kasai K, Iwanami A, Ohtani T, Yamada H, Abe O, Kuroki N, Fukuda R, Tochigi M, Furukawa S, Sadamatsu M, Sasaki T, Aoki S, Ohtomo K, Asukai N, Kato N. 2003. Voxel-based analysis of MRI reveals anterior cingulate gray-matter volume reduction in posttraumatic stress disorder due to terrorism. *Proceedings of the National Academy of Sciences of the United States of America*, **100**(15): 9039-9043.

Yuan TF, Li A, Sun X, Ouyang H, Campos C, Rocha NBF, Arias-Carrión O, Machado S, Hou GL, So KF. 2016. Transgenerational inheritance of paternal neurobehavioral phenotypes: stress, addiction, ageing and metabolism. *Molecular Neurobiology*, **53**(9): 6367-6376.

Zarrow MX, Philpott JE, Denenberg VH. 1970. Passage of ¹⁴C-4-corticosterone from the rat mother to the foetus and neonate. *Nature*, **226**(5250): 1058-1059.

Zhang TY, Bagot R, Parent C, Nesbitt C, Bredy TW, Caldji C, Fish E, Anisman H, Szyf M, Meaney MJ. 2006. Maternal programming of defensive responses through sustained effects on gene expression. *Biological Psychology*, **73**(1): 72-89.

Zhang TY, Meaney MJ. 2010. Epigenetics and the environmental regulation of the genome and its function. *Annual Review of Psychology*, **61**: 439-466.

Zheng J, Xiao XH, Zhang Q, Yu M. 2014. DNA methylation: the pivotal interaction between early-life nutrition and glucose metabolism in later life. *British Journal of Nutrition*, **112**(11): 1850-1857.

Ziabreva I, Schnabel R, Braun K. 2000. Parental deprivation induces N-methyl-D-aspartate-receptor upregulation in limbic brain areas of *Octodon degus*: protective role of the maternal call. *Neural Plasticity*, **7**(4): 233-244.

Ziabreva I, Poeggel G, Schnabel R, Braun K. 2003a. Separation-induced receptor changes in the hippocampus and amygdala of *Octodon degus*: influence of maternal vocalizations. *Journal of Neuroscience*, **23**(12): 5329-5336.

Ziabreva I, Schnabel R, Poeggel G, Braun K. 2003b. Mother's voice "buffers" separation-induced receptor changes in the prefrontal cortex of *octodon degus*. *Neuroscience*, **119**(2): 433-441.

Does mRNA structure contain genetic information for regulating co-translational protein folding?

Jian-Rong Yang*

Department of Biology, Zhongshan School of Medicine, Sun Yat-sen University, Guangzhou 510080, China

ABSTRACT

Currently many facets of genetic information are ill-defined. In particular, how protein folding is genetically regulated has been a long-standing issue for genetics and protein biology. And a generic mechanistic model with supports of genomic data is still lacking. Recent technological advances have enabled much needed genome-wide experiments. While putting the effect of codon optimality on debate, these studies have supplied mounting evidence suggesting a role of mRNA structure in the regulation of protein folding by modulating translational elongation rate. In conjunctions with previous theories, this mechanistic model of protein folding guided by mRNA structure shall expand our understandings of genetic information and offer new insights into various biomedical puzzles.

Keywords: Translational elongation rate; Protein folding; mRNA secondary structure; Codon usage bias

INTRODUCTION

With regards to the full contents of genetic information, the answer to this fundamental question in biology has been frequently updated as newly emerging techniques and growing data constantly challenge our existing understandings (Ramos & Laederach, 2014). On one hand, novel functions of non-coding genome are uncovered (ENCODE Project Consortium, 2012). On the other hand, our understandings of genetic information in coding regions have also extended beyond canonical schemes of how protein folding is regulated within cells.

Commonly known as “the second half of genetic code” (Kolata, 1986), a vast pool of information is required for ensuring correct folding of polypeptide into its native structure. However, little is known about how information stored in nucleotide sequence is transmitted from genome into polypeptide chain. According to the Central Dogma, messenger RNA is frequently targeted for searching regulatory signals for protein folding. Indeed, as

evidenced by unequal usage of synonymous codons and its correlation with efficiency and/or accuracy of translational elongation (Gingold & Pilpel, 2011), mRNA molecule obviously contains more information than primary protein sequence. This logic and decades of genomic sequencing have elucidated the association between codon usage bias and protein structures (Spencer & Barral, 2012; Tsai et al., 2008).

Nevertheless, secondary structure of mRNA is often overlooked, probably due to a lack of scalable experiments for detecting RNA structure (Eddy, 2014) and the complexity of *in silico* prediction for the structure of ribosome-bound mRNA. Recently, this viewpoint has altered due to the latest technical innovations, especially ribosome profiling (Ingolia et al., 2009) and several high-throughput assays for mRNA secondary structure (Graveley, 2016). As a result, a novel regulatory role of mRNA structure on protein folding emerges.

In this review, current models of co-translational protein folding were reviewed for elucidating the generic molecular mechanism for its linkage to mRNA structure. Several preliminary studies that correlate computationally predicted mRNA structures with protein conformation shall be discussed. A major focus was placed upon regulatory signals in major mRNA coding sequences rather than a specific mRNA fragment such as translational ramp at 5' end (Tuller et al., 2010). Furthermore, biologically relevant interpretations of this regulation were offered.

GENETIC INFORMATION GUIDING PROTEIN FOLDING

The complexity of protein folding have been conventionally summarized as the Levinthal's paradox. It states that the number of possible conformations of a small protein (around 100 residues) was so large that it would require more time than

Received: 01 December 2016; Accepted: 04 January 2017

Foundation items: This study was supported by the start-up grant from “Top 100 Talents Program” of Sun Yat-sen University to JRY (50000-31131114) and General Program of National Natural Science Foundation of China to JRY (31671320)

*Corresponding author, E-mail: yangjr27@mail.sysu.edu.cn

DOI:10.13918/j.issn.2095-8137.2017.004

the lifespan of universe (10^{16} seconds) to explore all possibilities and choose the native conformation (Zwanzig et al., 1992). How this astronomic eternity is reduced to biologically feasible range has been a long-standing puzzle of protein folding. As a theoretic milestone of protein folding, the Anfinsen's dogma, also known as thermodynamic hypothesis, suggests that protein conformation is solely determined by its amino acid sequence. In other words, assuming the validity of the Anfinsen's dogma, accurate protein folding requires no additional genetic input other than primary protein sequence. Nevertheless, multiple exceptions to the Anfinsen's dogma were detected later, including but not limited to prion (Fraser, 2014) and kinetically stable proteins (Xia et al., 2007). It has confirmed the existence of regulatory signals that guide protein folding, including *trans* factors such as chaperones and *cis* factors such as codon optimality (see below).

There are two major modes of protein folding. On the one hand, protein folding occurs after the entire coding sequence has been fully translated. Upon dissociation of mRNA with synthesized polypeptide, a generic molecular mechanism of conveying regulatory signals in mRNA into a remote polypeptide chain is unlikely, if not impossible. On the other hand, nascent polypeptide folds co-translationally while it is being synthesized. Many reports suggested that folding of many proteins was at least partially co-translational (Hardesty et al., 1999; Pechmann et al., 2013; Rodnina & Wintermeyer, 2016). More importantly, mRNA, ribosome and nascent polypeptide form a complex, allowing the transmission and/or realization of regulatory signals in mRNA for protein folding (Pechmann et al., 2013). In fact, current models linking mRNA structure to protein folding are all based upon co-translational folding pathway. Here the discussion of post-translational folding is skipped and only co-translational folding highlighted.

CONCEPTS OF CO-TRANSLATIONAL PROTEIN FOLDING PROCESS AND ITS REGULATION

Preliminary evidence of co-translational folding for at least some proteins appeared around the same time as Anfinsen performed his seminal experiments on ribonuclease (Cowie et al., 1961; Kiho & Rich, 1964). In theory, full-length unfolded polypeptide was energetically unfavorable (Fedorov & Baldwin, 1997) so that the folding process of most proteins should be more or less co-translational. How co-translational folding proceeds is another complicated point. As for the site of co-translational folding, there are two major steps of co-translational folding. Firstly, newly synthesized polypeptide had to travel through a ribosomal exit tunnel of approximately 80 angstrom in length (Fedyukina & Cavagnero, 2011), which is wide enough to accommodate α -helix formation. Indeed, α -helix within exit tunnel has been directly confirmed by FRET (fluorescence resonance energy transfer) (Woolhead et al., 2004). Additional evidence suggested that exit tunnel could entropically stabilize both α -helix (Ziv et al., 2005) and distinct conformations of nascent polypeptide via extensive contacts with ribosomal components (Bhushan et al., 2010). Secondly, other steps of co-translational folding, especially the higher

order ones impossible within confined space of exit tunnel, occurred after a partial exit of polypeptide from ribosome. For example, co-translational folding of cystic fibrosis transmembrane conductance regulator was dissected experimentally (Kim et al., 2015). And its α -subdomain compaction was delayed until all related polypeptides migrated into cytosol (Kim et al., 2015).

Consistent with our understandings of co-translation protein folding, many *cis* and *trans* regulators have been implicated. Some discovered *trans* regulators include ribosome-bound chaperones capable of operationally extending exit tunnel and providing additional space for protein folding (Kramer et al., 2009). Also co-translational recognition by signal recognition particle (SRP) induced a translocation of nascent peptide into endoplasmic reticulum with a distinct folding environment (Pechmann et al., 2014). As for *cis* regulators, two distinct and yet probably synergistic mechanisms affect co-translational folding (Pechmann et al., 2014). One mechanism operates by recruiting certain *trans* regulators through specific motifs such as SRP-binding elements (Pechmann et al., 2014) while another by modulating elongation speed (O'Brien et al., 2014, 2012). Two mRNA features have been implicated in modulating elongation speed and thus regulating co-translational protein folding, without altering peptide sequences, i.e. codon optimality (Pechmann et al., 2014; Zhang et al., 2009) and mRNA secondary structures (Faure et al., 2016; Jia et al., 2004; Liu & Liu, 1999; Zhang et al., 1998).

IMPACT OF CODON OPTIMALITY ON PROTEIN FOLDING

It was known that 18/20 amino acids are encoded by two or more synonymous codons. Among them, some are called "optimal" because of either their higher thermodynamic stability after pairing with anticodon or a higher abundance of their cognate tRNAs. The current mechanistic model of translational elongation dictates that codon optimality influences translational efficiency and/or accuracy (Gingold & Pilpel, 2011), i.e., optimal codons are translated faster (see below) and/or with higher fidelity. Unlike the sparse data for mRNA structure, accumulation of sequenced ORFs facilitated codon optimality profiling in a wide array of genes and species since 1970s, resulting in a large body of work investigating codon optimality-dependent modulation of translational elongation rate and its effect on co-translational protein folding. For example, *Escherichia coli* multidomain protein SufI was examined. Severe perturbation was reported for SufI folding efficiency by excessive tRNA *in vitro* or synonymous substitution into some clusters of non-optimal codons. It was assumed that the clusters of non-optimal codons transiently attenuated translational elongation, temporally separated the translation of segments of peptide chain and actively coordinated co-translational folding. Considering tRNA supply and demand, Pechmann and colleagues (Pechmann & Frydman, 2013) modeled efficiency of translational elongation in 10 closely related yeast species, and found evolutionarily conserved distribution of codon optimality that is associated with secondary structure of translated polypeptides. The authors suggested that mRNA sequences,

and in particular synonymous codon choices, are generally under selection to optimize the co-translational folding of corresponding polypeptides. Altogether, these and other reports (Komar, 2009) have hinted at an evolutionarily conserved link between clusters of non-optimal codons and pauses of translational elongation that facilitates co-translational protein folding, and more importantly, the existence of additional genetic information in an ORF beyond primary protein sequence. Nevertheless, the exact molecular mechanism for such regulatory effect remains elusive, as cluster of non-optimal codons could have been evolved due to selection for sequence features other than the non-optimality of codons.

MESSANGER RNA STRUCTURE AFFECTS TRANSLATIONAL ELONGATION

As a critical component of genetic information flow for a certain protein coding gene, messenger RNA extracts coding sequences from genome and applies it as a template for protein synthesis. Nevertheless, not merely a sequence of codons, mRNA has its own complex structures. In particular, its secondary structure of Watson-Crick pairing between nucleotides could regulate translational processes at multiple levels. For example, stable secondary structure at the 5' end of mRNA might suppress translational initiation and thus enhance overall translation efficiency in *E. coli* (Kudla et al., 2009). Excessive stable stem regions at the 3' UTR decreased the accessibility of miRNA response elements and interfered with miRNA-mediated translational repression (Kertesz et al., 2007). While the above examples were limited to either end of mRNA, other functional roles of mRNA secondary structure have been discovered for major coding sequences, such as regulating RNA editing (Nishikura, 2006) and splicing (Shepard & Hertel, 2008). Furthermore, secondary structure of nascent mRNA might lower local mutation rate (Chen et al., 2016). More importantly, specific mRNA structure could interfere with the movement of translating ribosome (Brierley et al., 1991; Chen et al., 2013; Qu et al., 2011; Wen et al., 2008). Given numerous reports (Ciryam et al., 2013; O'Brien et al., 2014, 2012; Wang et al., 2015) connecting translational elongation rate to co-translational folding, it is thus not surprising that theoretical studies have already linked mRNA secondary structure to protein folding.

EARLY EVIDENCE FOR THE REGULATORY ROLE OF mRNA STRUCTURE ON PROTEIN FOLDING

The earliest report correlating mRNA structure with the regulation of protein folding appeared in 1993 (Guisez et al., 1993). In the study of Guisez et al., several nascent polypeptide intermediates of coat protein of RNA bacteriophage MS2 were analyzed. And the sizes of nascent polypeptide intermediates were found corresponded to either the positions of rare codons or RNA regions with double-stranded secondary structures, both presumably decrease the velocity of translating ribosomes. It was thus hypothesized that discontinuous translational elongation rate generally facilitates optimal folding of polypeptides. The hypothesized regulated protein folding by

mRNA structure was later supported by two additional empirical analyses. On the one hand, the codons of hydrophobic and hydrophilic amino acids tend to respectively located in stem and loop regions of mRNA (Zhang et al., 1998). Given the crucial role of hydrophobic effect on stabilizing protein structure, such observation is suggestive for the information transfer between mRNA and protein structure (Zhang et al., 1998). On the other hand, experimentally determined protein secondary structures were directly compared with computationally predicted mRNA secondary structures (Jia et al., 2004). And α -helices and β -strands within a folded protein tend to be encoded by double-stranded mRNA regions whereas random coils within polypeptide were more likely to be encoded by single-stranded mRNA regions. Although these studies offered preliminary evidence for an intriguing link between mRNA and protein structure, their limitations were also obvious. On the one hand, due to the scarcity of experimentally determined mRNA structure, these studies resorted to computationally predicted mRNA secondary structure with a modest accuracy at best (Lange et al., 2012), let alone higher level structures. On the other hand, the proposed link between mRNA and protein structure was mediated by the capability of mRNA structure in modulating translational elongation speed, whose exact nature was mostly unknown by then. Recent genomic advances have enabled assessments of the above link.

NOVEL GENOMIC DATA QUESTION THE REGULATION OF PROTEIN FOLDING BY CODON OPTIMALITY

The advances of high throughput sequencing techniques have allowed experimental explorations for both translational elongation speed and mRNA structure at the genomic levels (Graveley, 2016; Ingolia et al., 2012). And the resulting datasets have triggered empirical tests of two major hypotheses for *cis*-regulatory signal in mRNA for co-translational folding, i.e. codon optimality and mRNA structure (Qian et al., 2012; Tuller et al., 2011; Yang et al., 2014). Overall, the effect of codon optimality failed to receive consistent supports. The rationales for the regulatory role of mRNA structure in protein co-translational folding and its relationship with codon optimality shall be summarized in the next section.

Essentially as a snapshot for the distribution of ribosomes within transcriptome, ribosome profiling (Ingolia et al., 2009) utilized high-throughput sequencing of segmental mRNA shielded by translating ribosomes from endonuclease digestion. Since translating ribosomes spend more time on stretches of nucleotides with higher coverage in ribosome profiling than other nucleotides in the same gene, such detail ribosomal kinetics allowed revelation/confirmation of several critical features of translational elongation. Firstly, translational elongation rate was not uniform among different mRNAs or along a single mRNA molecule (Ingolia et al., 2009, 2011). Secondly, strong ribosomal pauses lasting over a couple of seconds, >10 times slower than average elongation speed, were widely distributed (Ingolia et al., 2011). Thirdly, at least some variations of elongation speed within gene was obviously non-neutral and had evolved under natural selections (Tuller et

al., 2010). All the above findings were consistent with the model of co-translational protein folding as regulated by translational elongation rate, necessitating the validation of codon optimality and/or mRNA structure as a regulator of elongation rate.

Indeed existing data of ribosome profiling have enabled independent assessments of the role of codon optimality in the control of translational elongation speed. Unexpectedly, several attempts of confirming the slow translational speed of individual non-optimal codons failed to reveal any signal in genomic ribosome profiling data of several species, including yeast (Charneski & Hurst, 2013; Qian et al., 2012), worms (Stadler & Fire, 2011), rodents (Ingolia et al., 2011) and bacteria (Li et al., 2012). These studies explicitly tested the correlation between codon optimality and elongation speed, and found negative results so that other determinants of elongation speed were examined, such as positively-charged amino acids (Charneski & Hurst, 2013), wobble base-pairing (Stadler & Fire, 2011) and anti-Shine-Dalgarno sequence (Li et al., 2012). In one of these studies, observations were explained by balanced synonymous codon usage of transcriptome relative to the abundance of tRNA (Qian et al., 2012). When there was an overall shortage of translation-ready tRNAs, balanced codon usage makes tRNA shortage for all codons similar. It avoided long ribosomal pauses caused by extreme tRNA shortage for a few codons and thus minimizing transcriptome-wide total duration of ribosomal pauses. Further analyses confirmed such a balanced codon usage for multiple eukaryotic transcriptomes. It hinted at adaptive evolution towards balanced codon usage, which presumably provides optimal allocation of translational resources and alleviated ribosomal sequestering due to translational pauses. As experimentally validated, global translational efficiency increased after a heterologous gene with balanced codon usage was transfected into yeast cells, compared to a gene using only optimal or non-optimal codons. The experimental observations were consistent with codon harmonization (Angov et al., 2008), a strategy commonly employed to enhance heterologous protein expression in synthetic biology. More importantly, the model of balanced codon usage indicates that previous experimental results correlating non-optimal codons with halt of translational elongation could be artefactual since most of them involved transfecting a highly expressed heterologous gene into a host cell. A high expression of heterologous gene perturbed the balance between codon usage and tRNA supply. Since the absolute number of cognate tRNA for optimal codon was higher, tRNA shortage was thus proportionally less serious for optimal than non-optimal codons so that there was faster translational elongation for optimal codons in heterologous system. Collectively, the above results cast doubts over the conventional wisdom of faster translation of optimal codons (Charneski & Hurst, 2013; Ingolia et al., 2011; Qian et al., 2012; Yang et al., 2014).

Nevertheless, the results suggesting no correlation between codon optimality and ribosomal velocity are not without their own problems. Most notably, when cycloheximide was used for stabilizing ribosomes prior to position measurements, elongation re-occurred in the presence of cycloheximide but

with dramatically altered codon-specific elongation rates. And the measured positions of ribosomes failed to reflect the temporal durations of ribosomal pausing at each position *in vivo* (Hussmann et al., 2015). Meanwhile, other studies have independently examined the correlation between codon optimality and translational elongation rate, but inconsistent results were obtained (Gardin et al., 2014; Li, et al., 2012; Stadler & Fire, 2011). After analyzing multiple datasets of ribosome profiling, it was found that, regardless of using cycloheximide or not prior to cell lysis, the reproducibility of ribosome profiling was poor at codon resolution since signals at this level were not well-reproduced in experimental replicates (Diament & Tuller, 2016). Previous theoretical and experimental results have confirmed the regulation of co-translational protein folding by clusters of non-optimal codons through modulation of elongation speed. However, the exact molecular mechanism for non-optimal codon cluster stalling ribosomal movement has remained elusive. It left the possibilities of alternative or synergistic regulations other than codon optimality, such that the clusters of non-optimal codons are probably byproducts of other sequence features.

NOVEL GENOMIC DATA SUPPORTING THE REGULATION OF PROTEIN FOLDING BY mRNA STRUCTURE

The development of high-throughput sequencing has enabled multiple methods for examining RNA secondary structure at a genomic level. Early attempts of FragSeq (Underwood et al., 2010), PARS (Kertesz et al., 2010) and SHAPE-seq (Lucks et al., 2011) utilized P1 nuclease, RNase V1 & S1 nuclease and 1-methyl-7-nitroisatoic anhydride respectively, to probe the structures of a large pool of synthetic RNAs or total RNA after extraction from cells, which revealed *in vitro* pairing status of individual nucleotides on RNA molecules. These approaches were followed by the development of mod-seq (Talkish et al., 2014), DMS-seq (Rouskin et al., 2014), Structure-seq (Ding et al., 2014), icSHAPE (Spitale et al., 2015) and SHAPE-Map (Smola et al., 2015), which were capable of detecting *in vivo* RNA secondary structure. More recently, techniques have been developed for detecting pairing partners, including RPL (Ramani et al., 2015), PARIS (Lu et al., 2016), SPLASH (Aw et al., 2016) and LIGR-seq (Sharma et al., 2016). These advanced techniques have cleared the obstacles of genomic RNA structure investigation and enhanced the profiling accuracy of RNA secondary structure (Lange et al., 2012).

Combined with ribosomal profiling data, genomic profiles of RNA secondary structure supported the role of mRNA structure in modulating elongation speed (Tuller et al., 2010, 2011; Yang et al., 2014) (but see Charneski & Hurst, 2013). In particular, comparisons among genes revealed stronger mRNA secondary structure (Zur & Tuller, 2012) and slower translational elongation (Yang et al., 2014) for highly expressed genes, which is more sensitive to protein misfolding (Yang et al., 2010; Zhang & Yang, 2015). Additional comparisons within gene revealed that mRNA pairing status at the entrance of ribosome had the strongest impact upon elongation rate (Yang et al., 2014). This result was consistent with single molecule level

study of individual translating ribosome using optical tweezers, who found that pausing duration of ribosomal translocation was significantly dependent on mRNA secondary structure (Wen et al., 2008). More importantly, the regulatory effect of mRNA secondary structure on ribosome velocity seemed independent of codon optimality (Yang et al., 2014). Thus mRNA secondary structure might serve as a regulator of protein co-translational folding via modulating elongation speed. During translation, mRNA secondary structures were actively unfolded by ribosomes (Rouskin et al., 2014). However, the distance between adjacent ribosomes was approximately 20-35 nm in eukaryotes. And it was translated into 50-90 nt or 17-30 codons, allowing enough time for intervening mRNA to refold given the thousand fold difference between timescale of ribosomal elongation (~ 0.1 s per codon (Ingolia et al., 2009, 2011) and RNA folding kinetics (10^{-5} s for folding a simple hairpin (de Smit & van Duin, 2003)).

More recently, new genomic data were used for directly testing the connection between mRNA structure and protein folding by comparison among genes (Faure et al., 2016). Using protein structures from 2 eukaryotes and 3 prokaryotes, protein compactness was positively correlated with the stability of mRNA structure. Such correlations are more pronounced in ordered parts than disordered parts of protein. Thus it suggested an important role of mRNA secondary structure in modulating protein folding. More importantly, comparison with translational efficiency inferred from ribosome profiling data supported that stable mRNAs were translated slowly to allow more time for compact proteins to fold co-translationally (Faure et al., 2016).

Collectively, new genomic data of ribosome profiling and mRNA secondary structure suggested a mechanistic model (Fig. 1), where the co-translational protein folding is regulated by mRNA secondary structure through its modulation of translational elongation rate. Such a regulation was independent of elongation slowdown due to nonoptimality of synonymous codon usage, whose capacity of regulating co-translational protein folding has remained debated.

BIOMEDICAL IMPLICATIONS

In conclusion, elucidating the regulatory role of mRNA secondary structure for protein folding shall expand our understandings of the full contents of genetic information and the molecular mechanisms of its phenotypic expression. Gaining such insights offers broad implications for biological researches. For example, combined with proper bioinformatic algorithm for designing mRNA structure, it is bound to enhance our capability of expressing functionally heterologous proteins in cells. The generality of this regulatory role has raised questions on the neutrality of synonymous variations in coding sequences for molecular evolutionary analyses. The regulatory role of mRNA structure in protein folding has become the only model capable of explaining the stronger mRNA secondary structures in highly expressed genes (Yang et al., 2014; Yang & Zhang, 2015). It offers a fundamental tool of understanding how natural selection concert the optimality of synonymous codon




Figure 1 A mechanistic model of co-translational protein folding under the regulation of mRNA secondary structure

Accuracy of co-translational folding is affected by translational elongation speed so that optimal elongation speed varies for different regions within coding sequences. For regions where slow elongation is optimal, mRNA forms stable secondary structure to slow down the movement of ribosomes (highlighted by yellow box). On the contrary, single-stranded mRNA enables faster elongation in regions where higher elongation rate is preferred for accurate co-translational folding (highlighted by pink box).

usage and mRNA secondary structure and subsequently affects the evolution of coding sequences. Co-translational misfolding is a form of phenotypic mutation. The regulatory role of mRNA structure in both protein folding and mutation rate may lead to a quantitative coupling between genotypic and phenotypic mutation rates (Chen et al., 2016). Full biological ramifications of such intriguing coupling between processing and transmission fidelity of genetic information await further explorations.

Detailed modeling of co-translational folding modulated by mRNA structure can help us predict or interpret the phenotypic effects and elucidate the underlying mechanisms of synonymous variations ubiquitous in human genome (The 1000 Genomes Project Consortium, 2010). Implicated in human diseases (Kimchi-Sarfaty et al., 2007), it is considered as frequent driver mutations in human cancer (Supek et al., 2014). Altered mRNA structure might result in a dysregulation of co-translational protein folding, leading to protein misfolding and aggregation that is disproportionately involved in neurodegenerative diseases (Soto, 2003). Understanding the

subtle roles of mRNA structure in protein misfolding and aggregation shall reveal new therapeutic targets for neurodegenerative diseases.

ACKNOWLEDGEMENTS

Great appreciations were extended to Yong-Gang Yao, Gong-Wang Yu, Heng Zhang and two anonymous reviewers for valuable comments.

REFERENCES

- Angov E, Hillier CJ, Kincaid RL, Lyon JA. 2008. Heterologous protein expression is enhanced by harmonizing the codon usage frequencies of the target gene with those of the expression host. *PLoS One*, **3**(5): e2189, doi: 10.1371/journal.pone.0002189.
- Aw JGA, Shen Y, Wilm A, Sun M, Lim XN, Boon KL, Tapsin S, Chan YS, Tan CP, Sim AYL, Zhang T, Susanto TT, Fu ZY, Nagarajan N, Wan Y. 2016. In vivo mapping of eukaryotic RNA interactomes reveals principles of higher-order organization and regulation. *Molecular Cell*, **62**(4): 603-617, doi: 10.1016/j.molcel.2016.04.028.
- Bhushan S, Gartmann M, Halic M, Armache JP, Jarasch A, Mielke T, Berninghausen O, Wilson DN, Beckmann R. 2010. α -Helical nascent polypeptide chains visualized within distinct regions of the ribosomal exit tunnel. *Nature Structural & Molecular Biology*, **17**(3): 313-317, doi: 10.1038/nsmb.1756.
- Brierley I, Rolley NJ, Jenner AJ, Inglis SC. 1991. Mutational analysis of the RNA pseudoknot component of a coronavirus ribosomal frameshifting signal. *Journal of Molecular Biology*, **220**(4): 889-902.
- Charneski CA, Hurst LD. 2013. Positively charged residues are the major determinants of ribosomal velocity. *PLoS Biology*, **11**(3): e1001508, doi: 10.1371/journal.pbio.1001508.
- Chen CL, Zhang HB, Broitman SL, Reiche M, Farrell I, Cooperman BS, Goldman YE. 2013. Dynamics of translation by single ribosomes through mRNA secondary structures. *Nature Structural & Molecular Biology*, **20**(5): 582-588, doi: 10.1038/nsmb.2544.
- Chen X, Yang JR, Zhang J. 2016. Nascent RNA folding mitigates transcription-associated mutagenesis. *Genome Research*, **26**(1): 50-59, doi: 10.1101/gr.195164.115.
- Ciryam P, Morimoto RI, Vendruscolo M, Dobson CM, O'Brien EP. 2013. In vivo translation rates can substantially delay the cotranslational folding of the *Escherichia coli* cytosolic proteome. *Proceedings of the National Academy of Sciences of the United States of America*, **110**(2): E132-E140, doi: 10.1073/pnas.1213624110.
- Cowie DB, Spiegelman S, Roberts RB, Duerksen JD. 1961. Ribosome-bound beta-galactosidase. *Proceedings of the National Academy of Sciences of the United States of America*, **47**: 114-122.
- de Smit MH, van Duin J. 2003. Translational standby sites: how ribosomes may deal with the rapid folding kinetics of mRNA. *Journal of Molecular Biology*, **331**(4): 737-743.
- Diament A, Tuller T. 2016. Estimation of ribosome profiling performance and reproducibility at various levels of resolution. *Biology Direct*, **11**: 24, doi: 10.1186/s13062-016-0127-4.
- Ding YL, Tang Y, Kwok CK, Zhang Y, Bevilacqua PC, Assmann SM. 2014. In vivo genome-wide profiling of RNA secondary structure reveals novel regulatory features. *Nature*, **505**(7485): 696-700, doi: 10.1038/nature12756.
- Eddy SR. 2014. Computational analysis of conserved RNA secondary structure in transcriptomes and genomes. *Annual Review of Biophysics*, **43**: 433-456, doi: 10.1146/annurev-biophys-051013-022950.
- Faure G, Ogurtsov AY, Shabalina SA, Koonin EV. 2016. Role of mRNA structure in the control of protein folding. *Nucleic Acids Research*, **44**(22): 10898-10911, doi: 10.1093/nar/gkw671.
- Fedorov AN, Baldwin TO. 1997. Cotranslational protein folding. *Journal of Biological Chemistry*, **272**(52): 32715-32718.
- Fedyukina DV, Cavagnero S. 2011. Protein folding at the exit tunnel. *Annual Review of Biophysics*, **40**: 337-359, doi: 10.1146/annurev-biophys-042910-155338.
- Fraser PE. 2014. Prions and prion-like proteins. *Journal of Biological Chemistry*, **289**(29): 19839-19840, doi: 10.1074/jbc.R114.583492.
- Gardin J, Yeasmin R, Yurovsky A, Cai Y, Skiena S, Futcher B. 2014. Measurement of average decoding rates of the 61 sense codons in vivo. *eLife*, **3**, doi: 10.7554/eLife.03735.
- Gingold H, Pilpel Y. 2011. Determinants of translation efficiency and accuracy. *Molecular Systems Biology*, **7**: 481, doi: 10.1038/msb.2011.14.
- Graveley BR. 2016. RNA matchmaking: finding cellular pairing partners. *Molecular Cell*, **63**(2): 186-189, doi: 10.1016/j.molcel.2016.07.001.
- Guisez Y, Robbens J, Remaut E, Fiers W. 1993. Folding of the MS2 coat protein in *Escherichia coli* is modulated by translational pauses resulting from mRNA secondary structure and codon usage: a hypothesis. *Journal of Theoretical Biology*, **162**(2): 243-252, doi: 10.1006/jtbi.1993.1085.
- Hardesty B, Tsalkova T, Kramer G. 1999. Co-translational folding. *Current Opinion in Structural Biology*, **9**(1): 111-114.
- Husmann JA, Patchett S, Johnson A, Sawyer S, Press WH. 2015. Understanding biases in ribosome profiling experiments reveals signatures of translation dynamics in yeast. *PLoS Genetics*, **11**(12): e1005732, doi: 10.1371/journal.pgen.1005732.
- Ingolia NT, Ghaemmaghami S, Newman JR, Weissman JS. 2009. Genome-wide analysis in vivo of translation with nucleotide resolution using ribosome profiling. *Science*, **324**(5924): 218-223, doi: 10.1126/science.1168978.
- Ingolia NT, Lareau LF, Weissman JS. 2011. Ribosome profiling of mouse embryonic stem cells reveals the complexity and dynamics of mammalian proteomes. *Cell*, **147**(4): 789-802, doi: 10.1016/j.cell.2011.10.002.
- Ingolia NT, Brar GA, Rouskin S, McGeachy AM, Weissman JS. 2012. The ribosome profiling strategy for monitoring translation in vivo by deep sequencing of ribosome-protected mRNA fragments. *Nature Protocols*, **7**(8): 1534-1550, doi: 10.1038/nprot.2012.086.
- Jia MW, Luo LF, Liu CQ. 2004. Statistical correlation between protein secondary structure and messenger RNA stem-loop structure. *Biopolymers*, **73**(1): 16-26, doi: 10.1002/bip.10496.
- Kertesz M, Iovino N, Unnerstall U, Gaul U, Segal E. 2007. The role of site accessibility in microRNA target recognition. *Nature Genetics*, **39**(10): 1278-1284, doi: 10.1038/ng2135.
- Kertesz M, Wan Y, Mazar E, Rinn JL, Nutter RC, Chang HY, Segal E. 2010. Genome-wide measurement of RNA secondary structure in yeast. *Nature*, **467**(7311): 103-107, doi: 10.1038/nature09322.
- Kiho Y, Rich A. 1964. Induced enzyme formed on bacterial polyribosomes. *Proceedings of the National Academy of Sciences of the United States of*

America, **51**(1): 111-118.

Kim SJ, Yoon JS, Shishido H, Yang Z, Rooney LA, Barral JM, Skach WR. 2015. Protein folding. Translational tuning optimizes nascent protein folding in cells. *Science*, **348**(6233): 444-448, doi: 10.1126/science.aaa3974.

Kimchi-Sarfaty C, Oh JM, Kim IW, Sauna ZE, Calcagno AM, Ambudkar SV, Gottesman MM. 2007. A "silent" polymorphism in the MDR1 gene changes substrate specificity. *Science*, **315**(5811): 525-528, doi: 10.1126/science.1135308.

Kolata G. 1986. Trying to crack the second half of the genetic code. *Science*, **233**(4768): 1037-1039.

Komar AA. 2009. A pause for thought along the co-translational folding pathway. *Trends in Biochemical Sciences*, **34**(1): 16-24, doi: 10.1016/j.tibs.2008.10.002.

Kramer G, Boehringer D, Ban N, Bukau B. 2009. The ribosome as a platform for co-translational processing, folding and targeting of newly synthesized proteins. *Nature Structural & Molecular Biology*, **16**(6): 589-597, doi: 10.1038/nsmb.1614.

Kudla G, Murray AW, Tollervey D, Plotkin JB. 2009. Coding-sequence determinants of gene expression in *Escherichia coli*. *Science*, **324**(5924): 255-258, doi: 10.1126/science.1170160.

Lange SJ, Maticzka D, Möhl M, Gagnon JN, Brown CM, Backofen R. 2012. Global or local? Predicting secondary structure and accessibility in mRNAs. *Nucleic Acids Research*, **40**(12): 5215-5226, doi: 10.1093/nar/gks181.

Li GW, Oh E, Weissman JS. 2012. The anti-Shine-Dalgarno sequence drives translational pausing and codon choice in bacteria. *Nature*, **484**(7395): 538-541, doi: 10.1038/nature10965.

Liu SQ, Liu CQ. 1999. The relationship between mRNA'S sequence, structure, translational rate and protein structure. *Zoological Research*, **20**(6): 457-461. (in Chinese)

Lu ZP, Zhang QC, Lee B, Flynn RA, Smith MA, Robinson JT, Davidovich C, Gooding AR, Goodrich KJ, Mattick JS, Mesirov JP, Cech TR, Chang HY. 2016. RNA duplex map in living cells reveals higher-order transcriptome structure. *Cell*, **165**(5): 1267-1279, doi: 10.1016/j.cell.2016.04.028.

Lucks JB, Mortimer SA, Trapnell C, Luo SJ, Aviran S, Schroth GP, Pachter L, Doudna JA, Arkin AP. 2011. Multiplexed RNA structure characterization with selective 2'-hydroxyl acylation analyzed by primer extension sequencing (SHAPE-Seq). *Proceedings of the National Academy of Sciences of the United States of America*, **108**(27): 11063-11068, doi: 10.1073/pnas.1106501108.

Nishikura K. 2006. Editor meets silencer: crosstalk between RNA editing and RNA interference. *Nature Reviews Molecular Cell Biology*, **7**(12): 919-931, doi: 10.1038/nrm2061.

O'Brien EP, Vendruscolo M, Dobson CM. 2012. Prediction of variable translation rate effects on cotranslational protein folding. *Nature Communications*, **3**: 868, doi: 10.1038/ncomms1850.

O'Brien EP, Vendruscolo M, Dobson CM. 2014. Kinetic modelling indicates that fast-translating codons can coordinate cotranslational protein folding by avoiding misfolded intermediates. *Nature Communications*, **5**: 2988, doi: 10.1038/ncomms3988.

Pechmann S, Frydman J. 2013. Evolutionary conservation of codon optimality reveals hidden signatures of cotranslational folding. *Nature Structural & Molecular Biology*, **20**(2): 237-243, doi: 10.1038/nsmb.2466.

Pechmann S, Willmund F, Frydman J. 2013. The ribosome as a hub for protein quality control. *Molecular Cell*, **49**(3): 411-421, doi: 10.1016/j.molcel.2013.01.020.

Pechmann S, Chartron JW, Frydman J. 2014. Local slowdown of translation by nonoptimal codons promotes nascent-chain recognition by SRP *in vivo*. *Nature Structural & Molecular Biology*, **21**(12): 1100-1105, doi: 10.1038/nsmb.2919.

Qian WF, Yang JR, Pearson NM, Maclean C, Zhang JZ. 2012. Balanced codon usage optimizes eukaryotic translational efficiency. *PLoS Genetics*, **8**(3): e1002603, doi: 10.1371/journal.pgen.1002603.

Qu XH, Wen JD, Lancaster L, Noller HF, Bustamante C, Tinoco I, Jr. 2011. The ribosome uses two active mechanisms to unwind messenger RNA during translation. *Nature*, **475**(7354): 118-121, doi: 10.1038/nature10126.

Ramani V, Qiu RL, Shendure J. 2015. High-throughput determination of RNA structure by proximity ligation. *Nature Biotechnology*, **33**(9): 980-984, doi: 10.1038/nbt.3289.

Ramos SBV, Laederach A. 2014. Molecular biology: A second layer of information in RNA. *Nature*, **505**(7485): 621-622, doi: 10.1038/505621a.

Rodnina MV, Wintermeyer W. 2016. Protein elongation, Co-translational folding and targeting. *Journal of Molecular Biology*, **428**: 2165-2185, doi: 10.1016/j.jmb.2016.03.022.

Rouskin S, Zubradt M, Washietl S, Kellis M, Weissman JS. 2014. Genome-wide probing of RNA structure reveals active unfolding of mRNA structures *in vivo*. *Nature*, **505**(7485): 701-705, doi: 10.1038/nature12894.

Sharma E, Sterne-Weiler T, O'Hanlon D, Blencowe BJ. 2016. Global mapping of human RNA-RNA interactions. *Molecular Cell*, **62**(4): 618-626, doi: 10.1016/j.molcel.2016.04.030.

Shepard PJ, Hertel KJ. 2008. Conserved RNA secondary structures promote alternative splicing. *RNA*, **14**(8): 1463-1469, doi: 10.1261/ma.1069408.

Smola MJ, Rice GM, Busan S, Siegfried NA, Weeks KM. 2015. Selective 2'-hydroxyl acylation analyzed by primer extension and mutational profiling (SHAPE-MaP) for direct, versatile and accurate RNA structure analysis. *Nature Protocols*, **10**: 1643-1669, doi: 10.1038/nprot.2015.103.

Soto C. 2003. Unfolding the role of protein misfolding in neurodegenerative diseases. *Nature Reviews Neuroscience*, **4**(1): 49-60, doi: 10.1038/nrn1007.

Spencer PS, Barral JM. 2012. Genetic code redundancy and its influence on the encoded polypeptides. *Computational and Structural Biotechnology Journal*, **1**(1): e201204006, doi: 10.5936/csbj.201204006.

Spitale RC, Flynn RA, Zhang QC, Crisalli P, Lee B, Jung JW, Kuchelmeister HY, Batista PJ, Torre EA, Kool ET, Chang HY. 2015. Structural imprints *in vivo* decode RNA regulatory mechanisms. *Nature*, **519**(7544): 486-490, doi: 10.1038/nature14263.

Stadler M, Fire A. 2011. Wobble base-pairing slows *in vivo* translation elongation in metazoans. *RNA*, **17**(12): 2063-2073, doi: 10.1261/rna.02890211.

Supek F, Miñana B, Valcárcel J, Gabaldón T, Lehner B. 2014. Synonymous mutations frequently act as driver mutations in human cancers. *Cell*, **156**(6): 1324-1335, doi: 10.1016/j.cell.2014.01.051.

Talkish J, May G, Lin YZ, Woolford JL Jr., McManus CJ. 2014. Mod-seq: high-throughput sequencing for chemical probing of RNA structure. *RNA*, **20**(5): 713-720, doi: 10.1261/rna.042218.113.

The 1000 Genomes Project Consortium. 2010. A map of human genome variation from population-scale sequencing. *Nature*, **467**(7319): 1061-1073, doi: 10.1038/nature09534.

The ENCODE Project Consortium. 2012. An integrated encyclopedia of DNA elements in the human genome. *Nature*, **489**(7414): 57-74, doi: 10.1038/nature11247.

- Tsai CJ, Sauna ZE, Kimchi-Sarfaty C, Ambudkar SV, Gottesman MM, Nussinov R. 2008. Synonymous mutations and ribosome stalling can lead to altered folding pathways and distinct minima. *Journal of Molecular Biology*, **383**(2): 281-291, doi: 10.1016/j.jmb.2008.08.012.
- Tuller T, Carmi A, Vestsigian K, Navon S, Dorfan Y, Zaborse J, Pan T, Dahan O, Furman I, Pilpel Y. 2010. An evolutionarily conserved mechanism for controlling the efficiency of protein translation. *Cell*, **141**(2): 344-354, doi: 10.1016/j.cell.2010.03.031.
- Tuller T, Veksler-Lublinsky I, Gazit N, Kupiec M, Ruppin E, Ziv-Ukelson M. 2011. Composite effects of gene determinants on the translation speed and density of ribosomes. *Genome Biology*, **12**(11): R110, doi: 10.1186/gb-2011-12-11-r110.
- Underwood JG, Uzilov AV, Katzman S, Onodera CS, Mainzer JE, Mathews DH, Lowe TM, Salama SR, Haussler D. 2010. FragSeq: transcriptome-wide RNA structure probing using high-throughput sequencing. *Nature Methods*, **7**(12): 995-1001, doi: 10.1038/nmeth.1529.
- Wang EC, Wang J, Chen CJ, Xiao Y. 2015. Computational evidence that fast translation speed can increase the probability of cotranslational protein folding. *Scientific Reports*, **5**: 15316, doi: 10.1038/srep15316.
- Wen JD, Lancaster L, Hodges C, Zeri AC, Yoshimura SH, Noller HF, Bustamante C, Tinoco I. 2008. Following translation by single ribosomes one codon at a time. *Nature*, **452**(7187): 598-603, doi: 10.1038/nature06716.
- Woolhead CA, McCormick PJ, Johnson AE. 2004. Nascent membrane and secretory proteins differ in FRET-detected folding far inside the ribosome and in their exposure to ribosomal proteins. *Cell*, **116**(5): 725-736.
- Xia K, Manning M, Hesham H, Lin QS, Bystroff C, Colón W. 2007. Identifying the subproteome of kinetically stable proteins via diagonal 2D SDS/PAGE. *Proceedings of the National Academy of Sciences of the United States of America*, **104**(44): 17329-17334, doi: 10.1073/pnas.0705417104.
- Yang JR, Zhuang SM, Zhang JZ. 2010. Impact of translational error-induced and error-free misfolding on the rate of protein evolution. *Molecular Systems Biology*, **6**(1): 421, doi: 10.1038/msb.2010.78.
- Yang JR, Chen XS, Zhang JZ. 2014. Codon-by-codon modulation of translational speed and accuracy via mRNA folding. *PLoS Biology*, **12**(7): e1001910, doi: 10.1371/journal.pbio.1001910.
- Yang JR, Zhang JZ. 2015. Human long noncoding RNAs are substantially less folded than messenger RNAs. *Molecular Biology and Evolution*, **32**(4): 970-977, doi: 10.1093/molbev/msu402.
- Zhang G, Hubalewska M, Ignatova Z. 2009. Transient ribosomal attenuation coordinates protein synthesis and co-translational folding. *Nature Structural & Molecular Biology*, **16**(3): 274-280, doi: 10.1038/nsmb.1554.
- Zhang J, Gu BH, Peng SL, Liu CQ. 1998. Distributions of triplet codons in messenger RNA secondary structures. *Zoological Research*, **19**(5): 350-358.
- Zhang JZ, Yang JR. 2015. Determinants of the rate of protein sequence evolution. *Nature Reviews Genetics*, **16**(7): 409-420, doi: 10.1038/nrg3950.
- Ziv G, Haran G, Thirumalai D. 2005. Ribosome exit tunnel can entropically stabilize α -helices. *Proceedings of the National Academy of Sciences of the United States of America*, **102**(52): 18956-18961, doi: 10.1073/pnas.0508234102.
- Zur H, Tuller T. 2012. Strong association between mRNA folding strength and protein abundance in *S. cerevisiae*. *EMBO Reports*, **13**(3): 272-277, doi: 10.1038/embor.2011.262.
- Zwanzig R, Szabo A, Bagchi B. 1992. Levinthal's paradox. *Proceedings of the National Academy of Sciences of the United States of America*, **89**(1): 20-22.

Expression of plgR in the tracheal mucosa of SHIV/SIV-infected rhesus macaques

Dong Li, Feng-Jie Wang, Lei Yu, Wen-Rong Yao, Yan-Fang Cui, Gui-Bo Yang*

National Institute of AIDS/STD Control and Prevention, China-CDC, Beijing 102206, China

ABSTRACT

Polymeric immunoglobulin receptors (plgR) are key participants in the formation and secretion of secretory IgA (S-IgA), which is critical for the prevention of microbial infection and colonization in the respiratory system. Although increased respiratory colonization and infections are common in HIV/AIDS, little is known about the expression of plgR in the airway mucosa of these patients. To address this, the expression levels of plgR in the tracheal mucosa and lungs of SHIV/SIV-infected rhesus macaques were examined by real-time RT-PCR and confocal microscopy. We found that the levels of both *PIGR* mRNA and plgR immunoreactivity were lower in the tracheal mucosa of SHIV/SIV-infected rhesus macaques than that in non-infected rhesus macaques, and the difference in plgR immunoreactivity was statistically significant. IL-17A, which enhances plgR expression, was also changed in the same direction as that of plgR. In contrast to changes in the tracheal mucosa, plgR and IL-17A levels were higher in the lungs of infected rhesus macaques. These results indicated abnormal plgR expression in SHIV/SIV, and by extension HIV infections, which might partially result from IL-17A alterations and might contribute to the increased microbial colonization and infection related to pulmonary complications in HIV/AIDS.

Keywords: Tracheal mucosa; Lungs; plgR; SHIV/SIV infection; IL-17A

INTRODUCTION

The respiratory system is continuously exposed to foreign antigens from either airborne or commensal microbes. Due to vulnerability of the physical epithelial barrier of the respiratory system, most pathogens are stopped from entering the body by the mucosal immune system. A key component of the airway mucosal immune system that prevents microbial infections and colonization is secretory IgA (S-IgA), which is composed of

dimeric IgA produced in the lamina propria and extracellular part of the polymeric immunoglobulin receptors (plgR), also known as the secretory component (SC) expressed by mucosal epithelial cells (Johansen & Kaetzel, 2011).

Newly synthesized plgR is localized to the basolateral surfaces of mucosal epithelial cells, where it binds to dimeric IgA (dIgA) and mediates transcytosis of IgA to the apical surface of the epithelial cells (Johansen et al., 1999). The SC can be released to the mucosal surface alone (in the absence of IgA) or together with dIgA as S-IgA. In addition, SC bound to dIgA can elongate the life of S-IgA and enhance its immune exclusion ability. It can also stop microbial invasion. Mice deficient in plgR expression are reportedly unable to control infections of the airway by some bacteria, which could drive progressive chronic obstructive pulmonary disease (COPD) phenotype in these mice (Richmond et al., 2016).

Pulmonary complications are common and major causes of morbidity and mortality in HIV-infected individuals, even in the presence of highly active antiretroviral therapy (ART) (Grubb et al., 2006; Murray, 1996). Increased pulmonary infections and microbial colonization are common in HIV/AIDS patients (Zar, 2008). Whether and how the S-IgA/plgR system is involved in these alterations is not well addressed. Rhesus macaques are important in HIV/AIDS studies. In previous research, we found that plgR expression was altered in the gut mucosa of SHIV/SIV-infected rhesus macaques (Wang & Yang, 2016). To determine whether plgR is involved in the respiratory pathology of HIV/AIDS, we examined the expression of plgR in the tracheal mucosa of SHIV/SIV-infected rhesus macaques.

MATERIALS AND METHODS

Tissues

Tissue samples from the tracheas and lungs were collected from five normal and five SHIV/SIV-infected rhesus macaques

Received: 11 October 2016; Accepted: 10 November 2016

Foundation items: The study was supported by the Beijing Natural Science Foundation (7162136)

*Corresponding author, E-mail: guiboyang@chinaaids.cn

DOI:10.13918/j.issn.2095-8137.2017.007

(*Macaca mulatta*), as reported previously (Wang & Yang, 2016). The sites from which samples were collected were chosen randomly. Tissue samples for RNA isolation were frozen on dry ice immediately after collection and preserved in a freezer at -80 °C before use. Tissue samples for confocal microscopy were fixed in 4% paraformaldehyde immediately after collection, then washed and protected with 30% sucrose, and finally embedded in OCT and preserved in a freezer at -80 °C. All study animals were treated humanely per the state and local regulations on the care and use of experimental animals.

Real-time RT-PCR

Quantification of plgR and IL-17A mRNA levels was conducted by TaqMan® probe real-time RT-PCR, as performed previously (Wang & Yang, 2016; Zhang et al., 2014). Briefly, RNA was isolated using a RNeasy Pure Tissue Kit (Qiagen Biotech, China) per the manufacturer's protocols. Real-time PCR mixtures were established with a One Step PrimeScript® RT-PCR Kit (Takara, Japan) and primers and probes for plgR and IL-17A (Wang & Yang, 2016; Zhang et al., 2014). PCR was performed on a 7500 Real-Time PCR System with 7500 System SDS software version 1.4 (ABI, USA). GAPDH mRNA levels in all samples were used as internal controls.

Confocal microscopy

Tissue sections were cut with a cryostat LEICA CM 1850 (Leica Inc., Germany) to a thickness of 20 microns. After removing the OCT with PBS supplemented with 0.1% Triton X-100 and FSG, the slides were washed with PBST and blocked with 10% normal goat serum for 1 h before incubation in polyclonal antibody against human plgR (rabbit anti-PIGR, 4 µg/mL, Abcam, USA) at 4 °C overnight. Sections were incubated in secondary antibody (Alexa Fluor 488 conjugated goat anti-rabbit IgG, 2 µg/mL, Invitrogen, USA) for 1 h after washing off the extra primary antibody. Slides were washed and mounted with anti-fade mounting medium and observed with an Olympus FV1000D-ST confocal microscope (Olympus, Japan). Images (1024×1024) were acquired and morphometric measurements were obtained with Image-Pro Plus software version 6.0 (Media Cybernetics, Silver Springs, MD, USA).

Statistics

All quantitative parameters were expressed as mean±SD. Non-parametric Mann-Whitney *U* test was used to compare the means of parameters between normal and infected rhesus macaques. Spearman test was used to calculate the correlations between plgR mRNA and IL-17A mRNA levels. *P* values of less than 0.05 were considered statistically significant.

RESULTS

Localization of plgR immunoreactivity in the tracheal mucosa of rhesus macaques

To detect the expression of plgR in the tracheal mucosa of rhesus macaques, plgR immunoreactive cells were examined with confocal microscopy. As shown in Figure 1, plgR immunoreactivity was detected with a polyclonal antibody

against human plgR. In the epithelium, immunoreactivity to plgR was localized to both the apical and basolateral surfaces of the epithelial cells. It was also localized in the cytoplasm of the basal part (under the nucleus) of the epithelial cells. After SHIV/SIV infection, plgR immunoreactivity was lower in the tracheal mucosa of rhesus macaques.

Expression of plgR decreased in the tracheal epithelium of SHIV/SIV-infected rhesus macaques

To determine changes in plgR expression after SHIV/SIV infection, levels of plgR immunoreactivity were quantitatively examined with Image-Pro Plus software and plgR mRNA levels were examined by real-time PCR. As shown in Figure 2, levels of plgR immunoreactivity were 1.65 times higher in the tracheal epithelium of normal rhesus macaques than that in SHIV/SIV-infected rhesus macaques (Figure 2A), with statistical significance (Mann-Whitney *U* test, *P*=0.0079). The transcription levels of plgR genes in the tracheal mucosa of normal rhesus macaques were 1.57 times higher than that in infected rhesus macaques, although the difference was not statistically significant (Mann-Whitney *U* test, *P*=0.2544). Therefore, both the transcription and protein levels of plgR were about 1.6 times higher in normal than in infected rhesus macaques.

IL-17A is a regulator of plgR expression and is decreased in HIV and SIV infection. We examined the transcription levels of IL-17A in the tracheal mucosa of normal and infected rhesus macaques. IL-17A mRNA levels in the tracheal mucosa of normal rhesus macaques were 1.8 times higher than that in SHIV/SIV-infected rhesus macaques (Figure 2C), although the difference did not reach statistical significance (Mann-Whitney *U* test, *P*=0.5476). Positive correlation was observed between plgR and IL-17A mRNA levels in the tracheal mucosa of normal rhesus macaques (Figure 2D), though this trend was not found in SHIV/SIV-infected rhesus macaques.

Expression of plgR in the lungs of SHIV/SIV-infected rhesus macaques

To determine whether the lungs of SHIV/SIV-infected rhesus macaques were similarly affected, the expressions of plgR mRNA and IL-17A mRNA in the lungs of normal and infected rhesus macaques were examined. The mRNA levels of plgR and IL-17A were 50 and 32 times higher, respectively, in the tracheal mucosa than in the lungs. As shown in Figure 3, plgR and IL-17A mRNA were both detected in the lungs of normal and infected rhesus macaques. In contrast to the changes observed in the tracheal mucosa, the levels of plgR and IL-17A mRNA were 3 and 1.2 times higher, respectively, in infected rhesus macaques than in normal rhesus macaques, although the differences were not statistically significant (Mann-Whitney *U* test, *P*=0.4396 and 0.7857, respectively). Therefore, the expressions of plgR and IL-17A were higher in the tracheal mucosa than in the lungs, and were not reduced in the lungs of SHIV/SIV-infected rhesus macaques.

DISCUSSION

In the present study, we observed reduced expression of plgR




Figure 1 Distribution of pIgR immunoreactivity in the epithelium of tracheal mucosa from rhesus macaques

Immunoreactivities to pIgR (green) in the pseudo-stratified columnar epithelium of tracheal mucosa from normal (left column) and SHIV/SIV-infected (right column) rhesus macaques are shown. Nuclei (blue, stained with DAPI) and DIC images indicate that histological alterations also occurred in the tracheal mucosa of SHIV/SIV-infected rhesus macaques. Original magnification, $\times 800$.

in the tracheal mucosa of SHIV/SIV-infected rhesus macaques. Both the protein levels and mRNA levels of pIgR were decreased to almost the same degree, although the decrease in protein levels was statistically significant, whereas that of mRNA was not. It is possible that the effects of SHIV/SIV infection on pIgR expression were at the gene transcription level. In consistent with these results, previous research showed that pIgR mRNA levels were significantly reduced in the intestinal mucosa of SHIV/SIV-infected rhesus macaques (Wang & Yang, 2016). Downregulation of pIgR in airway mucosa has also been documented in other airway diseases (Gohy et al., 2014; Hupin et al., 2013). Since the pathology between SIV and HIV infection is similar, pIgR expression in the airway mucosa of HIV-infected patients could also be significantly affected.

The mechanism of decreased pIgR expression in SHIV/SIV

infection has not been addressed. There are many potential factors that could affect pIgR expression, among which IL-17A can significantly regulate pIgR expression (Jaffar et al., 2009). In the present study, a decrease in IL-17A expression in the tracheal mucosa of infected rhesus macaques was observed, suggesting a role of IL-17A in the downregulation of pIgR expression in the context of SHIV/SIV infection. The non-significant difference might be due to the large individual variability and small sample size. Significant correlation between pIgR and IL-17A mRNA has been observed in the intestinal mucosa of these animals and a significant decrease in IL-17A mRNA has also been observed in the intestinal mucosa (Wang & Yang, 2016; Zhang et al., 2014). Further studies are warranted to reveal the mechanism underlying the decrease of pIgR expression in HIV/AIDS.




Figure 2 Expression levels of plgR and IL-17A in the tracheal mucosa of rhesus macaques

Levels of plgR immunoreactivity (A), plgR mRNA (B) and IL-17A mRNA (C) in the tracheal mucosa of normal and infected rhesus macaques are shown. mRNA levels of IL-17A and plgR are positively correlated (D).




Figure 3 Expression of plgR and IL-17A in the lungs of rhesus macaques

mRNA levels of plgR (A) and IL-17A (B) in the lungs of normal and infected rhesus macaques are shown. Correlation between transcription levels of plgR and IL-17A in normal rhesus macaques is also shown (C).

The consequence of reduced plgR expression in the tracheal mucosa of SHIV/SIV-infected rhesus macaques is unknown. Nevertheless, these data indicate impaired immune exclusion of potential pathogenic and commensal microbes in the respiratory

system. In line with this, increased airway microbes and pulmonary infections have been documented in HIV/SIV infections (Nimmo et al., 2015; Twigg et al., 2016). Since elevated microbes can drive the COPD-like phenotype in plgR

deficient mice (Richmond et al., 2016) and downregulation of plgR is observed in COPD patients (Gohy et al., 2014), reduced plgR expression could be an underlying mechanism of the increased incidence of COPD in HIV/AIDS patients (Morris et al., 2011). COPD is the cause of death in a significant proportion of the HIV/AIDS population. ART treatment does not decrease the incidence of COPD, but is an independent predictor of increased airway obstruction (Gingo et al., 2010). Decreased expression of plgR might also be involved in other pathological processes of HIV/AIDS, such as lung cancer (Ocak et al., 2012). Therefore, abnormal expression of plgR should be taken into consideration in novel therapies for pulmonary complications such as COPD.

In summary, for the first time, reduced plgR expression was observed in the tracheal mucosa of SHIV/SIV-infected rhesus macaques, which might be linked to IL-17A reduction in the tracheal mucosa. The reduced expression of plgR might be the underlying mechanism of increased pulmonary microbiota and infections in HIV/AIDS. Rhesus macaques are a suitable model for future dissection of the mechanisms underlying respiratory complications in HIV/AIDS.

REFERENCES

- Gingo MR, George MP, Kessinger CJ, Lucht L, Rissler B, Weinman R, Slivka WA, McMahon DK, Wenzel SE, Sciurba FC, Morris A. 2010. Pulmonary function abnormalities in HIV-infected patients during the current antiretroviral therapy era. *American Journal of Respiratory and Critical Care Medicine*, **182**(6): 790-796.
- Gohy ST, Detry BR, Lecocq M, Bouzin C, Weynand BA, Amatngalim GD, Sibille YM, Pilette C. 2014. Polymeric immunoglobulin receptor down-regulation in chronic obstructive pulmonary disease. Persistence in the cultured epithelium and role of transforming growth factor- β . *American Journal of Respiratory and Critical Care Medicine*, **190**(5): 509-521.
- Grubb JR, Moorman AC, Baker RK, Masur H. 2006. The changing spectrum of pulmonary disease in patients with HIV infection on antiretroviral therapy. *AIDS*, **20**(8): 1095-1107.
- Hupin C, Rombaux P, Bowen H, Gould H, Lecocq M, Pilette C. 2013. Downregulation of polymeric immunoglobulin receptor and secretory IgA antibodies in eosinophilic upper airway diseases. *Allergy*, **68**(12): 1589-1597.
- Jaffar Z, Ferrini ME, Herritt LA, Roberts K. 2009. Cutting edge: lung mucosal Th17-mediated responses induce polymeric Ig receptor expression by the airway epithelium and elevate secretory IgA levels. *The Journal of Immunology*, **182**(8): 4507-4511.
- Johansen FE, Kaetzel CS. 2011. Regulation of the polymeric immunoglobulin receptor and IgA transport: new advances in environmental factors that stimulate plgR expression and its role in mucosal immunity. *Mucosal Immunology*, **4**(6): 598-602.
- Johansen FE, Pekna M, Norderhaug IN, Haneberg B, Hietala MA, Krajci P, Betsholtz C, Brandtzaeg P. 1999. Absence of epithelial immunoglobulin A transport, with increased mucosal leakiness, in polymeric immunoglobulin receptor/secretory component-deficient mice. *The Journal of Experimental Medicine*, **190**(7): 915-922.
- Morris A, George MP, Crothers K, Huang L, Lucht L, Kessinger C, Kleerup EC. 2011. HIV and chronic obstructive pulmonary disease: is it worse and why? *Proceedings of the American Thoracic Society*, **8**(3): 320-325.
- Murray JF. 1996. Pulmonary complications of HIV infection. *Annual Review of Medicine*, **47**: 117-126.
- Nimmo C, Capocci S, Honeyborne I, Brown J, Sewell J, Thurston S, Johnson M, Mchugh TD, Lipman M. 2015. Airway bacteria and respiratory symptoms are common in ambulatory HIV-positive UK adults. *European Respiratory Journal*, **46**(4): 1208-1211.
- Ocak S, Pedchenko TV, Chen H, Harris FT, Qian J, Polosukhin V, Pilette C, Sibille Y, Gonzalez AL, Massion PP. 2012. Loss of polymeric immunoglobulin receptor expression is associated with lung tumorigenesis. *European Respiratory Journal*, **39**(5): 1171-1180.
- Richmond BW, Brucker RM, Han W, Du RH, Zhang YQ, Cheng DS, Gleaves L, Abdolrasulnia R, Polosukhina D, Clark PE, Bordenstein SR, Blackwell TS, Polosukhin VV. 2016. Airway bacteria drive a progressive COPD-like phenotype in mice with polymeric immunoglobulin receptor deficiency. *Nature Communications*, **7**: 11240.
- Twigg HL 3rd, Knox KS, Zhou J, Crothers KA, Nelson DE, Toh E, Day RB, Lin HY, Gao X, Dong QF, Mi DM, Katz BP, Sodergren E, Weinstock GM. 2016. Effect of advanced HIV infection on the respiratory microbiome. *American Journal of Respiratory and Critical Care Medicine*, **194**(2): 226-235.
- Wang Y, Yang GB. 2016. Alteration of polymeric immunoglobulin receptor and neonatal fc receptor expression in the gut mucosa of immunodeficiency virus-infected rhesus macaques. *Scandinavian Journal of Immunology*, **83**(4): 235-243.
- Zar HJ. 2008. Chronic lung disease in human immunodeficiency virus (HIV) infected children. *Pediatric Pulmonology*, **43**(1): 1-10.
- Zhang WJ, Wang Y, Yu K, Duan JZ, Yao WR, Wang Y, Yang RG, Yang GB. 2014. Associated changes in the transcription levels of IL-17A and tight junction-associated genes in the duodenal mucosa of rhesus macaques repeatedly exposed to simian/human immunodeficiency virus. *Experimental and Molecular Pathology*, **97**(2): 225-233.

Nest survival rate of Reeves's pheasant (*Syrnaticus reevesii*) based on artificial nest experiments

Xu Luo, Yu-Ze Zhao, Jing Ma, Jian-Qiang Li, Ji-Liang Xu*

School of Nature Conservation, Beijing Forestry University, Beijing 100083, China

ABSTRACT

To explore the nest survival rate of Reeves's pheasant (*Syrnaticus reevesii*) and the nest-site factors that affect it, we conducted artificial nest experiments with reference to natural nests at Dongzhai National Nature Reserve (DNNR), Henan Province and Pingjingguan, Hubei Province from April to June 2014 simulating the situation in its early and later breeding season. We also determined distance characteristics of the nest sites by ArcGIS 10.0. Nest survival models were constructed in Program MARK for data analysis. Results indicated that in the early breeding season, the apparent survival rate (ASR) in DNNR (52.4%) was significantly greater than that in Pingjingguan (13.5%), and the ASR in the later breeding season in DNNR (26.7%) was not indistinctly correlated with Pingjingguan (3.2%). The daily survival rate (DSR) in the later breeding season was 93.8% in DNNR and 92.0% in Pingjingguan, respectively. The DSRs were both negatively correlated with nest distance to forest edges and settlements. The DSR in Pingjingguan was positively correlated with nest distance to paths and negatively correlated with nest distance to water sources. However, the DSR in DNNR was negatively correlated with nest distance to paths but positively correlated with nest distance to water sources.

Keywords: Reeves's pheasant; *Syrnaticus reevesii*; Nest survival rate; Artificial nest experiments.

INTRODUCTION

Reeves's pheasant (*Syrnaticus reevesii*) is a rare and endangered endemic species in China. It is listed as Grade II wildlife under national protection (State Council, 1988) and as a vulnerable species by the International Union for the Conservation of Nature and Natural Resources (IUCN, 2015). Due to habitat destruction (Wu & Xu, 1987), poaching (Xu et al., 1996) and predation pressure from natural enemies (Xu et al.,

1996), the Reeves's pheasant population is decreasing at a rate of 20% every ten years (Zhou et al., 2015). The threats to Reeves's pheasant survival are considered more severe than those of many Grade I protected species in China (Zhang et al., 2003; Zhou et al., 2015). The Reeves's pheasant is ground nesting and highly vigilant, which makes it difficult to carry out field research and nest tracking (Lei & Lu, 2006; Zhang et al., 2004). Direct observation of incubating females can induce nest abandonment and breeding failure (Collar et al., 1994), and thus only limited information is known in regards to their nest survival rate.

Nest building, egg laying and egg hatching are not only critical stages in bird breeding seasons, but are also relatively precarious and easily affected by adverse environmental factors (Welty, 1962). Nest success rate is one of the most important factors determining fertility and has significant effects on bird population quantities (Beale & Monaghan, 2004). Accurate estimates of nest success rate and relevant influencing factors are important to understand the quantitative dynamics of bird populations and meaningful to enact suitable protective policies for endangered species (Lindell et al., 2011).

Artificial nest experiments using quail (*Coturnix japonica*), chicken, or artificial plaster eggs to similar to natural nests have been used to investigate predation risks of nestling (Reitsma et al., 1990; Sun et al., 2011). In addition, because it is easy to manipulate and permitting to control the experimental conditions, artificial nest experiments have also been popular in the studies on nest survival rate (Major & Kendal, 1996; Martin, 1987; Sieving, 1992; Sun et al., 2011; Willebrand & Marcström, 1988), and are a valuable reference for natural nesting survival studies (Wilson et al., 1998). Wang et al. (2016) used artificial nest experiments to study predation of the Reeves's pheasant. Results indicated that, to a certain extent, artificial nest experiments can reflect the fates of natural nests. Referencing

Received: 28 September 2016; Accepted: 10 November 2016

Foundation items: This study is supported by the Forestry Commonweal Program (201404422) and the National Natural Science Foundation of China (31172115)

*Corresponding author, E-mail: xujiliang@bjfu.edu.cn

DOI:10.13918/j.issn.2095-8137.2017.008

the methods used in Wang et al. (2016), we conducted artificial nest experiments on Reeves's pheasant at Dongzhai National Nature Reserve (DNNR), Henan Province and Pingjingguan, Suizhou City, Hubei Province from April to June 2014. Experiments were based on the characteristics of natural nests collected from field surveys in recent years, with chicken eggs used as mock eggs. We aimed to provide evidence on the nest survival rates of Reeves's pheasant, and the effects of distances from nest sites to various habitat parameters (e.g., forest edges, paths, water sources, settlements) on survival rates.

MATERIALS AND METHODS

Study area

The experiments were carried out at Baiyun Protection Station, DNNR, Henan Province (N31°28'-32°09', E114°18'-114°30') and Pingjingguan, Suizhou City, Hubei Province (N31°51'-31°52', E113°54'-113°55'). The two areas are in the south foothills of Dabieshan Mountain, less than 40 km apart and both approximately 400 hm² in area. The geological conditions, precipitation, temperature and phenology are similar (details in Wang et al., 2016).

Locating natural nests

Natural Reeves's pheasant nests in Pingjingguan were located in two ways: (1) Fourteen nests were found by radio telemetry during field surveys from 2011 to 2013, seven nests were found during this study from ten captured females caught in March 2013; and, (2) Four nests were found by interviewing local residents undertaking farming and firewood, herb collection in the mountains. The natural nests included in DNNR were seven nests found by other research teams during 2011 to 2014.

Information on the structures and sites of the fourteen natural nests obtained from Pingjingguan during 2011 to 2013 were used as references for establishing artificial nests. The twenty-five natural nests from Pingjingguan and seven from DNNR during 2011 to 2014 were included in evaluating and comparing the survival rates between natural and artificial nests.

Artificial nest experiments

The locations of the artificial nests were chosen within the areas in which Reeves's pheasants were found by radio telemetry during previous research (Sun et al., 2003; Xu et al., 2007; Bai, 2013). The selection of the artificial nests was determined based on the habitat characteristics of fourteen natural nests found during field surveys from 2011 to 2013. The artificial nests were constructed by mimicking the structures of natural nests (Wang et al., 2016). The artificial nests were disk-like, padded with withered and yellow *Castanea mollissima* leaves, *Pinus tabulaeformis* pine needles or straw, and covered with 1-2 down feathers collected from the field. Four chicken eggs of similar size and color to that of Reeves's pheasant were placed in each artificial nest.

Two rounds of artificial nest experiments were conducted. The first round was carried out from late April. When choosing locations by ArcGIS 10.0 (Esri Inc. <<http://www.esri.com/>>), seventy random locations were picked in the two study areas. Except for the locations with rocks, roads, nudations, and ponds,

one artificial nest locus was picked within a 100 m² range around each of the remaining locations. A total of 46 and 42 artificial nests were placed at Pingjingguan and DNNR, respectively. Using the same method and procedure in the second round of experiments (May 2014), fifty random locations were determined, and a total of 31 and 30 artificial nests were placed at Pingjingguan and DNNR, respectively. As per the descriptions of the four stages of breeding season in Reeves's pheasant (Sun et al., 2003), our two experiments simulated nest survival situations in the early and later stages, respectively.

In natural environments, the hatching period of Reeves's pheasant is 26-27 d (Zhang et al., 2004). In this study, the period of the artificial nest experiments was set at 30 d. Due to adverse weather conditions during the first experiment, the nests were checked three times on day 10, day 15 and day 30. During the second experiment, the nests were checked every five days. The inspecting time was defined as interval of checked and setting up day (Dinsmore et al., 2002).

To reduce researcher influence on the experiments, human disturbance was minimized during nest checking, e.g., leaving few footprints around the artificial nest and avoiding behaviors that might affect nest, such as touching the eggs (Driscoll et al., 2005). In the wild, if a nest is disturbed or eggs in the nest are preyed, the female Reeves's pheasant abandon the nests (Johnsgard, 1999). If eggs inside the artificial nest were destroyed, removed or disappeared, the eggs were not replaced (Nour et al., 1993) and the nest was defined as failed (Noske et al., 2008). Otherwise, the nest was considered one survival nest. Moreover, because Reeves's pheasant is precocial, one success nest was defined if it was a survival nest at the last inspecting time.

Distance parameters of nest locations

The GPS locations of both natural and artificial nests were analyzed by ArcGIS 10.0. By referencing the definitions of habitat parameters by Xu et al. (2006) and Wang et al. (2016), the degree of slope (Sld), distance to water resources (Dwt), paths (Dho), forest edges (Dfb) and settlements (Dro) were determined. The degree of slope was calculated by Slope tool and the minimum distance parameters were obtained by Euclidean Distance tool.

Data analysis

To compare our findings with previous reports, the apparent survival rate, i.e. ASR was defined in accordance with the definitions of Driscoll et al. (2005):

$$ASR = n_s / (n_s + n_f) \quad (1)$$

Where, ASR is the apparent survival rate, n_s represents the number of successful nests, and n_f represents the number of failed nests.

χ^2 -tests were used to compare the differences in ASRs between artificial and natural nests, as well as the differences in ASRs between the two study areas and two different study periods. The results were examined by SPSS 22.0 and plotting was conducted by Sigma Plot 12.1.

Daily survival rates, i.e. DSRs were calculated by Program MARK (V8.0). Due to the irregular checking intervals in the first

experiment, only data from the second experiment were included in calculating the DSR. In the calculation, five basic variables were used for the nest survival rate obtained by the module Nest Survival within MARK program: (1) setting up day of artificial nests (all in "1"); (2) last survival day of nest; (3) the day of nest fate (failure or success) determined; (4) the fate of nest (failure or success); and (5) quantity of nests with same fate. None of the variables went through standardization. Selecting sin option in Link Function to construct the stable model of the DSR of the artificial nests in the later breeding season, and the DSRs of the two study areas in the later breeding season were calculated.


To clarify if degree of slope and distance parameters of nest location to various habitat factors affected DSR, five variables, Sld, Dho, Dfb, Dwt and Dro, were added into the model Nest Survival. By combining the five previously described basic variables, a model regarding the correlations among DSR, degree of slope and distance parameters was constructed. The model average estimated value ($\theta \pm SE$) of each variable and 95% confidence interval, i.e. 95% CI were obtained. The plus-minus estimated value excluding "0" indicated that this variable had positive and negative effects on model construction.

In the results, data of successional variables were presented as mean $\pm SE$, where, mean is the arithmetical mean and SE is the standard deviation. A P -value of <0.05 represented significant differences.

RESULTS

Apparent survival rates between artificial and natural nests

The ASRs of the artificial nests in the two study areas decreased with time (Figure 1). At the end of the experiments, the ASRs of artificial nests in Pingjingguan and DNNR were 11.6% ($n=9$) and 41.7% ($n=30$), respectively; that of the twenty-five natural nests in Pingjingguan was 20.0%, and that of the seven natural nests in DNNR was 28.5%. The χ^2 -test results showed no significant differences in the ASRs between artificial and natural nests during the two experiments (Pingjingguan: $\chi^2=0.67$, $df=1$, $P=0.41$; DNNR: $\chi^2=0.20$, $df=1$, $P=0.65$).



Figures 1 Apparent survival rates over the two experimental periods in the two study areas

Apparent survival rates of artificial nests at the two study areas in different time periods

At the end of the early breeding season, the ASR of the artificial nests in Pingjingguan was 13.5% ($n=6$), significantly lower than that in DNNR (52.4%, $n=22$) ($\chi^2=7.65$, $df=1$, $P<0.01$). At the end of the later breeding season, the ASR of artificial nests in Pingjingguan was 3.2% ($n=1$), also lower than that of DNNR (26.7%, $n=8$), though the difference was not significant ($\chi^2=3.81$, $df=1$, $P>0.05$). In the two study areas, the ASRs of artificial nests during later breeding season were both lower than that during the early breeding season. However, significant differences were only found in DNNR (Pingjingguan: $\chi^2=4.75$, $df=1$, $P>0.05$; DNNR: $\chi^2=5.12$, $df=1$, $P<0.05$).

Daily survival rate during the later breeding season and its variations with distance and degree of slope

In the later breeding season, the stable model results showed that the DSRs in DNNR and Pingjingguan were 93.8% ($n=30$, $CI_{95}=90.8\%-95.9\%$) and 92.0% ($n=31$, $CI_{95}=88.7\%-94.3\%$), respectively.

The distance parameter analyses of model Nest Survival showed that in the later breeding season, DSRs of artificial nests in the two study areas were negatively correlated with distances to forest edges and settlements (Figure 2). In DNNR, the DSR decreased considerably with increasing distance from nest location to forest edges. Moreover, different correlation patterns of DSRs with distance to paths and water resources were found in DNNR and Pingjingguan (Figure 2). In Pingjingguan, the DSR was positively correlated with distance to paths, but negatively correlated with distance to water resources. However, these parameters effects showed the opposite pattern in DNNR.

DISCUSSION

Reeves's pheasants are highly vigilant birds, which makes it difficult to track their nests in the field and carry out monitoring (Lei & Lu, 2006; Zhang et al., 2004). Artificial nest experiments can decrease human disturbance to hatching in the field and provide sufficient samples for research purposes (Major & Kendal, 1996; Sieving, 1992; Sun et al., 2011; Wang et al., 2016; Wilson et al., 1998). In this study, the ASRs of artificial nests during the two experiments were comparable with that of natural nests, indicating their ability to mimic, to some extent, the survival situation of Reeves's pheasant nests in the field.

In the two artificial nest experiments, the ASRs during the later breeding season were both lower than those during the early breeding season. This phenomenon that nest survival rates vary or decrease with time has been reported in previous research (Daan et al., 1990). Our results also support this finding, which could relate to the increase in predation pressure and human disturbance with time (Hatchwell, 1991; Becker & Zhang, 2011).

In the two study areas, during the artificial nest experiments in late May (the later breeding season), the number of raptors, such as *Butastur indicus* and *Accipiter soloensis*, increased, and as they also began to breed, predation pressure to the




Figure 2 Daily survival rates of artificial nests in the later breeding season with different nest distances to habitat parameters

Reeves's pheasant increased as well. Therefore, predation threats to artificial nests were more severe than that during the early breeding season. According to the local DNNR survey conducted by Ma et al. (2012), because the later breeding season of Reeves's pheasant overlaps with the peak period of busy farming, e.g., pasturing, chestnut weeding, human disturbance in the later breeding season was more intense than that in the early breeding season. In Ma's survey to the community, 35.3% of interviewees claimed to collect Reeves's pheasant eggs from April to June. Sun et al. (2011) stated that repetitive artificial nest experiments in one breeding season could allow predators to adapt to human traces, with increases in predation risks in later experiments.

In this study, in the later breeding season, the DSRs in the two study areas both decreased with increasing distance from nests to the forest edges. The edge effect hypothesis indicates that the edging areas of habitats usually have more enriched vegetation resources and more complicated environments than central areas, so predation pressure at edging areas is higher (Ewers & Didham, 2007; Fahrig, 2003). Our results are discordant with the edge effect hypothesis. It is possible that because human activities are more intense in edging areas, predators, especially those that mammalian predators, are forced into the less disturbed central areas. Therefore, predation pressures in central areas are higher than that in edging areas. Moreover, because it is more common for a nest to be destroyed by a predator than by a human, the effect of

predation pressure on nest survival is higher than that of human disturbance. Wang et al. (2016) found that in the later breeding season, the major predation pressure came from predators preying on animals, which supports our assumption to a certain extent. Moreover, the negative correlation between DSR and nest distance to settlements in both study areas also suggests that human disturbance relieved potential predation pressures of artificial nests.

We also found that in the later breeding season, with increasing distance from the nests to paths, the DSR of artificial nests in Pingjingguan increased, whereas that in DNNR showed the opposite trend. The possible reasons for this are that the paths in Pingjingguan are also the main roads used by residents entering the mountain area; therefore, compared with DNNR, the habitats near paths in Pingjingguan have smaller vegetation coverage, fewer herbaceous plants, and higher human disturbance. Therefore, the nests closer to the paths were under higher predation pressure and human disturbance. Moreover, the distance of nests to water resources also had different effects on DSRs in the two study areas. In DNNR, the DSR increased with increasing distance to water resources. The reason might be that predators are more active near water resources, and thus threats from natural enemies decreased with distance to water resources and the artificial nests closer to water resources were under higher predation pressure. However, the situation in Pingjingguan exhibited the opposite pattern, and it was assumed that because water resources, e.g.,

river, penstock, were in areas with more frequent farming activities, disturbance from livestock and herdsman suppressed activities of predators nearby.

In summary, artificial nest experiments were conducted to mimic the survival rates of Reeves's pheasant's nests. Results showed low nest survival rates and high impacts of nest location. Our findings indicate that although data obtained from natural nests allow for more persuasive assumptions (Paton, 1994), to vigilant and endangered species, such as Reeves's pheasant, using artificial nests to mimic natural nests is a useful method in research.

ACKNOWLEDGEMENTS

We thank Qin-Yun Wang, Peng Zhao, and Jun-Qin Hua for help in collecting data in the fields; residents Yuan-Jun Zhang and Jin-Dong Chen for their help; and staff from the Administration Department of Dongzhai National Nature Reserve and Pingjingguan village committee for supporting this study.

REFERENCES

- Bai J. 2013. The effect of artificial plantation on home range of Reeves's pheasants (*Symptotus reevesii*). Master thesis, Beijing Forestry University. (in Chinese)
- Beale CM, Monaghan P. 2004. Behavioural responses to human disturbance: a matter of choice?. *Animal Behaviour*, **68**(5): 1065-1069.
- Becker PH, Zhang H. 2011. Renesting of Common Terns *Sterna hirundo* in the life history perspective. *Journal of Ornithology*, **152**(S1): 213-225.
- Collar NJ, Crosby MJ, Stattersfield AJ. 1994. Birds to Watch 2: The World List of Threatened Birds (BirdLife Conservation Series No. 4): The Official Source for Birds on the IUCN Red List. Cambridge, UK: Bird Life International.
- Daan S, Dijkstra C, Tinbergen JM. 1990. Family planning in the kestrel (*Falco tinnunculus*): the ultimate control of covariation of laying date and clutch size. *Behaviour*, **114**(1): 83-116.
- Dinsmore SJ, White GC, Knopf FL. 2002. Advanced techniques for modeling avian nest survival. *Ecology*, **83**(12): 3476-3488.
- Driscoll MJL, Donovan T, Mickey R, Howard A, Fleming KK. 2005. Determinants of wood thrush nest success: a multi-scale, model selection approach. *Journal of Wildlife Management*, **69**(2): 699-709.
- Ewers RM, Didham RK. 2007. The effect of fragment shape and species' sensitivity to habitat edges on animal population size. *Conservation Biology*, **21**(4): 926-936.
- Fahrig L. 2003. Effects of habitat fragmentation on biodiversity. *Annual review of ecology, evolution, and systematics*, **34**(1): 487-515.
- Hatchwell BJ. 1991. An experimental study of the effects of timing of breeding on the reproductive success of common guillemots (*Uria aalge*). *Journal of Animal Ecology*, **60**(3): 721-736.
- IUCN. 2015. IUCN red list of threatened species. Version 2014.3. http://www.iucnredlist.org/.
- Johnsgard PA. 1999. The Pheasants of the World: Biology and Natural History. 2nd ed. Washington: Smithsonian Institution Press.
- Lei FM, Lu TC. 2006. Endemic Bird Species of China. Beijing: Science Press. (in Chinese)
- Lindell CA, O'Connor RS, Cohen EB. 2011. Nesting success of neotropical thrushes in coffee and pasture. *The Wilson Journal of Ornithology*, **123**(3): 502-507.
- Ma J, Luo X, Bai J, Zhao Y, Xu JL, Li Z. 2012. The impacts from the local communities on Reeves's pheasant conservation—a preliminary case study in Dongzhai National Nature Reserve. *Forest Resources Management*, (3): 126-130. (in Chinese)
- Major RE, Kendal CE. 1996. The contribution of artificial nest experiments to understanding avian reproductive success: a review of methods and conclusions. *Ibis*, **138**(2): 298-307.
- Martin TE. 1987. Artificial Nest experiments: effects of nest appearance and type of predator. *The Condor*, **89**(4): 925-928.
- Noske RA, Fischer S, Brook BW. 2008. Artificial nest predation rates vary among habitats in the Australian monsoon tropics. *Ecological Research*, **23**(3): 519-527.
- Nour N, Matthysen E, Dhondt AA. 1993. Artificial nest predation and habitat fragmentation: different trends in bird and mammal predators. *Ecography*, **16**(2): 111-116.
- Paton PWC. 1994. The effect of edge on avian nest success: how strong is the evidence?. *Conservation Biology*, **8**(1): 17-26.
- Reitsma LR, Holmes RT, Sherry TW. 1990. Effects of removal of red squirrels, *Tamiasciurus hudsonicus*, and eastern chipmunks, *Tamias striatus*, on nest predation in a northern hardwood forest: an artificial nest experiment. *Oikos*, **57**(3): 375-380.
- Sieving KE. 1992. Nest predation and differential insular extinction among selected forest birds of central Panama. *Ecology*, **73**(6): 2310-2328.
- State Council. 1988. The list of wild animals under national protection issued by the State Council of the People's Republic of China on 10 December 1988. Beijing, China: State Council. (in Chinese)
- Sun JJ, Wang SY, Wang YP, Shao DY, Ding P. 2011. Effects of habitat fragmentation on avian nest predation risk in Thousand Island Lake, Zhejiang Province. *Biodiversity Science*, **19**(5): 528-534. (in Chinese)
- Sun QH, Zhang ZW, Zheng GM, Zhang KY, Ruan XF, Zhu J G. 2003. Ranging behaviour of territorial male Reeves's pheasants in the breeding season. *Acta Zoologica Sinica*, **49**(3): 318-324. (in Chinese)
- Wang QY, Zhao YZ, Luo X, Hua JQ, Li Z, Xu JL. 2016. Potential nest predators of *Symptotus reevesii* based on camera traps and artificial nests. *Chinese Journal of Applied Ecology*, **27**(6): 1968-1974. (in Chinese)
- Welty JC. 1962. The life of birds. *The Quarterly Review of Biology*, **85**(4): 299-417.
- Willebrand T, Marcström V. 1988. On the danger of using dummy nests to study predation. *The Auk*, **105**(2): 378-379.
- Wilson GR, Brittingham MC, Goodrich LJ. 1998. How well do artificial nests estimate success of real nests?. *The Condor*, **100**(2): 357-364.
- Wu ZK, Xu WS. 1987. On the geographical distribution and abundance of white crowned long tailed pheasants in Guizhou province. *Zoological Research*, **8**(1): 13-19. (in Chinese)
- Xu JL, Zhang ZW, Zheng GM, Zhang XH, Sun QH, McGowan P. 2007. Home range and habitat use of Reeves's Pheasant *Symptotus reevesii* in the protected areas created from forest farms in the Dabie Mountains, central China. *Bird Conservation International*, **17**(4): 319-330.
- Xu JL, Zhang XH, Zhang ZW, Zheng GM, Ruan XF, Zhu JG, Xi B. 2006. Multi-scale analysis on wintering habitat selection of Reeves's pheasant (*Symptotus reevesii*) in Dongzhai National Nature Reserve, Henan Province, China. *Acta Ecologica Sinica*, **26**(7): 2061-2067. (in Chinese)
- Xu YG, Yin ZH, Lei FM, Ding WN, Liu RS, Yu Q. 1996. The status of

Reeves's pheasant and suggestions for conservation. *Acta Zoologica Sinica*, **42**(S): 155. (in Chinese)

Zhang XH, Xu JL, Zhang ZW, Xie FL, Zhang KY, Zhu JG. 2004. A Study on the incubation behavior of Reeves's pheasant (*Symaticus reevesii*) by radio tracking. *Journal of Beijing Normal University: Natural Science*, **40**(2): 255-259. (in Chinese)

Zhang ZW, Ding CQ, Ding P, Zheng GM. 2003. The current status and a conservation strategy for species of Galliformes in China. *Chinese Biodiversity*, **11**(5): 414-421.

Zhou CF, Xu JL, Zhang ZW. 2015. Dramatic decline of the vulnerable Reeves's pheasant *Symaticus reevesii*, endemic to central China. *Oryx*, **49**(3): 529-534.

Zoological Research Editorial Board

EDITOR-IN-CHIEF

Yong-Gang Yao

Kunming Institute of Zoology, CAS, China

ASSOCIATE EDITORS-IN-CHIEF

Wai-Yee Chan

The Chinese University of Hong Kong, China

Xue-Long Jiang

Kunming Institute of Zoology, CAS, China

Bing-Yu Mao

Kunming Institute of Zoology, CAS, China

Yun Zhang

Kunming Institute of Zoology, CAS, China

Yong-Tang Zheng

Kunming Institute of Zoology, CAS, China

MEMBERS

Le Ann Blomberg

Beltsville Agricultural Research Center, USA

Jing Che

Kunming Institute of Zoology, CAS, China

Biao Chen

Capital Medical University, China

Ce-Shi Chen

Kunming Institute of Zoology, CAS, China

Gong Chen

Pennsylvania State University, USA

Jiong Chen

Ningbo University, China

Xiao-Yong Chen

Kunming Institute of Zoology, CAS, China

Michael H. Ferkin

University of Memphis, USA

Nigel W. Fraser

University of Pennsylvania, USA

Colin P. Groves

Australian National University, Australia

Wen-Zhe Ho

Wuhan University, China

David Irwin

University of Toronto, Canada

Nina G. Jablonski

Pennsylvania State University, USA

Prithwiraj Jha

Raiganj Surendranath Mahavidyalaya, India

Xiang Ji

Nanjing Normal University, China

Le Kang

Institute of Zoology, CAS, China

Ren Lai

Kunming Institute of Zoology, CAS, China

Bin Liang

Kunming Institute of Zoology, CAS, China

Wei Liang

Hainan Normal University, China

Hua-Xin (Larry) Liao

Duke University, USA

Si-Min Lin

Taiwan Normal University, China

Huan-Zhang Liu

Institute of Hydrobiology, CAS, China

Meng-Ji Lu

University Hospital Essen, University Duisburg Essen, Germany

Masaharu Motokawa

Kyoto University Museum, Japan

Victor Benno Meyer-Rochow

University of Oulu, Finland

Monica Mwale

South African Institute for Aquatic Biodiversity, South Africa

Neena Singla

Punjab Agricultural University, India

Bing Su

Kunming Institute of Zoology, CAS, China

Wen Wang

Kunming Institute of Zoology, CAS, China

Fu-Wen Wei

Institute of Zoology, CAS, China

Jian-Fan Wen

Kunming Institute of Zoology, CAS, China

Richard Winterbottom

Royal Ontario Museum, Canada

Jun-Hong Xia

Sun Yat-sen University, China

Lin Xu

Kunming Institute of Zoology, CAS, China

Jian Yang

Columbia University, USA

Xiao-Jun Yang

Kunming Institute of Zoology, CAS, China

Hong-Shi Yu

University of Melbourne, Australia

Li Yu

Yunnan University, China

Lin Zeng

Academy of Military Medical Science, China

Xiao-Mao Zeng

Chengdu Institute of Biology, CAS, China

Guo-Jie Zhang

University of Copenhagen, Denmark

Ya-Ping Zhang

Chinese Academy of Sciences, China

ZOOLOGICAL RESEARCH
动物学研究
Bimonthly, Since 1980



Editor-in-Chief: Yong-Gang Yao

Executive Editor-in-Chief: Wai-Yee Chan

Editors: Su-Qing Liu Long Nie

Edited by Editorial Office of Zoological Research

(Kunming Institute of Zoology, Chinese Academy of Sciences, 32 Jiaochang Donglu, Kunming,
Yunnan, Post Code: 650223 Tel: +86 871 65199026 E-mail: zoores@mail.kiz.ac.cn)

Sponsored by Kunming Institute of Zoology, Chinese Academy of Sciences; China Zoological Society©

Supervised by Chinese Academy of Sciences

Published by Science Press (16 Donghuangchenggen Beijie, Beijing 100717, China)

Printed by Kunming Xiaosong Plate Making & Printing Co, Ltd

Domestic distribution by Yunnan Post and all local post offices in China

International distribution by China International Book Trading Corporation (Guoji Shudian) P.O.BOX 399,
Beijing 100044, China

Advertising Business License 广告经营许可证: 滇工商广字66号

Domestic Postal Issue No.: 64-20

Price: 10.00 USD/60.00 CNY Post No.: BM358



ISSN 2095-8137

